

# EFFECT OF BIOSTIMULANTS ON THE GROWTH ATTRIBUTES AND YIELD OF RICE

---

## ABSTRACT

A field investigation was conducted at Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu during September - February 2022 to study the effect of biostimulants on the growth attributes and yield of rice. The experiment was laid out in a randomized block design with nine treatments and three replications. The treatments consisted of T<sub>1</sub> - 100% RDF (150:50:50 kg of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O), T<sub>2</sub> - 100% RDF + 3% Panchagavya spray at 30, 45 and 60 DAT, T<sub>3</sub> - 100% RDF + 5% Jeevamirtham spray at 30, 45 and 60 DAT, T<sub>4</sub> - 100% RDF + 3% Dasagavya spray at 30, 45 and 60 DAT, T<sub>5</sub> - 100% RDF + 2% Humic acid spray at 30, 45 and 60 DAT, T<sub>6</sub> - 100% RDF + 5% Seaweed liquid fertilizer (*Sargassum wightii*) spray at 30, 45 and 60 DAT, T<sub>7</sub> - 100% RDF + 2% Effective microorganism inoculation spray at 30, 45 and 60 DAT, T<sub>8</sub> - 100% RDF + 1% Liquid biofertilizer spray at 30, 45 and 60 DAT and T<sub>9</sub> - 100% RDF + 5% Seaweed liquid fertilizer (*Ascophyllum nodosum*) spray at 30, 45 and 60 DAT. Among these treatments, it is observed that the application of a recommended dose of fertilizers along with 2 per cent humic acid spray at 30, 45 and 60 DAT had a remarkable effect on the resulted in enhanced values of growth attributes and yield viz., plant height (95.6 cm), number of tillers hill<sup>-1</sup> (25.01), leaf area index (5.43), dry matter production (12927 kg ha<sup>-1</sup>), root length (24.63 cm), root volume (34.73 cc), grain yield (5540 kg ha<sup>-1</sup>) and straw yield (8200 kg ha<sup>-1</sup>) of rice.

**Keywords:** Growth, yield, rice, biostimulants and humic acid

## 1. INTRODUCTION

Agriculture has a critical role in the Indian economy. It serves as its foundation and is acknowledged as the most common employment globally. Rice (*Oryza sativa* L.) is one of the world's most significant staple food grain crops, both in terms of farmed area and consumption levels. Rice is the most common ancient crop, growing in 117 countries and accounting for over 90% of global rice production and consumption in Asia [1]. Rice is cultivated in 114 countries across the world with an area of 165.25 million hectares with a production of 505.4 million tonnes and average productivity is 4.66 t ha<sup>-1</sup>[2]. In India, rice is grown on approximately 44.6 million hectares, with a production of around 130.5 million tonnes and an average productivity of 2927 kg ha<sup>-1</sup>[3].

Rice requires a large amount of nutrients to get its maximum production. Continuous and indiscriminate use of inorganic fertilizers in intensive cropping systems causes nutrient imbalances in the soil as well as environmental contamination, which has a negative impact on soil health, ecological dangers, and crop yields [4] and [5]. To maintain current food security and fulfill future food demands, innovative management strategies must be devised and tested to boost production. The use of biostimulants in modest doses improves physiological processes in plants, resulting in high-quality product [6]. They improve plants metabolic and enzymatic activities, particularly during the early stages of crop development. They also improve nutrient absorption, making plants more resistant to stress and increasing output. As a result, these biostimulants play a vital role in plant structural processes that promote growth, tolerance to abiotic and biotic stress, and increased product yield and quality. The use of biostimulants in agriculture is a sustainable approach that boosts output while minimizing environmental impact [7]. As a result, the current study was designed to investigate the influence of biostimulants on rice growth and yield.

## 2. MATERIALS AND METHODS

The field experiment was conducted at Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu during September - February, 2022 to study the effect of biostimulants on the growth and yield of rice (Fig. 1). The experimental farm is geographically located at 11° 24' N latitude and 79° 44' E longitude, at an altitude of +5.79 m above MSL. The soil of the experimental field was clay loam in texture. The initial nutrient status of the experimental field soil was low in available nitrogen, medium in available phosphorus and high in available potassium. The field experiment was laid out in a randomized block design with three replications. The treatments consisted of T<sub>1</sub> - 100% RDF (150:50:50 kg of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O), T<sub>2</sub> - 100% RDF + 3% Panchagavya spray at 30, 45 and 60 DAT, T<sub>3</sub> - 100% RDF + 5% Jeevamirtham spray at 30, 45 and 60 DAT, T<sub>4</sub> - 100% RDF + 3% Dasagavya spray at 30, 45 and 60 DAT, T<sub>5</sub> - 100% RDF + 2% Humic acid spray at 30, 45 and 60 DAT, T<sub>6</sub> - 100% RDF + 5% Seaweed liquid fertilizer (*Sargassum wightii*) spray at 30, 45 and 60 DAT, T<sub>7</sub> - 100% RDF + 2% Effective microorganism inoculation spray at 30, 45 and 60 DAT, T<sub>8</sub> - 100% RDF + 1% Liquid biofertilizer spray at 30, 45 and 60 DAT and T<sub>9</sub> - 100% RDF + 5% Seaweed liquid fertilizer (*Ascophyllum nodosum*) spray at 30, 45 and 60 DAT. BPT-5204 was the variety chosen for the experiment, and 30 kg of seed was sown per hectare at a spacing of 20 × 10 cm in a field that had been levelled and well-puddled. The foliar application of 3% Panchagavya, 5% Jeevamurtha, 3% Dasagavya, 2% Humic acid, 5% Seaweed liquid fertilizer, 2% Effective Microorganism inoculation and 1% Liquid Biofertilizer were sprayed as per the treatment schedule at 30, 45, 60 days after transplanting. Two hand weeding were done on 20 DAT and 45 DAT. The quantitative assessment of growth parameters and yield were recorded at respective stages of crop growth. The data collected from the field experiment was statistically analyzed [8].



**Fig. 1 – Overview of the experimental field**

### **3. RESULT AND DISCUSSION**

#### **3.1 Growth attributes**

The results of the field experiment involving inorganic fertilizers with biostimulants exhibit an impact on the growth and growth parameters on rice are furnished in **Table 1.**

Among the various treatments, 100 per cent recommended dose of fertilizer and 2 per cent humic acid spray at 30, 45 and 60 DAT ( $T_5$ ) performed the highest growth attributes *viz.*, plant height (95.6 cm), number of tillers hill<sup>-1</sup> (25.01), leaf area index (5.43), dry matter production (12927 kg ha<sup>-1</sup>), root length (24.63 cm), root volume (34.73 cc) than rest of the treatments. It was followed by the application of a recommended dose of fertilizer along with Seaweed liquid fertilizer (*Sargassum wightii*) 5 per cent spray at 30, 45 and 60 DAT ( $T_6$ ). The increased plant height could be attributed to an adequate supply of inorganic and humic acid to the plant, which resulted in rapid growth through good root establishment and various metabolic processes, ultimately performing

**Table 1. Effect of biostimulants on the growth attributes of rice crop at harvest**

Treatments	Plant height (cm)	Number of tillers hill <sup>-1</sup> (At maximum tillering stage)	Leaf area index	Dry matter production (kg ha <sup>-1</sup> )	Root length (cm) (At flowering)	Root volume (cc) (At flowering)
T <sub>1</sub> - 100% RDF	82.1	13.10	3.47	8153	17.47	26.33
T <sub>2</sub> - 100% RDF + 3% Panchagavya spray at 30,45 and 60 DAT	86.3	15.60	4.31	9526	19.70	27.73
T <sub>3</sub> - 100% RDF + 5% Jeevamurtha spray at 30,45 and 60 DAT	86.6	15.50	4.29	9489	18.63	29.23
T <sub>4</sub> - 100% RDF + 3% Dasagavya spray at 30,45 and 60 DAT	88.8	15.33	4.18	9390	20.80	30.87
T <sub>5</sub> - 100% RDF + 2% Humic acid spray at 30,45 and 60 DAT	95.6	25.01	5.43	12927	24.63	34.73
T <sub>6</sub> - 100% RDF + 5% Seaweed liquid fertilizer ( <i>Sargassum wightii</i> ) spray at 30,45 and 60 DAT	90.3	22.90	5.16	11120	24.00	34.47
T <sub>7</sub> - 100% RDF + 2% Effective microorganism inoculation spray at 30,45 and 60 DAT	87.6	19.02	4.73	10788	22.17	33.40
T <sub>8</sub> - 100% RDF + 1% Liquid biofertilizer spray at 30,45 and 60 DAT	88.7	17.02	4.47	9820	21.67	30.47
T <sub>9</sub> - 100% RDF + 5% Seaweed liquid fertilizer ( <i>Ascophyllum nodosum</i> ) spray at 30,45 and 60 DAT	90.6	22.01	5.13	11160	23.33	33.27
<b>S.Ed</b>	<b>0.83</b>	<b>0.55</b>	<b>0.07</b>	<b>110.98</b>	<b>0.87</b>	<b>0.49</b>
<b>CD (P=0.05)</b>	<b>1.76</b>	<b>1.17</b>	<b>0.15</b>	<b>235.29</b>	<b>1.88</b>	<b>1.04</b>

better mobilization of synthesized carbohydrates into amino acids and protein, stimulating rapid cell division and cell elongation. Humic acid's auxin action enhances plant membrane permeability and intensifies enzyme systems, resulting in enhanced plant height and crop dry matter. These results are presented with the findings of [9] and [10].

The number of tillers hill<sup>-1</sup> and LAI also significantly increased with the combination of 100 per cent RDF along with 2 per cent humic acid at 30, 45 and 60 DAT (T<sub>5</sub>). The increased number of tillers might be attributed to the rapid availability of nutrients through inorganics in the latter phases of crop growth, as well as the delayed release of nitrogen and consistent release of other nutrients over a prolonged period of crop growth by organics. Productive tillers naturally stabilize as crop age increases. Humic acid improves nitrogen availability, which is required for vegetative development, leading in an increase in the number of tillers hill<sup>-1</sup> and maximum LAI. The results are similar to the findings of [11]; [12] and [13].

### 3.2 Grain and straw yield

All the treatments significantly influenced the grain and straw yield of rice are presented in **Table 2**. The treatment 100 per cent RDF + 2 per cent humic acid spray at 30, 45 and 60 DAT significantly **increased the grain yield** (5540 kg ha<sup>-1</sup>) and straw yield (8200 kg ha<sup>-1</sup>) of rice (**Fig. 2**). This might be attributed to **the rapid absorption** of more nitrogen, phosphorous, potassium, and micronutrients found in inorganic fertilizers and organic compounds. This leads to improved physiological and morphological characteristics, which are then reflected in better yield.

Similarly, higher growth metrics such as plant height, the number of tillers per hill, the leaf area index, and dry matter production resulted in the maximum straw output. This results in similar findings of [14] and [15].



**Fig. 2 - Best Treatment (T<sub>5</sub>) – 100% RDF + 2 per cent Humic acid spray at 30, 45 and 60 DAT) & Least Treatment (T<sub>1</sub>) – Application of 100% RDF alone (150:50:50 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O)**

**Table 2. Effect of biostimulants on the grain yield and straw yield of rice crop at harvest**

Treatments	Grain yield (kg ha <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )
T <sub>1</sub> - 100% RDF	4150	6990
T <sub>2</sub> - 100% RDF + 3% Panchagavya spray at 30,45 and 60 DAT	4490	7220
T <sub>3</sub> - 100% RDF + 5% Jeevamurtha spray at 30,45 and 60 DAT	4500	7280
T <sub>4</sub> - 100% RDF + 3% Dasagavya spray at 30,45 and 60 DAT	4450	7210
T <sub>5</sub> - 100% RDF + 2% Humic acid spray at 30,45 and 60 DAT	5540	8200
T <sub>6</sub> - 100% RDF + 5% Seaweed liquid fertilizer ( <i>Sargassum wightii</i> ) spray at 30,45 and 60 DAT	5240	7820
T <sub>7</sub> - 100% RDF + 2% Effective microorganism inoculation spray at 30,45 and 60 DAT	4820	7610
T <sub>8</sub> - 100% RDF + 1% Liquid biofertilizer spray at 30,45 and 60 DAT	4630	7400
T <sub>9</sub> - 100% RDF + 5% Seaweed liquid fertilizer ( <i>Ascophyllum nodosum</i> ) spray at 30,45 and 60 DAT	5150	7840
<b>S.Ed</b>	<b>58.00</b>	<b>89.79</b>
<b>CD (P=0.05)</b>	<b>122.95</b>	<b>190.36</b>

## 4. CONCLUSION

Based on the result of the field experiment, it is observed that the application of a recommended dose of fertilizers along with 2 per cent humic acid spray at 30, 45 and 60 DAT had a remarkable effect on the growth attributes and yield of rice. It is an effective practice for augmenting higher yield and has also been found to be agronomically feasible and economically viable. So, this practice can be recommended to the rice cultivating farmers of Tamil Nadu.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text to image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Debbarma VICTOR, Abraham V, Abraham T, Debbarma S, Debbarma H. Influence of different planting methods and organic nutrients on growth and yield of rice (*Oryza sativa* L. sub sp. *Japonica*). The Bioscan. 2015;9:1039-1044.
2. USDA. World Agricultural Production. United States Department of Agriculture, Foreign Agricultural Service. Washington, D.C. 2022.
3. Ministry of Agriculture & Farmers Welfare, Government of India. Second Advance Estimates of Production of Foodgrains for 2023-24. New Delhi, India.
4. Sivanantha J, Sunil JA. The paddy cultivation in the era of corona virus (COVID-19) pandemic crises and its prompts changes in paddy cultivation. In Book: Latest innovation for future education (LIFE-2021), Agriculture and Food, ESN Publications, 2021;199-208.
5. Sivanantha J, Durga CS, Suriya SG. A Review on Environmental Impact of Pesticide Usage and Its Agronomic Management Strategies. Trends in Agricul. Sci. 2024;3(Spl1):102-105.DOI:10.5281/zenodo.10551188
6. Vasconcelos ACF, Chaves LHG. Biostimulants and their role in improving plant growth under abiotic stresses. In Book: Biostimulants in plant science. 2019;3-16.
7. Rodrigues M, Baptistella JLC, Horz DC, Bortolato LM, Mazzafera P. Organic plant biostimulants and fruit quality-A review. Agron. 2020;10(7):988.
8. Gomez KA, Gomez AA. Statistical procedure for agricultural research.2nd Edition, John Wiley and Sons, New York, 1984;680.
9. Eshwar M, Srilatha M, Bhanu Rekha K, Harish Kumar Sharma S. Effect of humic substances (humic, fulvic acid) and chemical fertilizers on nutrient uptake, dry matter production of aerobic rice (*Oryza sativa* L.). J. Pharma. Phytochem. 2017;6(5):1063-1066.

10. Rajnesh T, Singh NB, Singh V, Kumar D. Effect of planting methods and integrated nutrient management on growth parameters, yield and economics of rice. *J. Pharma. Phytochem.* 2018;7(2):520-527.
11. Osman EAM, El-Masry AA, Khatab KA. Effect of nitrogen fertilizer sources and foliar spray of humic and / or fulvic acids on yield and quality of rice plants. *Adv. Appl. Sci. Res.* 2013;4(4):174-183.
12. Rasool G, Wahla AJ, Nawaz M, Rehman MA. Determination and evaluation of the effect of different doses of humic acid on the growth and yield of wheat (*Triticum aestivum* L.). *J. Agri. Veter. Sci.* 2015;8(2):5-7.
13. Balamurugan R, Babu R, Swaminathan C, Baskar K, Gopal, NO. Performance of organic nutrition on physiological characters and yield of aerobic rice. *Int. J Cur. Micro. Appl. Sci.* 2018;6:2936-2947.
14. Priyanka B, Ramesh T, Rathika S, Balasubramaniam P. Foliar application of fish amino acid and egg amino acid to improve the physiological parameters of rice. *Int. J. Curr. Microbio. Appl. Sci.* 2019;8(2):3005-3009.
15. Vinod SR, Jawahar S, Kowsalya M, Kumar CS. Effect of integrated nutrient management practices on productivity and profitability of transplanted rice. *Plant Archives.* 2019;19(1):1219-1222.