

## Minireview Article

### Ectopic tooth eruption – Integrative Literature Review

**Abstract:** Tooth eruption is characterized by the axial movement of the tooth element, from the position of formation in the alveolar bone to its place of function in the occlusal plane. The term ectopic eruption is used to designate teeth that deviate from their normal eruption pattern. Its etiology is multifactorial including local and systemic factors, somatic disorders, and genetics. To discuss the main etiological and clinical management aspects of ectopic teeth. A literature search of scientific articles indexed in the PUBMED® and VHL databases was carried out in September 2023, with the following descriptors: "Tooth Eruption; Ectopic; Dentistry". 42 articles published in the last 5 years were found. Of these, 16 addressed ectopic eruption and were included. The reviewed literature highlights the following clinical aspects: delayed eruption, partial appearance of the crown, changes in the eruption axis, transpositions and gyro versions. It is essential to diagnose ectopic eruption early and identify etiological factors, in order to obtain an early treatment, to prevent the development of this pathology and, consequently, the onset of malocclusion.

**Key words:** Tooth Eruption; Ectopic; Dentistry.

#### 1. INTRODUCTION

Tooth eruption is a phenomenon characterized by the axial movement of the tooth from its position of development in the alveolar bone until it reaches its functional site within the dental arches [1]. When there is a deviation from the normal tooth eruption pattern, and one or more teeth emerge out of anatomical position, ectopic eruption occurs [2].

The interaction between the oral epithelium and germ cells plays important roles in tooth development [2]. Developmental disorders, pathologies, cellular processes with

mutations or iatrogenic activity during embryonic development can cause an abnormal relationship or migration of these cells to an atypical position [2,3].

In this context, the pathogenesis of ectopic eruption may be directly related to the processes of odontogenesis [2,3]. However, the etiology is multifactorial and not yet well understood [1,4]. Somatic disorders and genetic factors such as hypothyroidism, Down Syndrome and Cleido Cranial Dysplasia are described as predisposing [1,4].

In addition, local factors such as supernumerary teeth, atresia and clefts in the bone bases, discrepant tooth morphology and sizes, and prolonged retention of deciduous teeth are conditions commonly associated with ectopic eruption [1,4,18].


Epidemiology establishes that the ectopic eruption most often affects the first permanent molars and canines, both in the mandible and maxilla [1,5]. In other regions outside the oral cavity this condition is rare, although there are reports of teeth in the maxillary sinus, mandibular condyle, coronoid process, palate, chin and nasal cavity [1,5,19,20].

Considering the functional and aesthetic impacts related to ectopic eruption, the objective of this study was, based on an integrative review of the literature, to discuss the main etiological and clinical management aspects of ectopic teeth.

## **2. METHODOLOGY**

A bibliographic search of scientific articles indexed in the PUBMED® and VHL databases (Virtual Health Library), carried out in July 2024, was carried out with the following descriptors: "Tooth Eruption; Ectopic; Dentistry". Articles published in the last 5 years were included as potentially relevant. A total of 30 were identified on PUBMED® and 12 on VHL, totaling 42 articles. After the analysis of all articles, 17 were excluded by title, 7 were excluded after reading the abstract and 2 were removed because they were duplicates, so 16 articles were selected for full reading and all were included in this study. The exclusion criteria are articles that have been published for more than 5 years, articles for which it was not possible to obtain the full text, articles with subjects that blur their theme in relation to the central theme.

**Figure 1:** Flowchart of the selection process of the included articles



**Table 1:** Characteristics of the included studies

Article name	Authors/references	Incidence/prevalence	Tooth	Study type	Summary of the article
--------------	--------------------	----------------------	-------	------------	------------------------

<b>Treatment of ectopic eruption of the maxillary first permanent molar in children and adolescents: A scoping review</b>	[1]	The prevalence of ectopic eruption is up to 6% and affects both sexes equally.	1st molar	Scoping review	To conduct a scoping review to collect the most relevant studies published in the last 30 years regarding ectopic eruption of the first permanent molar. Its local eruptive behavior is disturbed during the development of mixed dentition.
<b>Ectopic Tooth in Mandibular Canal, Maxillary Sinus, and Mandibular Condyle</b>	[17]	Uncertain	Teeth in general	Case description	To describe the ectopic eruption, which although rare, there have been cases in the maxillary sinus, mandibular condyle, nasal cavity, chin, palate and orbital floor. It has been reported that odontogenic tumors and cysts can develop around the ectopic tooth. Therefore, ectopic teeth can be monitored regularly if they do not present signs and symptoms.
<b>Management of rare ectopic teeth eruption: case series</b>	[6]	80.0% male, with an average of 23.3 years of age.	3rd molar	Case series	To report a case series of ectopic teeth in rare locations in the mandible and highlight the associated pathology and experience in surgical management. Of these, ten cases of ectopic teeth were identified during the study period. The antrum and lower border of the mandible represented 50.0% and 40.0% of ectopic locations, respectively.
<b>Laser-assisted management of ectopic eruption of permanent incisors</b>	[7]	The most frequently influenced teeth are the upper first molars, accompanied by the upper canines.	Incisives	Case report	The use of Er, Cr: YSGG lasers for correction of hyperplastic soft tissues involving ectopically erupting permanent incisors is portrayed considering two cases.
<b>Ectopic tooth in maxillary sinus compressing the nasolacrimal canal: A case report</b>	[2]	Uncertain	Tooth with dentigerous cyst	Case report	To present a rare case of an ectopic tooth with a dentigerous cyst in the maxillary sinus compressing the nasolacrimal duct.
<b>Ectopic Eruption of First Permanent Molars for Pediatric Patients Attended King Saud University.</b>	[8]	Male 28 (2.9%), female 17 (1.6%). 47 (78.33%) maxilla, 13 (21.66%) mandible	1st molar	Case report	Study aimed to determine the prevalence of ectopic eruption of the upper first permanent molar in the pediatric population and analyze whether there is a relationship between this anomaly and dento-skeletal characteristics using Ricketts cephalometry.

<b>Ectopic Eruption of Maxillary First Permanent Molars: Preliminary Results of Prevalence and Dentoskeletal Characteristics in Spanish Pediatric Population</b>	[5]	Prevalence of maxillary ectopic eruption (MEE), 8.7%. Age group 7 years of age (18.8%) and bilateral.	1st molar	Prevalence study (Spain)	To determine the prevalence of EEM in children and analyze whether there is a relationship between EEM and dento-skeletal characteristics. 322 children were analyzed using the Ricketts cephalometric study and descriptive and analytical statistical analysis were performed.
<b>Ectopic Permanent Molars: A Review</b>	[9]	7 to 8 years old	Molars	Literature review	Elucidate the etiology, clinical characteristics and consequences of ectopic molar eruption, in addition to highlighting the importance of future research on the variables that contribute to irreversible results of ectopic molars.
<b>Ectopically Positioned Tooth Piercing the Philtrum</b>	[10]	Uncertain	Incisive	Case report	Case report of a 70-year-old woman who presented to the Department of Oral and Maxillofacial Surgery, Karnavati School of Dentistry, Uvardsad, Gandhinagar, Gujarat, India reporting an abnormally positioned tooth piercing her upper lip. According to the patient's history, the labially erupted tooth had been perforating the philtrum for 60 years, it was visible extra orally and was painful.
<b>Ectopic Mandibular Canine with Partial Anodontia - A Rare Case Report</b>	[11]	Rare case	Canine	Rare case report	To present a rare case of non-syndromic partial anodontia and ectopic eruption of the permanent lower canine in a female patient.
<b>Evaluation of the position of the permanent maxillary canine and its relationship to dental anomaly patterns in the pediatric patient</b>	[12]	The prevalence of ectopic eruption of the permanent upper canine was 9.23%.	Superior canine	Descriptive/transversal/observational and descriptive	To evaluate the prevalence of ectopic eruption of the permanent upper canine in patients between 6 to 10 years of age and its relationship with other dental anomalies, age and sex of the patient.
<b>Management of ectopically erupting permanent molars in a seven-year-old girl: A case report</b>	[13]	Uncertain	Permanent molar	Case report	To report the case of a seven and a half year old girl with ectopic eruption of the permanent upper and lower first molars, treated with orthodontic separators and cutting the distal surface of the adjacent deciduous second molar separately. Regular examination and timely interception are important during the management of children's oral health and are of great importance in alleviating the side effects of ectopic eruption.

<p><b>Case Report: Ectopic third molar in the maxillary sinus with infected dentigerous cyst assessed by cone beam CT</b></p>	<p>[14]</p>	<p>Uncertain</p>	<p>3rd molar/case report</p>	<p>Case report</p>	<p>Report the case of a 13-year-old student who presented with purulent secretion behind her right upper second molar with a bad taste and foul odor. Radiographic examination revealed a third upper molar located on the posterosuperior surface of the right maxillary sinus with a hyperdense lesion surrounding the crown, obliterating the sinus cavity. Both the tooth and the dentigerous cyst were surgically removed under general anesthesia via a Caldwell-Luc antrostomy.</p>
<p><b>Canine Transposition - Prevalence, Distribution and Treatment Considerations among Orthodontic Patients</b></p>	<p>[15]</p>	<p>3 (20%) men and 12 (80%) women with a prevalence of 0.5%.</p>	<p>Canine</p>	<p>Prevalence /cross-sectional</p>	<p>To determine the prevalence and distribution of canine transposition in a sample of orthodontic patients and present treatment alternatives and outcomes.  Study design: The records of 3,000 consecutive orthodontic patients treated at a university clinic were searched to detect transposition of canines in both dental arches.</p>
<p><b>Corrective Treatment of Ectopic Eruption of Permanent First Molars: Case Report / Tratamiento correctivo de erupción ectópica de primeros molares permanentes: Reporte de casos</b></p>	<p>[3]</p>	<p>Uncertain</p>	<p>1st molar</p>	<p>Case report</p>	<p>The objective of the present study was to report two clinical cases where ectopic eruption of first permanent molars was corrected using elastic separators.</p>
<p><b>Tooth transposition: a multidisciplinary approach</b></p>	<p>[16]</p>	<p>Uncertain</p>	<p>Teeth in general</p>	<p>Review with case report</p>	<p>Discuss the treatment modalities for dental transposition, their advantages and disadvantages, and present a clinical case of transposition between canine and lateral incisor in the upper right hemiarch. The therapeutic choice was the extraction of one of the transposed teeth, requiring a multidisciplinary approach with the participation of Orthodontics, Dentistry and Periodontics, so that adequate aesthetic and functional results were achieved.</p>

### 3. RESULT AND DISCUSSION

A variety of eruption disorders can occur during the period of intraosseous tooth formation up to the occlusion position. This change is more frequent during the transitional period, in mixed dentures, in which the deciduous dentition is replaced by the permanent one [16].

An ectopic tooth is one that is displaced at a considerable distance from its normal anatomical location [6]. One of the most common types of ectopic eruption is tooth transposition, which results in a change in the position of the teeth and an abnormal sequence of permanent teeth in the dental arch [15].

Transposition can be complete when all the crowns and root structures of the teeth involved are in transposed malposition, or incomplete (also described as "pseudo" or "partial") when only the crowns are transposed, but the root apices remain in normal positions [15]. Transposition has been reported to occur most frequently between the maxillary permanent canine and the first premolar, and occasionally involving the maxillary canine and lateral incisor or the mandibular canine and lateral incisor. This anomaly is more often found unilaterally than bilaterally, with a predominance of the left side and more often in women than in men [15] (Figure 1).

The prevalence of eruption abnormality has been reported to be between 1.8% to 6%, affecting both sexes equally. It is believed that there is a familial tendency (higher occurrence in siblings) [1]. It is also evidenced as an expression of numerous syndromes and in children who present concomitant orofacial and dental anomalies, such as cleft palate (with a four-times higher prevalence), congenital absence of teeth, supernumerary teeth and dental size anomalies [1].

The teeth most affected by ectopy are the first permanent molars, comprising 2 to 6%, and the canines with approximately 1.5 to 4.6%. Ectopy is identified in the incisors, second molars, premolars and, more uncommonly, in the third molars (these are more frequently reported as impaction) [1,9,15].

The etiology of an ectopic eruption is still uncertain and may occur due to local and systemic factors. Several theories have been presented to describe the appearance of this condition, such as: developmental disorders like cleft palate; pathological processes such as large cysts, which displace the tooth buds to other areas; odontogenic and rhinogenic infections; or iatrogenesis [12,14].

In the oral cavity, a clinical variety of inclinations, angulations and transpositions are observed. And it can be present even in other regions of the body such as the maxillary sinus, condylar process, coronoid process, nasal cavity and orbit [6].

The atypical rash is usually a painless abnormality and commonly not noticed by patients. However, it may be associated with recurrent infections characterized by pain, swelling, and purulent discharge, dentigerous cysts, bone expansion, or may be discovered incidentally during other investigations [6].

The diagnosis is essentially clinical and radiographic. Evaluation is done through panoramic radiographs and teleradiography, or in more complex cases computed tomography. Clinically, more frequent changes in position are observed in the initial mixed dentition from 6 to 7 years of age. This local condition is grouped into self-correcting or reversible when the ectopic tooth eventually goes into occlusion after the end of its eruption; or irreversible when the ectopic element remains in an altered position even after the irruptive period [1,14].

In this context and to avoid functional and aesthetic problems as a result of this condition, an early diagnosis is necessary. The management sequence includes careful anamnesis, clinical and radiographic diagnosis, and appropriate treatment. The individual tooth position should be evaluated before the irruptive process and its intermaxillary relations in the three-dimensional context of the maxilla and mandible [1,6,12].

Preventive treatment should focus on three objectives: favoring tooth eruption in the normal position, maintenance of the irruptive sequence, and arch perimeter. These measurements can create space and bone direction so that the teeth can occupy their correct position in the dental arch. In this sense, the planning can include the use of orthopedic appliances for the expansion of the dental arches, as well as serial and planned extractions for the correct alignment and leveling of the teeth at the bone bases [17].

As a consequence of this process and the lack of early diagnosis and treatment, there is a tendency for prolonged retention or early loss of deciduous teeth, root resorption, caries lesions in adjacent teeth, migration of neighboring teeth, referred pain, alignment problems and alteration in normal occlusion [1,12,15].

Therefore, eruption guidance during deciduous and mixed dentitions is an integral part of comprehensive oral health care to achieve occlusal harmony, function, and aesthetics [1,12,15].

#### 4. CONCLUSION

The orientation of the eruption during deciduous and mixed dentitions is an integral part of comprehensive oral health care to achieve occlusal harmony, function, and aesthetics.

As a result of the failure of the eruption or the inadequate positioning of the teeth, there may be changes in the ideal pattern of mandibular excursive movements, prolonged retention of deciduous teeth, root resorption in the teeth proximal to this condition, among other functional and aesthetic problems. Thus, it is essential to diagnose ectopic eruption early and identify etiological factors in order to obtain early treatment to prevent the development of this pathology and, consequently, the onset of malocclusion.

#### 6. REFERENCES

- [1] Garrocho-Rangel A, Benavídez-Valadez P, Rosales-Berber MÁ, Pozos-Guillén A. Treatment of ectopic eruption of the maxillary first permanent molar in children and adolescents: A scoping review. *Eur J Paediatr Dent.* junho de 2022;23(2):94–100.
- [2] Yang P, Liang H, Zou B, Liu J, Yuan D, Meng Z, et al. Ectopic tooth in maxillary sinus compressing the nasolacrimal canal: A case report. *Medicine (Baltimore).* 7 de maio de 2021;100(18):e25514.
- [3] Gutiérrez-Marín N. Corrective Treatment of Ectopic Eruption of Permanent First Molars: Case Report. *Odvotos International Journal of Dental Sciences.* 20 de fevereiro de 2020;23(2):19–26.
- [4] Ciuraj M, Lipowicz A, Graja K, Zwolska P, Myśliwiec A. Physiotherapeutic methods of treatment of mandibular distal occlusion and the progress of therapy: A case report. *J Bodyw Mov Ther.* janeiro de 2022;29:26–32.
- [5] Helm A, Martín-Vacas A, Molinero-Mourelle P, Caleyá AM, Gallardo NE, Mourelle-Martínez MR. Ectopic Eruption of Maxillary First Permanent Molars: Preliminary Results of Prevalence and Dentoskeletal Characteristics in Spanish Paediatric Population. *Children (Basel).* 6 de junho de 2021;8(6):479.

- [6] James O, Suleiman IK, Ahmad MM, Olasoji HO. Management of rare ectopic teeth eruption: case series. *J Korean Assoc Oral Maxillofac Surg.* 30 de abril de 2023;49(2):86–90.
- [7] Ciftci V, Uguz HN, Ozcan M. Laser-assisted management of ectopic eruption of permanent incisors. *Niger J Clin Pract.* fevereiro de 2019;22(2):276–80.
- [8] Aldowsari MK, Alsaidan M, Alaqil M, BinAjian A, Albeialy J, Alraawi M, et al. Ectopic Eruption of First Permanent Molars for Pediatric Patients Attended King Saud University, Riyadh, Saudi Arabia: A Radiographic Study. *Clin Cosmet Investig Dent.* 2021;13:325–33.
- [9] Alfuriji S, Alamro H, Kentab J, Alosail L, Alali L, Altuwaijri N, et al. Ectopic Permanent Molars: A Review. *Dent J (Basel).* 29 de agosto de 2023;11(9):206.
- [10] Mehta PD, Desai N, Matariya R, Patel Y. Ectopically Positioned Tooth Piercing the Philtrum. *Ann Maxillofac Surg.* 2019;9(2):475–7.
- [11] Hemamalini R, Mahabob MN, Anbuselvan GJ, Sivagami M, Yamunadevi A, Vijayalakshmi D, et al. Ectopic Mandibular Canine with Partial Anodontia - A Rare Case Report. *J Pharm Bioallied Sci.* julho de 2023;15(Suppl 1):S795–6.
- [12] Díaz-González L, Guinot F, García C, Baltà L, Chung-Leng I. Evaluation of the position of the permanent maxillary canine and its relationship to dental anomaly patterns in the paediatric patient. *Eur J Paediatr Dent.* dezembro de 2022;23(4):281–7.
- [13] Meng M, Zhou X, Zhang Q, Zou J. Management of ectopically erupting permanent molars in a seven-year-old girl: A case report. *J Pak Med Assoc.* dezembro de 2021;71(12):2809–11.
- [14] Elmorsy K, Elsayed LK, El Khateeb SM. Case Report: Ectopic third molar in the maxillary sinus with infected dentigerous cyst assessed by cone beam CT. *F1000Res.* 2020;9:209.

- [15] Finkelstein T, Shapira Y, Pavlidi AM, Davidovitch M, Blumer S, Schonberger S, et al. Canine Transposition - Prevalence, Distribution and Treatment Considerations among Orthodontic Patients. *J Clin Pediatr Dent*. 1º de agosto de 2020;44(4):268–73.
- [16] Matsumoto MAN, Stuani MBS. Tooth transposition: a multidisciplinary approach. *Dental Press J Orthod*. fevereiro de 2018;23:97–107.
- [17] Akbas M, Karabıyık Z, Varol A. Ectopic Tooth in Mandibular Canal, Maxillary Sinus, and Mandibular Condyle. *Case Rep Dent*. 2022;2022:3118998.
18. Chintakanon K, Boonpinon P. Ectopic eruption of the first permanent molars: prevalence and etiologic factors. *The Angle Orthodontist*. 1998 Apr 1;68(2):153-60.
19. Bjerklin K, Kurol J, Valentin J. Ectopic eruption of maxillary first permanent molars and association with other tooth and developmental disturbances. *The European Journal of Orthodontics*. 1992 Oct 1;14(5):369-75.
20. Ismail MQ, Lauridsen E, Andreasen JO, Hermann NV. Ectopic eruption of the second premolar: an analysis of four different treatment approaches. *European Archives of Paediatric Dentistry*. 2020 Feb;21:119-27.