

Evaluation of alternate tillage and weed management options on production potential and weed incidences for dry seeded rice (DSR)

ABSTRACTS

The study which aims at the evaluation of conservation tillage and to identify appropriate management options for dry seeded rice (DSR) under conservation tillage system was carried out at the Bangladesh Rice Research Institute (BRRI), Regional station, Rajshahi during wet season of 2010 and 2011. The experiment was laid out in split plot design with three replications. The main plot treatments were Zero tillage (ZT), Strip tillage (ST) using power tiller operated seeder (PTOS), Minimum tillage using PTOS (MT), Permanent bed (PB) using bed planter, Fresh bed (FB) using bed planter and Conventional tillage (CT). Except conventional tillage, other alternate tillage treatments considered as conservation tillage treatments. The subplot treatments were Weed free (W_1); Post emergence herbicide + 1 hand weeding (W_2), Pre-emergence herbicide + post-emergence herbicide (W_3), Post emergence herbicide alone (W_4) and Weedy control (W_5). Pandimethylin and Bispyribac Sodium were the pre and post emergence herbicide, respectively. Among the tillage treatments, conventional tillage (CT) registered comparatively lower weeds population as well as weeds biomass. Alternate tillage treatments showed about similar result in terms of weed incidences. Again, crop weed competition almost remained same among the weed free and herbicide applied treatments. Although the number of panicle m^{-2} was comparatively lower, the higher panicle length associated with higher grains panicle⁻¹ was achieved in fresh bed and permanent bed treatments. The grain yield of rice was also comparable among the conventional tillage, bed planting system and also the zero or strip tillage treatments. On the other-hand, the post emergence herbicide W_4 treatment (Bispyribac Na⁺) was found more effective in controlling weeds where W_1 (weed free) and W_2 treatment (Bispyribac Na⁺ + 1 hand weeding) showed identical effect in relation to weed intensity, grain and straw yield of rice.

Keywords: Alternate tillage, Conservation tillage, Conventional tillage, Bispyribac Na⁺ and Herbicides

INTRODUCTION

“Rice, a cereal crop plays a vital role in global food security and acts as a main source of nutrition for millions of farm families living in Asia especially in Bangladesh. Puddling associated with conventional tillage consumes a high proportion (30-35%) of the total operational energy for rice production. At present, conventional tillage poses some serious concerns e.g. high fuel, time and labour requirement, soil compaction and deterioration in soil structure” (Mitchell *et al.*, 2009, Kumar *et al.* 2019). “Intensive tillage practices may also result in increased greenhouse gas emissions and energy requirements” (Gathala *et al.*, 2011). “It also caused long-term sustainability of intensive cultivation systems in South Asia” (Joshi *et al.*, 2005; Timsina *et al.*, 2011). “Moreover, loss of fertile topsoil from the arable land through runoff and wind erosion adds to the issues found under conventional farming practices” (Singh *et al.* 2020). “Therefore, it has drawn worldwide attention to revert the land degradation processes through sustainable land management practices” (Somasundaram *et al.* 2020). “Now the challenges of crop production are to produce more food at less cost and to improve water, labor and land productivity. One of the strategies is to grow crop with the introduction of alternate tillage systems with use of direct drill seeder. Alternate tillage options reduce the production costs, time and labor. With the advent of effective herbicides and rising concern over natural resource degradation, conservation tillage systems have gained

more attention in recent years. Moreover, the application of machineries in conservation agricultural system reduces drudgery of farm work and sustain crop production at more economic levels” (Osunbitana *et al.*, 2005). “The zero or strip tillage, reduced or minimum tillage and raised bed practices are to be considered as conservation tillage. Compared with conventional, conservation agriculture (CA) has been found to maintain or increase crop yields, improve soil fertility and reduce soil erosion” (Hobbs *et al.*, 2008; Kassam *et al.*, 2009). Conservation tillage can improve soil physical, chemical and biological properties (Saikia *et al.* 2020; Rice and Smith, 1984; Franzluebbbers *et al.*, 1994) and decrease input costs for labor, tractor and other equipment’s. Benefits from conservation tillage, include improvement of soil properties e.g. organic matter and soil water content (Somasundaram *et al.* 2019; Moreno *et al.*, 1997), aggregate stability, soil aggregation (McQuaid and Olsen, 1998), time savings and controls wind and water erosion. Growers can save 30-40% of time, labor, and fossil fuels through CA compared to conventional cropping (Hobbs and Gupta, 2004).

“Conservation tillage practices, such as zero tillage and permanent raised beds combined with straw mulching, may offset the production costs and other constraints associated with land preparation” (Hobbs, 2001). “In zero tillage, the minimal soil disturbance is achieved with special implements like tine openers, litho or coulter etc. Strip tillage decreases both the volume of soil disturbance and reduces fuel, labor and equipment costs when compared with traditional broadcast tillage” (Johansen *et al.*, 2012). “Bed planting is a technique of conservation tillage for improving resource use efficiency and increasing the yield. The practice of planting rice, high water requirement crop in the rice-wheat cropping system, on narrow raised bed was introduced recently in South Asia” (Connor *et al.*, 2003).

“The direct dry seeded rice (DSR) is a resource conservation technology as it incurs low labor expenses and is conducive to mechanization” (Bhuiyan *et al.*, 1995; Mahajan *et al.*, 2012a,b; Jeghangir *et al.*, 2005). Farooq *et al.* (2011) summarized “the benefits of DSR i.e. substantially reduce crop water requirements, soil organic matter turn over and reduce greenhouse gas emissions because of its low-input demand”. “Earlier planted DSR matures 1-2 weeks before transplanted rice (TPR), thus reducing the risk of terminal drought and allowing earlier planting of a following non-rice crop” (Saleh *et al.*, 2000). “Aerobic rice systems can reduce water application by 44% relative to conventionally transplanted systems by reducing percolation, seepage and evaporative losses, while maintaining yield at an acceptance level” (Bouman *et al.*, 2005). “Typically, DSR is established earlier than TPR; which hastens physiological maturity and reduces vulnerability to late season drought” (Tuong *et al.*, 2005). “But the traditional agronomic management practices for DSR specially gigh weed infestation is the major bottleneck in DSR especially in dry field conditions” (Harda and Shibayama, 1996; Rao *et al.*, 2007; Tomita *et al.*, 2003). “Estimated losses from weeds in rice are around 10% of total grain yield; however, such losses may be up to 53% for DSR” (Ramzan, 2003). “Successful rice cropping by dry DSR is critically dependent on weed control” (Mahajan *et al.*, 2011). “Since hand weeding is becoming expensive due to shortage of labor, use of herbicides can be a better cost-effective alternative to control weeds. Any single method of weed control may not provide season long and effective weed control. Several pre or post emergence herbicides or supplemented with hand weeding may act as good weed control in DSR” (Singh *et al.*, 2005). “Therefore, a combination of different weed management strategies should be evaluated for sustainable crop production. DSR was not expanded earlier due to lack of appropriate machinery” (Gupta *et al.*, 2002). “Now, mechanized direct seeding opens the window for expanding of DSR in the Globe. Machine sowing permits crops to take advantage of residual moisture and reduces labor requirements to establish DSR. In unfavorable rainfed environments, mechanical seeding reduces drought stress of seedlings by providing better seed covering after seeding” (Bakker *et al.*, 2000). Maurye *et al.* (1998) concluded that “mechanical row seeding led to better emergence, more uniform plant stand, faster and more efficient weeding between rows”. Thus, DSR along with better weed management options and adoption of alternate of tillage-based

practices need to be evaluated for sustainable crop production in this region.

Materials and methods

The experiment was carried out at the Bangladesh Rice Research Institute (BRRI), Regional station, Rajshahi (24°69' latitude N, 88°30' longitude E) in the wet season of 2010-11 and 2011-12 under Maize-Mungbean-Aman (wet season) rice cropping pattern. The experiment was laid out in split plot design with three replications. The main plot treatments were Zero tillage (ZT), Strip tillage (ST) using power tiller operated seeder (PTOS), Minimum tillage using PTOS (MT), Permanent bed (PB) using bed planter, Fresh bed (FB) using bed planter and Conventional tillage (CT). Except conventional tillage, other alternate tillage treatments were considered as conservation tillage treatments. The subplot treatments were Weed free (W₁); Post emergence herbicide + 1 hand weeding (W₂), Pre-emergence herbicide + post-emergence herbicide (W₃), Post emergence herbicide (W₄) and Weedy control (W₅). Pendimethalin and bispyribac Na⁺, respectively were used as pre and post emergence herbicide. The soil of the experimental plot was low in organic matter (1.35%), very low in available N (0.07%) and low in P (10 µg/g soil), K (0.19 meq/100 g soil), S (9.80 µg/g soil) and Zn (0.40 µg/g soil). Soil reaction was in slightly alkaline range (pH=8.3). Except conventional tillage (CT), rice was established in dry direct seeding in all five alternate tillage treatments.

Power tiller operated seeder (PTOS): The PTOS is a single-pass shallow-tillage seed and fertilizer drill which is attached in 2-wheel tractor (2WT) and it is 120-cm wide, allowing six rows of rice at 20-cm spacing. Operating capacity is typically 0.14–0.20 ha/hr (Hossain *et al.*, 2004). This seeder accomplishes three operations in a single pass, including tillage (up to 5cm), placement of seed and fertilizer in a furrow, and seed covering by a post-furrow opener roller bar. Compared with traditional broadcast sowing using 2-wheel

tractor (WT) full tillage, the PTOS requires less than half the time and fuel for sowing cereals, and is sometimes referred to as “reduced tillage” because of shallow seeding and the reduction in number of passes, though the 400–480 rotor RPM speed results in considerable soil surface disturbance (Wohab *et al.*, 2007).

Bed planter: Bed planter is a single 2WT attachable unit that tills the soil, places seed and fertilizer by inclined plates and finally shapes beds in a trapezoidal shape. Once beds are established, the seasonal reshaping of beds involves only reshaping of beds with minimal soil disturbance. A 2 WT operated bed planter, developed jointly by BARI, Cornell University, and CIMMYT in Bangladesh, and based on CIMMYT models developed in Mexico, can make and shape beds and place seed and fertilizer in furrows on the bed in one pass (Hossain *et al.*, 2004; Krupnik *et al.*, 2013; Wohab *et al.*, 2009). Beds of 54-cm furrow to furrow (30-35 cm top of bed width) distance are produced which can accommodate two rows of rice with about 25 cm distance.

Description of tillage treatments and establishment of crops: In CT, conventionally transplanted rice was grown on puddled soil. In such case, the land was cultivated with power tiller 3 times followed by consecutive irrigation and laddering. As a result, puddled condition was prevailed in soil. In case of all five conservation tillage treatments, glyphosate (Roundup) was applied in untilled soil @ 1.0 kg a.i. ha⁻¹ using 500 L water 5 days before sowing of seeds. In case of ZT, little slits apart from 20 cm distance rows were made in untilled soil by hook like iron made structure locally called lithao. Then rice seeds were sown manually with discontinuous seeding. Strip tillage was made by removing 2/3rd tines from PTOS and then strips were made in untilled soil through pass of PTOS. During ST, seeds and fertilizers were dropped in furrow of strips with 20 cm row to row distance where 3 rows of seeds were sown in one pass of PTOS. Minimum tillage was done through one pass with PTOS where seeds and fertilizers were also placed. Fresh beds were prepared by a bed planter in untilled soil. Permanent beds and fresh beds were remained same for the first crop; the next crops were grown in PB with only renovating the beds. Except CT and ZT, seeds were sown during tillage operation either by power tiller operated seeder (PTOS) or by

bed planter.

Herbicide Used

Glyphosate: Glyphosate was applied on untilled soil in alternate tillage treatments (except CT) @ 1.0 kg a.i./ha using 500 L water 5-6 days before sowing of seeds.

Pandimethalin: Pandimethalin is a selective pre-emergence herbicide for rice cultivation and its chemical form is “2-chloro-2: 6-diethyl-N-(2-propoxyethyl) acetanilide”. It was applied in the field after 5 days after seeding (DAS)/days after transplanting (DAT). Pandimethalin forms a thin layer of 2 cm deep on soil surface. Roots or emerging shoots of weed seedlings absorb the herbicide when they pass through this layer and are killed in the process. Necessary moisture was maintained in the field for better effectiveness of herbicide to kill the germinating weeds in the soil surface.

Bispyribac Sodium: Bispyribac Sodium 10% SC is a post emergence herbicide and was applied in the field after 15 DAS/DAT. The chemical form of Bispyribac Sodium is the “Sodium 2,6-bis [(4,6-dimethoxy-2-pyrimidinyl)oxy] benzoate”. Bispyribac Sodium is the broad spectrum systemic herbicide for control of weeds in direct sown rice, rice nursery and transplanted rice field. Bispyribac Sodium controls major grasses, sedges and broadleaf weeds of rice and offers a wide application window from 2-5 leaf stages of weeds.

Crop management practices: A medium-duration variety BRRI dhan39 which matures at 118 days was used in Aman season. All DSR plots were sown at the seed rate of 25–30 kg ha⁻¹ after initial precipitation (between 10 and 15 June) in each year. Seeds for all transplanted rice (TPR) plots were sown on seedbeds at the same time. Twenty-five days old seedlings were transplanted between 05 and 10 July. Nitrogen, Phosphorus, Potassium, Sulphur and Zinc fertilizers rates for rice in the elemental forms were 76, 12.5, 25, 8 and 7.5 kg ha⁻¹, respectively, applied as per the recommendation of the Bangladesh Rice Research Institute (BRRI). In DSR, N was broadcasted as urea in 3 equal splits (10 days after seeding (DAS), 25–30 DAS, and 45–50 DAS). In TPR, urea was broadcasted at basal, at 15–20 days after transplanting (DAT) and 35–40 DAT. Required gap filling was done after 10-15 days of seeding/transplanting. Thinning of all crops was also done after 15 DAS. Pre and post emergence herbicides were applied as per the treatments. In weed free treatment, two hand weeding was done for transplanted rice and the first hand weeding was done at 21 DAT and 2nd hand weeding was done at 42 DAT. Two times hand weeding also done in dry seeded rice and the first and second weeding hand weeding were done at 28 and 56 DAS, respectively.

Soil Analysis: The initial nine soil sub-samples were collected from the experimental field and mixed in a composite sample and then the soil sample was analyzed for determining the initial properties of soil. Analysis of variance (ANOVA) for different parameters of the treatments was also done by software package CROP STAT version 7.2 developed by International Rice Research Institute (IRRI).

Data collection: Weed data were computed in population and dry matter basis at first weeding (21 DAT /28 DAS) and at 2nd weeding (42 DAT/56 DAS). Two weed samples in each plot were collected (sampling area 0.4 m x 0.4m), separated by species and counted the number. After counting, roots of weeds were discarded and removed and other portion was oven dried at 70°C to constant weight and then weighed. At maturity stage of rice, panicles were counted from one square meter area of each treatment. Panicle length and grains/panicle were taken from 10 randomly selected plants at maturity stage of crop. Panicle length was measured in centimeter from the first neck to tip of the panicle. Then, filled and sterile spikelets were separated by submerging them in tap water. Six square meter areas in each plot were harvested to record grain yield and the yield was adjusted with the moisture content of 14% for rice. Straw was sun dried properly and measured weight. Sub samples from straw were taken and placed it in oven at 70°C temperature up to 72 hours to take oven dry weight. Straw yield adjusted to 3 % moisture content and express in t/ha.

Statistical Analysis The data were analyzed statistically through computer based statistical program

package software package CROP STAT version 7.2 developed by International Rice Research Institute (IRRI). Analysis of variance (ANOVA) for different parameters of the treatments was also done by software package CROP STAT version 7.2.

RESULTS

Effect of tillage on weed density and dry biomass before first hand weeding

The broadleaf weeds population before first hand weeding responded significantly due to tillage options in both the years (**Table 1**). The maximum broadleaf weeds population was found in MT (55.02 m⁻² and 36.67 m⁻² in 2010 and 2011, respectively) in both the years. The lowest broadleaf weeds population in first year was recorded in ZT which was statistically identical with all other treatments except MT. In 2nd year, broadleaf weed density was found lower in CT followed by ZT. It was also noticed that all the alternate tillage treatments gave statistically similar number of broadleaf weeds population during 2nd year. Unlike population, the biomass of broadleaf was not influenced due to tillage options over the years. Sedge weeds population as well as dry biomass was not affected by the tillage options in both the years (**Table 1**). The density and dry biomass of grass weeds before 1st hand weeding in 2010 were not influenced significantly by the tillage options (**Table 1**). The maximum number of weeds m⁻² during 2011 was found in ZT which was statistically resembled with ST and PB treatments. The lowest dry mass of grass weeds in 2011 was found in CT (6.54 g m⁻²) which was statistically comparable with rest of the treatments except ZT and ST while the highest biomass was recorded in ST (16.10 g m⁻²) followed by ZT and PB.

Effect of weed management options on weed density and dry biomass before first hand weeding

There was a wide variation among the weed management options on weed density and dry biomass of every category of weeds in both the years (**Table 2**). For each category of weed at first hand weeding, the lowest and the highest weed density and dry biomass were recorded in W₃ (Pandimethalin + Bispyric Na⁺) and W₅ (Weedy) treatments, respectively in both the years. It was also noticed that all herbicide treated treatments (W₂, W₃ and W₄) gave statistically lower weed density and biomass over non treated treatments across the years. W₁ and W₅ had dissimilar effect on broadleaf weeds population although those treatments showed identical effect on broadleaf weeds biomass in both the years. The sedge weeds population and dry matter were remained unchanged between W₁ and W₅ in both the years. The W₁ and W₅ treatments showed significant variation on grass weed density in both the years while grass weed biomass remained same during 2010 and those treatments gave identical effect during 2011.

Table 1: Population and dry matter of weeds as affected by tillage options before first hand weeding in Aman rice, 2010-11

Tillage	Broadleaf (No. m ⁻²)	Broadleaf wt. (g m ⁻²)	Sedge (No. m ⁻²)	Sedge wt. (g m ⁻²)	Grass weed (No. m ⁻²)	Grass weed wt. (g m ⁻²)
2010-11						
ZT	32.93	8.11	40.44	8.70	35.83	23.22
ST	50.44	14.68	32.53	6.43	35.42	23.57
MT	55.02	13.99	35.44	9.01	30.83	20.07
PB	43.76	11.09	27.53	7.34	30.00	17.23
FB	49.19	14.06	34.61	9.48	31.67	13.84
CT	37.53	10.10	26.69	5.09	28.33	10.10
LSD (0.05)	20.62	NS	NS	NS	NS	NS
CV (%)	47.5	64.3	67.1	89.7	50.6	48.3
2011-12						
ZT	31.67	6.44	20.83	3.52	27.92	13.29
ST	34.58	7.98	17.50	4.57	27.90	16.10
MT	36.67	6.85	18.33	3.71	16.67	9.63

PB	35.24	7.45	16.67	3.15	22.50	11.69
FB	34.17	8.55	20.85	3.85	14.58	7.49
CT	25.00	5.23	17.08	3.59	15.42	6.54
LSD (0.05)	15.29	NS	NS	NS	7.64	9.46
CV (%)	43.10	62.40	70.10	96.60	52.90	72.30

Table 2: Population and dry matter of weeds as affected weed management options before first hand weeding in Aman rice

Tillage	Broadleaf (No. m ⁻²)	Broadleaf wt. (g m ⁻²)	Sedge (No. m ⁻²)	Sedge wt. (g m ⁻²)	Grass weed (No. m ⁻²)	Grass weed wt. (g m ⁻²)
2010-11						
W ₁	64.93	19.21	59.72	14.05	46.88	31.03
W ₂	27.43	7.09	12.50	2.04	21.53	10.64
W ₃	14.93	3.15	7.99	1.17	13.89	6.33
W ₄	19.79	5.65	13.89	2.18	17.36	8.27
W ₅	96.88	24.88	70.14	18.92	60.42	33.74
LSD (0.05)	17.13	6.99	11.75	5.82	10.37	7.23
CV (%)	47.5	64.3	67.1	89.7	50.6	48.3
2011-12						
W ₁	51.39	11.44	32.3	7.31	30.21	17.93
W ₂	19.44	4.43	9.72	1.61	9.36	3.38
W ₃	10.76	2.12	5.90	0.95	8.33	3.20
W ₄	14.24	3.03	9.38	1.28	10.24	5.19
W ₅	68.60	14.40	35.4	7.51	45.83	24.29
LSD (0.05)	10.63	3.77	5.97	4.69	9.02	4.19
CV (%)	43.10	62.40	70.10	96.60	52.90	72.30

NS=Not significant, W₁=Weed free (2 Hand weeding), W₂=Bispyribac Na⁺ + 1HW, W₃=Pandimethylin + Bispyric Na⁺, W₄=Bispyribac Na⁺, W₅= Weedy (Control)

Effect of tillage on weeds density and dry biomass before second hand weeding

Table 3 shows the density and dry matter of weeds in different years. In 2010, the main effect of tillage did not cause significant effect on population and dry matter of broadleaf weeds before 2nd hand weeding but it did little influence on broadleaf weeds biomass in 2011. The maximum broadleaf population was found in ZT (37.08 m⁻² and 32.9 m⁻², respectively after the years in 2010 and 2011) in both the years while the highest biomass was recorded in MT (6.42 g m⁻²) and ZT (6.31 g m⁻²), respectively after the years in 2010 and 2011. Though FB recorded (25.0 m⁻² and 23.3 m⁻², respectively after the years in 2010 and 2011) minimum number of broadleaf weeds but CT (4.19 m⁻² and 3.68 g m⁻² respectively after the years in 2010 and 2011) obtained the lowest biomass in all the years. Before 2nd hand weeding, population of sedge weeds as well as biomass in 2010 remained higher in FB which was statistically comparable with other alternate tillage treatments. In 2011, the density of sedge weeds was not influenced although its biomass was affected due to tillage options. The maximum sedge weeds biomass in 2011 was also found in FB (9.8 g m⁻²) followed by MT, CT and PB treatments while the lowest biomass was obtained in ST which was statistically similar with all other treatments except MT and FB. The population and dry biomass of grass weeds before 2nd hand weeding were affected significantly by the tillage options over the years. In 2010, the maximum grass weeds population was found in ST (37.08 m⁻²) which was statistically resembled with other alternate tillage treatments. On the other-hand the highest biomass of grass weed in 2010 was observed in MT (28.07 g m⁻²), differed with only CT. In 2011, the population and biomass of grass weed were the highest in PB (20.42 m⁻², 16.88 g m⁻²) which was also statistically alike with other treatments except CT.

Table 3: Population and dry matter of weeds as affected by tillage options before 2nd hand weeding

in Aman rice

Tillage	Broadleaf (No. m ⁻²)	Broadleaf wt. (g m ⁻²)	Sedge (No. m ⁻²)	Sedge wt. (g m ⁻²)	Grass weed (No. m ⁻²)	Grass weed wt. (g m ⁻²)
2010-11						
CT	37.08	6.26	64.58	9.75	32.50	26.75
ZT	33.33	5.21	59.58	9.72	37.08	20.30
ST	36.67	6.42	62.50	16.40	31.67	28.07
MT	28.75	4.63	47.92	13.68	26.67	17.48
FB	25.00	4.51	65.00	18.93	30.83	24.70
PB	32.08	4.19	38.33	7.42	21.25	8.75
LSD (0.05)	NS	3.96	23.56	10.36	13.50	18.30
CV (%)	59.80	90.90	57.00	111.5	67.10	122.8
2011-12						
ZT	32.92	6.31	35.42	5.28	21.67	15.98
ST	25.83	4.80	33.75	5.19	18.75	12.34
MT	25.83	5.74	48.75	9.00	16.25	15.88
PB	25.42	4.92	40.42	7.59	20.42	16.88
FB	23.33	4.38	49.17	9.80	13.75	16.36
CT	27.50	3.68	47.50	8.11	10.00	6.83
LSD (0.05)	10.57	1.73	19.60	3.28	9.54	7.10
CV (%)	39.80	43.56	35.08	42.21	48.41	51.80

NS=Not significant, ZT=Zero tillage, ST=Strip tillage, MT=Minimum tillage, PB=Permanent bed, FB=Fresh bed, CT=Conventional tillage

Effect of weed management options on weeds density and dry biomass before second hand weeding

Irrespective of tillage options, there was a wide variation among the weed management treatments on population as well as dry biomass for all categories of weeds in both the years. (Fig. 1 and Fig. 2). It was found that the density and dry matter of all categories of weeds were top ranked in W₅ which was statistically differed with rest of the treatments in both the year. The population and biomass of broadleaf were lower in W₁ in both the years. The density of broadleaf weeds was remained same among W₁, W₂ and W₃ treatments in both the years. The broadleaf biomass was lower in W₁ followed by W₂ during 2010. On the other-hand, all the treatments except W₄ and W₅ produced similar broadleaf weed biomass in 2011.

The population of sedge weeds was the lowest in W₁ followed by W₂ in 2010 whereas its biomass yield was lower in W₂ which was statistically similar with other treatments except W₅. In 2011, sedge weeds population as well biomass was the lowest in W₁ which was statistically alike with W₂ and W₃ treatments. In 2010, the W₂ registered the lowest grass weed population followed by W₁ and W₃ treatments. The lowest biomass yield of grass weed during 2010 was also found in W₂ which was statistically comparable with rest of the treatments except W₅.

Fig. 1: Effect of weed management on weeds density and dry biomass of Aman rice before 2nd hand weeding, 2010-11

Fig. 2: Effect of weed management on weeds density and dry biomass of Aman rice before 2nd hand weeding, 2011-12

W₁=Weed free (2 Hand weeding), W₂=Bispyribac Na⁺ + 1HW, W₃=Pandimethylin + Bispyribac Na⁺, W₄=Bispyribac Na⁺, W₅= Weedy (Control)

Effect of tillage and weed management on yield components

Panicle m⁻²

The tillage, weed management options and tillage x weed management interactions showed significant effect on number of panicles across the years (**Table 4 and 5**). The number of panicles m⁻² remained maximum in ST and that was the minimum in FB in both the years where FB and PB showed the identical effect. Irrespective of tillage options, the W₁ recorded the maximum and W₅ obtained the significantly lower number of panicle m⁻² in both the years. The interaction result showed that the highest number of panicle m⁻² was recorded ST × W₁ which was statistically similar with all other weed free and Bispyriban Na⁺ + 1 HW weeding treatments in both the years.

Table 4: Yield components of Aman rice as affected by tillage and weed management options, 2010

	ZT	ST	MT	PB	FB	CT	Mean
Panicles m ⁻² (No.)							
W ₁	252	260	249	241	239	252	249
W ₂	236	232	226	205	206	248	226
W ₃	236	243	235	209	216	228	228
W ₄	218	226	214	223	216	227	221
W ₅	194	195	208	187	180	196	193
Mean	227	231	226	213	211	230	
LSD (0.05)	T = 10.9		W = 12.4		T x W = 30.3		
Panicle length (cm)							
W ₁	24.3	23.6	23.4	25.0	24.5	24.0	24.1
W ₂	23.6	24.0	23.3	23.7	25.1	24.0	24.0
W ₃	23.8	23.4	23.6	24.1	25.3	23.9	24.1
W ₄	23.5	22.6	23.5	24.2	23.4	23.6	23.5
W ₅	21.8	21.9	22.9	24.1	24.3	21.4	22.7
Mean	23.4	23.1	23.3	24.2	24.5	23.4	
LSD (0.05)	T = 0.93		W = 0.70		T x W = 1.73		
Grains panicle ⁻¹ (No.)							
W ₁	124	127	117	137	140	131	129
W ₂	124	125	119	141	146	135	131
W ₃	115	117	113	148	148	135	129
W ₄	125	123	120	139	142	128	130
W ₅	116	114	120	116	123	119	118
Mean	121	121	118	136	140	129	
LSD (0.05)	T = 11.7		W = 7.0		T x W = 14.2		

ZT=Zero tillage, ST=Strip tillage, MT=Minimum tillage, PB=Permanent bed, FB=Fresh bed,
CT=Conventional tillage

W₁=Weed free (2 Hand weeding), W₂=Bispyribac Na⁺ + 1HW, W₃=Pandimethylin + Bispyric Na⁺, W₄=Bispyribac Na⁺, W₅= Weedy (Control)

Panicle length

Inverse relation was prevailed between the number of panicle m⁻² and the panicle length (**Table 4 and 5**). The larger panicle length was obtained in FB followed by PB in both the years whereas the shorter panicle length was obtained in ZT and MT, respectively after the years 2010 and 2011. The panicle length of rice was remained same in W₁, W₂, and W₃ and those treatments were significantly superior over control (W₅) in both the years. The tillage and weed management showed positive interaction, the longer panicle length

was found in FB × W₃ and that was the shorter in CT × W₅ in both the years.

Table 5: Yield components of Aman rice as affected by tillage and weed management options, 2011

	ZT	ST	MT	PB	FB	CT	Mean
Panicles m² (No.)							
W ₁	246	271	264	237	226	257	250
W ₂	260	254	244	244	204	258	244
W ₃	254	267	246	230	244	223	244
W ₄	241	238	222	237	244	251	239
W ₅	194	214	204	193	174	220	200
Mean	239	249	236	228	218	242	
LSD (0.0%)	T = 18.0		W = 16.7		T x W = 28.1		
Panicle length (cm)							
W ₁	24.3	23.6	23.4	24.6	24.2	23.9	24.0
W ₂	23.7	24.2	23.6	23.2	25.1	24.0	23.9
W ₃	23.8	23.4	23.3	23.8	25.3	23.8	23.9
W ₄	23.5	22.6	23.5	24.2	23.4	23.5	23.4
W ₅	21.4	22.3	22.1	24.0	23.8	21.5	22.5
Mean	23.3	23.2	23.2	24.0	24.3	23.3	
LSD (0.05)	T = 1.11		W = 0.73		T x W = 1.50		
Grains panicle⁻¹ (No.)							
W ₁	143	152	120	147	159	156	146
W ₂	131	140	137	141	161	141	142
W ₃	113	127	120	152	157	143	135
W ₄	124	130	110	154	149	135	134
W ₅	121	123	130	132	136	121	128
Mean	127	134	124	145	153	139	
LSD (0.05)	T = 23.3		W = 10.7		T x W = 31.0		

ZT=Zero tillage, ST=Strip tillage, MT=Minimum tillage, PB=Permanent bed, FB=Fresh bed,
CT=Conventional tillage

W₁=Weed free (2 Hand weeding), W₂=Bispyribac Na⁺ + 1HW, W₃=Pandimethylin + Bispyribac Na⁺, W₄
=Bispyribac Na⁺, W₅= Weedy (Control)

Grains panicle⁻¹

Like panicle length, the higher number of filled grains panicle⁻¹ was recorded in FB followed by PB in both the years (Table 4 and 5). The MT obtained lower number of filled grains per panicle over the years. Irrespective of tillage options, the maximum number of filled grains panicle⁻¹ was found in W₂ and W₁, respectively after the years during 2010 and 2011. The lowest number of panicle was obtained in W₅ which was statistically differed from rest of the treatments in both the years. The number of filled grains panicle⁻¹ also remained unchanged among weed free and herbicide applied plots.

Effect of tillage and weed management on grain and straw yield

The grain yield of rice was little affected by the tillage options in both the years (Table 6). The CT (4.28 and 4.37 t ha⁻¹, after the years in 2010 and 2011) obtained the highest and the MT (3.64 and 3.71 t ha⁻¹, after the years in 2010 and 2011) recorded the lowest grain yield in both the years. The grain remained statistically similar among CT, FB and PB treatments in first year and among CT, ZT, FB and PB in the succeeding year. It was also noticed that the MT, ZT and ST also gave comparable grain yield of rice in both the years. A remarkable effect was observed in grain yield of rice by the weed management options across the years (Table 7). In both years, the superior grain yield of rice was found in W₁ closely followed by W₂ while the significantly lower grain yield was recorded in W₅. Among herbicide applied treatments, grain yield of rice was significantly superior in W₂ over W₃ and W₄ in both the years. The tillage and

weed management options showed a highly significant interaction on grain yield of rice. All the weedy plots proved inferior causing significantly lower grain yield compared with other plots in all the years. The grain yield was found higher in CT×W₁ followed by FB×W₁ in 2010 while it remained maximum in PB×W₁ in 2011.

The tillage options did not produce significant effect on straw yield in first year while it had done slight effect in succeeding year. Straw yield ranges from 5.44 t ha⁻¹ in ST to 5.83 t ha⁻¹ in FB in 2010. In following year, the highest straw yield was found in PB treatment which is statistically identical with all other treatments except MT. Weed management options produced significant influence on straw yield of rice in both the years. Like grain yield, the highest straw yield of rice during 2010 was found in W₁ followed by W₂. In 2011, W₁ and W₃ recorded the highest straw yield of rice and those treatments were statistically identical with W₂. The straw yield of rice was significantly lower in W₅ than other treatments in both the years. The highest straw yield (6.86 t ha⁻¹) was achieved in ST×W₁ during 2010. In 2011, the highest straw yield of rice was found in ST×W₃ which was statistically identical with all of weed free and herbicide applied plots with little exception. The ZT plus W₅ (4.30 t ha⁻¹) and MT plus W₅ (4.45 t ha⁻¹) recorded the lowest straw yield of rice during 2010 and 2011, respectively.

Table 6: Grain and straw yield of Aman rice as affected by tillage and weed management options, 2010 and 2011

	ZT	ST	MT	PB	FB	CT	Mean
2011							
Grain yield (t ha⁻¹)							
W ₁	4.56	4.54	4.04	4.89	4.89	4.99	4.65
W ₂	4.47	4.50	4.34	4.76	4.75	4.71	4.59
W ₃	4.21	3.88	3.81	4.61	4.54	4.46	4.25
W ₄	3.81	3.71	3.12	4.16	4.17	4.12	3.85
W ₅	2.72	2.85	3.33	2.69	2.70	3.13	2.90
Mean	3.95	3.89	3.73	4.22	4.21	4.28	
LSD (0.05)	T = 0.37		W = 0.59		T x W = 0.73		
Straw yield (t ha⁻¹)							
W ₁	6.28	6.86	6.23	6.21	6.46	6.01	6.34
W ₂	6.47	5.78	5.80	6.30	5.81	6.34	6.08
W ₃	4.80	5.04	5.06	5.74	6.56	6.11	5.55
W ₄	5.69	5.10	5.14	6.00	5.66	5.48	5.51
W ₅	4.30	4.41	5.46	4.46	4.65	5.14	4.74
Mean	5.51	5.44	5.54	5.74	5.83	5.82	
LSD (0.05)	T = 0.31		W = 0.56		T x W = 1.33		
2011							
Grain yield (t ha⁻¹)							
W ₁	4.99	4.87	3.85	5.11	5.01	5.12	4.82
W ₂	4.50	4.64	4.54	4.87	5.01	5.00	4.76
W ₃	3.79	4.11	4.04	4.49	4.44	4.35	4.20
W ₄	3.40	3.55	3.76	4.37	4.07	4.27	3.90
W ₅	2.95	3.08	2.36	2.82	2.36	3.13	2.78
Mean	3.93	4.05	3.71	4.33	4.18	4.37	
LSD (0.05)	T = 0.39		W = 0.60		T x W = 0.81		
Straw yield (t ha⁻¹)							
W ₁	6.84	6.74	5.86	6.25	6.13	6.63	6.41
W ₂	6.10	6.08	6.22	6.48	6.13	5.99	6.17
W ₃	6.00	6.92	5.50 ₁₀	6.60	6.89	6.55	6.41

W ₄	5.64	5.68	6.08	6.75	5.66	5.80	5.94
W ₅	5.34	4.80	4.45	5.27	5.07	5.48	5.07
Mean	5.99	6.04	5.62	6.27	5.98	6.09	
LSD (0.05)	T = 0.34	W = 0.47	T x W = 1.22				

DISCUSSION

Effect of tillage on weed incidences

The density of broadleaf weeds at first weeding affected significantly due to tillage options and the weeds population remained higher in MT during both the years while it was the lowest in ZT and CT, after the years during 2010 and 2011. In some cases, abundance of weeds especially broadleaf weeds was comparatively lower in zero tillage or some other alternate tillage treatments. Fewer annual broadleaf weeds in zero tillage fields than CT were reported by Singh, 2007; Malik *et al.* 2002. Similarly, Swanton *et al.* (1999) reported that the conventional tillage system was associated with some broadleaf weeds. Cussans (1976) found the higher broadleaf weeds with increase in tillage intensity. The number of broadleaf weeds at 2nd weeding was remained higher in ZT in both the years while the highest biomass was recorded in MT and ZT, after years during 2010 and 2011. Although FB recorded lower numbers of broadleaf weeds but CT obtained the lowest biomass at 2nd weeding in both the years. Therefore, contrasting results was found regarding broadleaf weeds abundance. In some cases, CT and zero or strip tilled plots showed a small difference in weeds intensity which was supported by Derksen *et al.* (1993) and Wrucke and Arnold (1985). The sedge weeds abundance at 2nd weeding was also found lower in CT where all conservation tillage obtained higher weeds compared with CT. Similarly, the density and biomass of grassy weeds at 1st weeding remained lower in CT. Similar trend of grassy weeds intensity remained at 2nd weeding. The lower density and dry matter of grassy weeds in CT were in same line with the findings of Ozpinar, 2006; Gill and Arshad, 1995; Froud-Williams *et al.* 1981; Moody and Mian 1979. Majid *et al.* (1998) and Das and Choudhury (1985) reported that tillage is one of most practical methods for perennial weed control. Compared with CT, the density and dry mass of grassy weeds were remained higher in all alternate conservation tillage treatments. Many researchers stated that reduced or no-till system increased perennial grass weeds densities and diversity (Chokkor *et al.*, 2007; Swanton *et al.*, 1999; Moody, 1992; Gill and Arsad, 1995; Moyer *et al.*, 1994; Nyarko and De Datta, 1991; Feldman *et al.*, 1997).

Effect of weed management options on weed incidences

There was a wide variation among the weed management options on density and dry biomass of weeds at first and 2nd weeding. The crop weed competition at 28 DAS in both the years was markedly reduced by two herbicide (W₃) applied treatment. It was noticed that all of the herbicide applied treatments (W₂, W₃ and W₄) recorded statistically lower weed density and biomass over non treated treatments at 1st weeding across years. Herbicide use can markedly affect crop-weed competition was reported by Chauhan *et al.*, 2012, 2006; Rao *et al.*, 2007; Catheart *et al.*, 2004. Weed management using herbicide has become an integral part of modern agriculture was also reported by Maity and Mukherjee, 2008; Rao *et al.*, 2007. Although the weed control at 1st weeding was found better in W₃ than W₂ and W₄ in both the years but this difference was insignificant indicated that the pre-emergence herbicide (pendimethalin) did least effect on weed suppression. Poor weed control due to pre-emergence herbicide (such as pendimethalin) under CA system was also reported by Chauhan *et al.*, 2012; Mahajan and Timsina, 2011; Walker and Bond 1977; Savage and Barrentine, 1969. In CA systems, effectiveness of pre-emergence herbicides was found lower (Hartzler and Owen, 1997). Similarly, Chauhan *et al.*, (2006) reported that within CA

systems of low soil disturbance, the pre-emergence herbicide on soil surface was susceptible to loss through volatilization and photodecomposition. In contrast, Jayadeva and Bhairappanavar (2002) and Singh *et al.* (2006) reported that the pendimethalin as pre-emergence herbicide was quite effective for weeds suppression.

More weeds suppression at 1st weeding by W₃ might be due to the more suppression of weeds by post emergence-Bispyribac Na⁺. Moreover, weeds at 2nd weeding remained similar between W₃ and W₄ in both the years. This happened due to more suppression of weeds by Bispyribac Na⁺. The effective weeds suppression by Bispyribac Na⁺ were also reported by Mahajan and Timsina, 2011; Mahahjan *et al.*, 2009; Risi *et al.*, 2004, Schmidt *et al.*, 1999. At 1st and 2nd weeding, the population and dry biomass of weeds were found lower in W₁ followed by W₂ over the years. The longer weed free period obviously recorded the lower weeds density as well as dry biomass supported by the findings of Sanjay *et al.* (2008). The W₂ also performed better in regards to weeds suppression than other herbicide applied treatments at 2nd weeding. Integration of recommended herbicide with one hand weeding significantly reduced the weeds density and dry matter compared to application of one or more herbicide which was supported by Mahajan *et al.* (2009). Bispyribac Na⁺ supplemented with one hand weeding has been found very effective against most of the weeds (Misra and Singh, 2012; Rao *et al.* 2007). Similar findings were also reported by Mahajan and Timsina (2011) and Risi *et al.*, (2004).

Effect on yield and yield components

The number of panicle m⁻² remained lower in FB in both the years while there was inverse relation was prevailed between the number of panicle m⁻² and the panicle length. The shorter panicle length was obtained in ZT and MT, after the years during 2010 and 2011. The larger panicle length was obtained in FB followed by PB in both the years. Connor *et al.* (2003) reported that panicle length of rice was found to be longer in transplanted rice on beds than conventional tillage. length. Like panicle length, the higher number of filled grains per panicle was recorded in FB followed by PB in both the years. The MT obtained lower number of filled grains per panicle over the years. The results also indicated that the raised bed plots (either fresh bed or permanent bed) gave comparatively higher number of filled grains per panicle in both the years. Connor *et al.* (2003) also reported that numbers of grains panicle⁻¹ was higher on beds (173) than conventional tillage (163). The grain yield of rice was little affected by the tillage options in both the years. The CT and MT recorded the highest and the lowest grain yield in both the years. In first year, the grain yield remained statistically similar in CT, FB and PB treatments while in succeeding year the grain yield remained in similar in CT, ZT, FB and PB treatments. Cavalaris and Gemtus (2004) showed that the reduced tillage gave similar yield with the conventional tillage presented a lower energy productivity indicating that it was the most possible method to substitute the plough and obtain significant benefits. Fahong *et al.* (2003) and Grover *et al.* (2004) also reported that bed planting technology increased the yield and net returns to the farmers.

Irrespective of tillage options, the maximum number of filled grains panicle⁻¹ was found in W₂ and W₁ after the years during 2010 and 2011. The lowest number of panicle m⁻² was obtained in W₅ which was statistically differed from rest of the treatments in both the years. The number of filled grains panicle⁻¹ was also remained unchanged among weed free and herbicide applied plots.

The interaction result showed that the highest number of panicle m⁻² was recorded ST with W₁ which was statistically similar with all other weed free and Post-emergence + 1 HW weeding plots in both the years. It was noticed that the weedy (control) plots showed the comparatively lower number of panicle m⁻² over the years. The panicle length of rice was remained same in W₁, W₂, and W₃ and those treatments were significantly superior over control (W₅) in both the years. The tillage and weed management showed positive interaction, the longer panicle length was found in FB with W₃ and shorter in CT with W₅ in both

the years. The results also revealed that the weedy plots showed inferiority in decreasing panicle length compared with the other treatments across the years. Irrespective of tillage treatments, the superior grain yield of rice was found in W₁ closely followed by W₂ across the years while the significantly lower grain yield was recorded in W₅. Among herbicide applied treatments, grain yield of rice was significantly superior in W₂ over W₃ indicated that post-emergence herbicide supplemented with one hand weeding is necessary to obtain better grain yield. Similar result was also reported by Singh *et al.* (2005). The interactions result also indicated that almost all the weed free and Post-emergence + 1 hand weeding plots showed identical effect in relation to grain yield of rice.

Conclusion

The CT obtained the comparatively lower weeds population as well as weed biomass across the years where alternate tillage treatments showed about similar result. Some cases weed intensity was also less in ZT and FB treatments. Again, crop weed competition was almost remained same among the weed free and herbicide applied treatments. Although the number of panicles was comparatively lower, the higher panicle length associated with higher number of grains per panicle was achieved in fresh bed or permanent bed. The grain yield of rice was also comparable among the conventional tillage, bed planting system and also the zero or strip tillage treatments. On the other-hand, the post emergence herbicide (Bispyribac Na⁺) was more effective in controlling weeds where weed free (W₁) and W₂ (Bispyribac Na + 1 HW) treatments showed identical effect in relation to weed intensity, grain and straw yield of rice. Among herbicide applied treatments, grain yield of rice was significantly superior in W₂ (Nominee gold + 1 HW) over W₃ indicated that post-emergence herbicide supplemented with one hand weeding is necessary to obtain better grain yield.

REFERENCES

- Bakker RR, Bell MA, Rickman JF. 2000; Mechanization issues in tillage and crop establishment for dry direct-seeded rice. Proceedings of International Workshop on Direct Seeding in Asian Rice Systems: Strategic Research Issues and Opportunities, 25-28 January 2000, Bangkok, Thailand. Los Banos (Philippines): IRRI. pp. 219-230.
- Bhuiyan SI, Sattar MA, Khan MAK. 1995; Improving water use efficiency in rice irrigation through wet seeding. *Irrigat. Sci.* 16:1-8.
- Bouman BAM, Pend S, Castaneda AR, Visperas RM. 2005; Yield and water use of tropical aerobic rice systems. *Agric. Water Manage.* 74: 87-105.
- Catheart RJ, Chandler K, Swanton CJ. 2004; Fertilizer N rate and the response of weeds to herbicide. *Weed Sci.* 52:578-583.
- Cavalaris CC, Gemotus TA. 2004; Evaluations of four reduced tillage methods with different rotations. The proceedings of CGIAR International Conference, 11-14 October 2004, Beijing, China.
- Chauhan BS, Gill G, Preston C. 2006; Tillage systems affect trifluralin bioavailability in soil. *Weed Sci.* 54:941-947.
- Chauhan BS, Singh RG, Mahajan G. 2012; Ecology and management of weeds under conservation agriculture: A review. *Crop Protect.* 38:57-65.

- Chhokar RS, Sharma RK, Jat GR, Pundir AK, Gathala MK. 2007; Effect of tillage and herbicides on weeds and productivity of wheat under rice-wheat growing system. *Crop Protection*, **26** 1689–1696.
- Conner DJ, Timsina J, Humphreys E. 2003; Prospects for permanent beds for the rice-wheat system. In: *Improving Productivity and Sustainability of Rice-Wheat systems: Issues and Impact*. Am. Soc. Agron. Spec. Publ. 65:197-210.
- Cussans GW. 1976; The influence of changing husbandry on weeds and weed control in arable crops. In: *Proceedings of British Crop Protection Conference-Weeds*, Brighton, UK. pp. 1001-1008.
- Das JC, Chaudhury AK. 1985; Effect of soil management practices and weeds control in Rice in Southeast Asia: Methods and trends. *Philipp. Weed Sci. Bull.* 4:39-65
- Derksen DA, Lafond GP, Thomas AG, Loeppky HA, Swanton CJ. 1993; Impact of agronomic practices on weed communities: tillage systems. *Weed Sci.* 41:409-417.
- Fahong W, Xuqing W, Sayre K. 2003; Comparison study on two different planting methods for winter wheat in China. In *Bed planting training course*, 19 May-21 June 2003. Int. Maize and Wheat Impr. Cen. Mexico. D.F. 06600, Mexico.
- Farooq M, Siddique KHM, Rehman H, Aziz T, Dong-Jin Lee, Wahid A. 2011; Rice direct seeding: Experiences, challenges and opportunities. *Soil Tillage Res.* 111:87-98.
- Feldman SR, Alzugaray C, Torres PS, Lewis P. 1997; The effect of different tillage systems on the composition of the seed bank. *Weed Res.* 37:71–76.
- Franzluebbers AJ, Hons FM, Zuberer DA. 1994; Long-term challenges in soil carbon and nitrogen pools in wheat management systems. *Soil Sci. Soc. Am. J.* 58:1639-1645.
- Froud-Williams RJ, Chancellor RJ, and Drennan DSH. 1981; Potential changes in weed flora associated with reduced cultivation systems in cereal production in temperate regions. *Weed Res.* 21:99-109.
- Fukai S. 2002; Rice cultivar requirements for direct seeding in rainfed lowlands. In: Pandey S, Mortimar M, Wade L, Tuong TP, Lopes K, Hardey B. (Eds.): *Direct Seeding: Research Strategies and Opportunities*: International Rice Research Institute, Los Banos, Philippines, pp. 257-269.
- Gathala MK, Ladha JK, Kumar V, Saharawat YS, Sharma PK, Pathak H. 2011; Tillage and crop establishment affects sustainability of South Asian rice-wheat system. *Agronomy J.* 103:961–971.
- Gill KS, Arshad MA. 1995; Weed flora in the early growth period of spring crops under conventional, reduced and zero tillage systems on a clay soil in northern Alberta, Canada. *Soil Tillage Res.* 33:65-79.
- Grover DK, Singh J, Singh R, Dhillon SS. 2004; Socio-economic impact of Rice-wheat system. In *Improving Productivity and Sustainability of Rice-Wheat-systems: Issues and Impact*. Assessment of bed planting technology in Punjab. Presented in Nat. Conf. Conservation

- Agriculture: Conservation Resources-Enhancing Productivity. 22-23 September 2004. NASC complex, Dev Prakash Marg, Pusa, New Delhi 110012. Abst. pp 7-8.
- Gupta RK, Hobbs PR, Ladha JK, Probhakar SVRK. 2002; Resource conserving technologies. Transforming the rice-wheat systems of the Indo-Gangetic Plains. Rice-Wheat consortium- a success story. Asia pacific association of agricultural research institutes, Bangkok, Thailand.
- Harda J, Shibayama H. 1996; Weeds in Tropics. Association for International Co-operation of Agriculture and Forestry (AICAF), Tokyo, Japan.
- Hartzler RG, Owen MD. 1997; Weed management in Conservation Tillage Systems. Iowa State University. Ames. Available online www.extension.edu/Publications/PM1176
- Hobbs PR. 2003; Zero tillage technology: troubleshooting tips. In Addressing Resource Conservation Issues In Rice-Wheat systems of South Asia: A Resource Book, Rice-Wheat Consortium for the Indo-Gangetic Plains. Intl. Maize and Wheat. Impr. Cent., New Delhi, India. pp. 119-122.
- Hobbs PR. 2001; Tillage and crop establishment in South Asian rice-wheat system: present practices and future options. J. Crop Prod. 4:1-22.
- Hobbs PR, Gupta RK. 2004; Problem and challenges of no-till farming for rice-wheat systems of the Indo-Gangetic Plains in South Asia. In: Lal R et al. (Eds): Sustainable Agriculture and the Rice-Wheat System. Ohio State University and Marcel Dekker, Inc. Columbus, Ohio and New York, USA. pp. 101-119.
- Hobbs PR, Sayre K, Gupta R. 2008; The role of conservation agriculture in sustainable agriculture. Phil. Trans. R. Soc. Lond., B, Biol. Sci. B. 363:543-555.
- Hossain I, Sufian MA, Sarkar MAZ, Haque ME, Rahman ABMM. 2004; Power tiller operated bed planter for improved crop establishment. J. Sci. Technol. 2:17-23.
- Jayadeva HM, Bhairappanavar ST. 2002; Chemical weed control in drum seeded rice. Indian J Weed Sci. 34:290-292
- Jehangir WA, Mahish I, Ahmed S, Gill MA, Ahmed M, Mann R A, Chaudhary, MR, Turrall H. 2005; Sustaining crop water productivity in rice-wheat systems of South Asia: A case study from Punjab Pakistan. In: Draft Working Paper, Inter. Water Management Institute, Lahore, Pakistan.
- Johansen CJ, Haque ME, Bell RW, Thierfelder C, Esdaile RJ. 2012; Conservation agriculture for small holder rainfed farming: Opportunities and constraints of new mechanized seeding systems. Field Crops Res. 132:18-32.
- Joshi PK, Singh NP, Singh NN, Gerpacio RV, Pingali PL. 2005; Maize in India: Production Systems, Constraints, and Research Priorities. Mexico, D.F. CIMMYT.
- Kassam A, Friederich T, Shaxson F, Pretty J. 2009; The spread of conservation agriculture: justification, sustainability and uptake. Int. J. Agric. Sust. 7:292-320.
- Kumar N, Nath CP, Hazra KK, Das K, Venkatesh MS, Singh MK, Singh SS, Praharaj CS, Singh NP. 2019; Impact of zero-till residue management and crop diversification with legumes on soil

- aggregation and carbon sequestration. *Soil Tillage Res* 189:158–167. <https://doi.org/10.1016/j.still.2019.02.001>
- Krupnik TJ, Santos vale S, McDonald AJ, Justice S, Hossain I, Gathala MK. 2013; Made in Bangladesh: Scale-Appropriate Machinery for Agricultural Resource Conservation. CIMMYT, Mexico D.F. pp. 126.
- Linquist BA, Fischer A, Godgrey L, Greer C, Hill J, Koffler K, Moeching M, Mutters R, Van Kessel C. 2008; Minimum tillage could benefit California rice farmers. *Cal. Agric.* 62(1):24-29.
- Mahajan G, Johnson DE, Chauhan BS. 2009; Weed management in aerobic rice in north western Indo-Gangetic Plains. *J. Crop Improve.* 23:366-382.
- Mahajan G, Chauhan BS, Timsina J, Singh PP, Singh K. 2012a; Crop performance and water- and nitrogen-use efficiencies in dry-seeded rice in response to irrigation and fertilizer amounts in northwest India. *Field Crops Res.* 134:59-70
- Mahajan G, Timsina J, Jhanji S, Sekhon NK, Singh K. 2012b; Cultivar response, dry matter partitioning and nitrogen-use efficiency in dry direct-seeded rice in north-west India. *J. Crop Improve.* 26(6):767-790.
- Mahajan G, Timsina J. 2011; Effect of nitrogen rates and weed control methods on weeds abundance and yield of direct-seeded rice. *Archives Agron. Soil Sci.* 57(3):239-250.
- Mahajan G, Timsina J, Singh K. 2011; Performance and water use efficiency of rice relative to establishment methods in northwestern Indo-Gangetic Plains. *J. Crop Improve.* 25:597-617.
- Maity DJ, Mukherjee PK. 2008; Integrated weed management in dry direct seeded rainy season rice. *Indian J. Agron.* 53:116-120.
- Majid A, Farooq M, Ahmed SI, Multi AI. 1998; Wet and dry tillage practices in paddy production, *AMA.* 19(3):29-31.
- Malik RK, Yadov A, Singh S, Malik RS, Balyan RS, Banga RS, Sardana PK, Jaipal S, Hobbs PR, Gill G, Singh S, Gupta RK, Bellinder R. 2002; Herbicide resistance management and evaluation of zero tillage- A success story. Chaudhury Charan Singh Haryana Agricultural University, Hisar, India.
- Maurye PR. 1998; Comparison of zero tillage and conventional tillage in wheat and maize production under different soils and climatic conditions in Nigeria. *AMA.* 19:30-32.
- McQuaid BF, Olsen GL. 1998; Soil quality indices of piedmont sites under different management systems. *Adv. Soil. Sci.* 21:427-433.
- Misra JS, Singh VP. 2012; Tillage and weed control effects on productivity of a dry seeded rice-wheat system on a vertisol in central India. *Soil Tillage Res.* 123:11-20.
- Mitchell JP, Shrestha A, Campbell-Mathews M, Giacomazzi D, Goyal S, Bryant D, Herrera I. 2009; Strip-Tillage in California's Valley University of California, Division of Agriculture and Natural Resources. ANR Publication p. 8361.
- Moody 1992; Weed management in wet seeded rice in tropical Asia. *Food Fertil. Technol. Centre,* 1-9.

- Moody K, Mian AL. 1979; In: Rainfed Lowland Rice. Selected Papers for IRRI Conference, IRRI, Los Banos, Philippines, pp 235-245.
- Moreno F, Pelegrin F, Fernande JE, Murillo JM. 1997; Soil physical properties, water depletion and crop development under traditional and conservation tillage in southern Spain. *Soil Tillage Res.* 41:25-42
- Moyer JR, Rosman ES, Lindwall CW, Blackshaw RE. 1994; Weed management in conservation tillage systems for wheat production in North and South America. *Crop Protect.* 13:243-259
- Nyarko KA, De Datta SK. 1991; A Hand Book for Weed Control in Rice. International Rice Research Institute, pp 109-115.
- Osunbitana JA, Oyedeleb DJ Adekalu KO. 2005; Tillage effects on bulk density, hydraulic conductivity and strength of a loamy sand soil in southwestern Nigeria. *Soil Tillage Res.* 82:57-64.
- Ozpinar, S. 2006; Effect of tillage systems on weed population and economics for winter wheat production under the Mediterranean dry-land conditions. *Soil Tillage Res.* 87:1-8.
- Ramzan M. 2003; Evaluation of various planting methods in rice-wheat cropping system. Punjab, Pakistan, Rice Crop Report, 2003-04, pp. 4-5.
- Rao AN, Johnson DE, Sivaprasad B, Ladha JK, Mortimar AM. 2007; Weed management in direct seeded rice. *Adv. Agron.* 99:153-255.
- Rice CW, Smith MS. 1984; Short term immobilization of fertilizer nitrogen at the surface no-till and plowed soils. *Soil Sc. Soc. Am. J.* 48:295-297.
- Risi C, Arcangeli G, Cantoni A, Campani E. 2004; Nominee (Bispyribac Na sodium): A new post emergence herbicide for rice. *Informat Fitopatol.* 54:44-49.
- Saikia R, Sharma S, Thind HS, Singh Y (2020); Tillage and residue management practices affect soil biological indicators in a rice wheat cropping system in north-western India. *Soil Use Manag.* 36:157-172.
- Saleh AFM, Mazid MA, Bhuiyan SI. 2000; Agrohydrologic and drought-risk analyses of rain-fed cultivation in northwest Bangladesh. In: Tuong TP, Kam SP, Wade LJ, Pandey S, Bouman BAM, Hardy B, editors. *Characterizing and understanding rainfed environments*, Manila (Philippines): International Rice Research Institute. pp. 233-244.
- Sanjay MT, Setty TK, Prabhakaran, Nanjapa HV. 2008; Investigation of crop establishment methods and weed management practices on productivity and economics of rice. *Mysore J. Agril. Sci.* 42 (1):60-66.
- Savage K, Barrentine W. 1969; Trifluralin persistence as affected by depth of soil incorporation. *Weed Sci.* 17:349-352.
- Schmidt LA, Talbert RE, Baldwin FL, Rutledge EFS, Wheeler SS. 1999; Performance of Bispyribac Na+ in rice weed control programs. *Proc. South Weed Sci. Soc.* 52:49.
- Singh KK, Sharma SK. 2005; Conservation Tillage and Crop Residue Management in Rice-Wheat

- Cropping System. In: Abrol IP, Gupta RK, Malik RK (Eds.) 2005. Conservation Agriculture: Status and Prospects. Centre for Advancement of Sustainable Agriculture, New Delhi. pp. 242.
- Singh S. 2007; Role of management practices on control of isoproturon resistant little seed canary grass (*Phalaris minor*) in India. Weed Technol. 21:339-346.
- Singh S, Bhusan L, Ladha JK, Gupta RK, Rao AN, Siva Prasad B. 2006; Weed management in dry seeded rice (*Oryza sativa*) cultivated in the furrow-irrigated raised-bed planting system. Crop Protect. 25 487-495.
- Singh Y, Singh G, Johnson D, Mortimer M. 2005; Changing from transplanted rice to direct seeding in the rice-wheat cropping system in India. In: Rice is Life: Scientific Perspective for the 21st Century, Tsukuba, Japan: Proceedings of the World Rice Research Conference, 4-7 November 2004. pp. 198-201.
- Somasundaram J, Sinha NK, Mohanty M, Chaudhary RS, Hati KM, Singh RK, Biswas AK, Shukla AK, Dalal RC, Patra AK (2019; Soil hydrothermal regimes as affected by different tillage and cropping systems in a rainfed Vertisol. J Ind. Soc. Soil Sci. 66(4): 362–369
- Somasundaram J, Sinha NK, Dalal RC, Lal R, Mohanty M, Naorem AK, Hati KM, Chaudhary RS, Biswas AK, Patra AK, Chaudhari SK. 2020; No-till farming and conservation agriculture in South Asia—issues, challenges, prospects and benefits. Crit. Rev. Plant Sci. 39(3):236–279
- Swanton CJ, Shrestha A, Roy RC, Ball-Coelho BR, Knezevic SZ. 1999; Effect of tillage systems, N, and cover crop on the composition of weed flora. Weed Sci. 47:454-461.
- Timsina J, Buresh RJ, Dobermann A, Dixon J. 2011; Rice-maize systems in Asia: current situation and potential. Book published by IRRI as an IRRI-CIMMYT joint publication, p. 235.
- Tomita S, Miyagawa S, Kono Y, Noichana C, Inamura T, Nagata Y, Sributta A, Nawata E. 2003; Rice yield losses by competition with weeds in rainfed paddy fields in north-east Thailand. Weed Biol. Manag. 3:162-171.
- Tuong TP, Bouman BAM, Mortimer M. 2005; More rice, less water integrated approaches for increasing water productivity in irrigated rice-based systems in Asia. Plant Prod. Sci. 8:231–241.
- Walker A, Bond W. 1977; Persistence of the herbicide AC 92,553 [N-(1-ethylpropyl)-2,6-dinitro-3,4-xylidine] in soils. Pesticide Sci. 8:359-365.
- Watkins KB, Anders MM, Windham TE. 2004; An economic comparison of alternative rice production systems in Arkansas. J. Sust. Agric. 24(4):57-78.
- Wrucke MA, Arnold WE. 1985; Weed species distribution as influenced by tillage and herbicides. Weed Sci. 33:853-856.
- Wohab MA, Roy KC, Haque ME, Abedin MJ, Islam MS. 2007; Field performance evaluation of BHT seeder for sowing and tilling. J. Socioecon. Res. Dev. 4:116–119.
- Wohab MA, Ziauddin ATM, Haque ME, Sayre KD, Hossain MI. 2009; Development of single pass bed planter for small holder farming, crop diversification and resource conservation. In: Proc. Int.

Conf. Fourth World Congress of Conservation Agriculture, New Delhi, India, p. 302.

Yadvinder-Singh, Humphreys E, Kubal SS, Dhillon SS, Balwinder-Singh, Amanpreet-Kaur, Thaman S, Prashar A, Yadav S, Navneet-Kaur, Smith DJ, Timsina J, Gajri PR. 2008; Crop performance in a permanent raised bed rice-wheat cropping system in Punjab, India. *Field Crops Res.* 111:1-20.