

Prevalence and treatment outcomes of cutaneous leishmaniasis in Baluchistan, Pakistan

Abstract:

Leishmaniasis a diseased condition is caused by approximately 20 species of the genus *Leishmania*. It is a neglected vector-borne parasitic infection that results in a wide range of illnesses with various symptoms. Numerous risk factors such as poverty, hunger, migration, and substandard housing, attributes to its occurrence and prevalence. It has been documented that in Pakistan, the prevalence of Cutaneous leishmaniasis (CL) is on the rise. Hence it has been considered as a serious public health issue for the nation. Furthermore, areas that border Afghanistan and in the towns that have seen the greatest number of incoming migrants has shown more incidence of CL. Considering the geographical location and spread of disease across the border area this review aims to highlight the current reported prevalence and treatment outcomes of cutaneous leishmaniasis in Baluchistan, Pakistan. The review was carried out to report the findings of studies (original articles) conducted to highlight the prevalence of CL in Baluchistan Pakistan in last 5 years along with its treatment outcomes. The articles were recruited by entering “Leishmaniasis”, “Cutaneous leishmaniasis (CL)”, “Prevalence”, “Treatment outcome”, “Baluchistan” and, “Pakistan” as key words in different literature search engines including PubMed, Scopus, Medline search and Google Scholar. The reported study in last 5 years have demonstrated scattered prevalence of the CL in the province Baluchistan. Every district has shown diverse course of disease and different data. Though the consistent thing identified was the trend of increase in the incidence and prevalence across the province. The male gender and first 3 decades of life has been reported as the high-risk group. The most frequent sites of the lesions were reported to be the arms and legs. The review was carried out to report the findings of studies (original articles) conducted to highlight the prevalence of CL in Baluchistan Pakistan in last 5 years along with its treatment outcomes.

Introduction:

Leishmaniasis a diseased condition is caused by approximately 20 species of the genus *Leishmania*. It is a neglected vector-borne parasitic infection that results in a wide range of

Comment [MF1]: delete

illnesses with various symptoms. Numerous risk factors such as poverty, hunger, migration, and substandard housing, attributes to its occurrence and prevalence^{1, 2}. It has been reported that “Cutaneous leishmaniasis (CL)” is the most prevalent type of leishmaniasis, which affects 0.6 - 1 million new individuals globally each year. CL is not life-threatening, but it is nonetheless crucial to identify and treat since it can result in stigma, long-term psychological effects, irreversible scarring, and a worse quality of life^{3,4}.

It has been documented that in Pakistan, the prevalence of CL is on the rise. Hence it has been considered as a serious public health issue for the nation. Furthermore, areas that border Afghanistan and in the towns that have seen the greatest number of incoming migrants has shown more incidence of CL^{5,6}. Additionally It has been highlighted by Postigo who pointed out that outbreaks of both anthroponotic cutaneous leishmaniasis (anthroponotic CL) caused by *Leishmania tropica* (*L. tropica*) and zoonotic CL produced by *Leishmania major* (*L. major*) have been observed in Pakistan in particular. While *L. major* is more prevalent in the nation's rural parts, *L. tropica* is mostly found in metropolitan areas^{7,8}.

Comment [MF2]: Reeated in abstract

It has been reported that following malaria, CL is the second most common vector-borne illness in Pakistan and is endemic in a number of the country's regions. Pakistan's endemic regions for illness include parts of South Punjab, Khyber Pakhtunkhwa, Baluchistan, and Interior Sindh^{9, 10}. Considering the geographical location and spread of disease across the border area this review aims to highlight the current reported prevalence and treatment outcomes of cutaneous leishmaniasis in Baluchistan, Pakistan.

Comment [MF3]: Reeated in abstract

Methodology:

The review was carried out to report the findings of studies (original articles) conducted to highlight the prevalence of CL in Baluchistan Pakistan in last 5 years along with its treatment outcomes. The articles were recruited by entering “Leishmaniasis”, “Cutaneous leishmaniasis (CL)”, “Prevalence”, “Treatment outcome”, “Baluchistan” and, “Pakistan” as key words in different literature search engines including PubMed, Scopus, Medline search and Google Scholar.

Results & Discussion:

Studies from Baluchistan, the largest province by land located in southwest Pakistan have reported that it has the highest burden of CL followed by Khyber Pakhtunkhwa¹¹. The geographic dispersion, as stated by Firdous et al., is a result of the sandfly vector, coupled with activities affecting their environment, such as wars, deforestation, and agricultural activities. Animal breeding is anticipated to have a contributing role, along with "dark niches and cracks in the ground providing suitable habitat for the sandflies"⁸. Currently to estimate the accurate prevalence of CL in the province is difficult as the hospitals and clinics function as significant teaching/tertiary care institutions not only treats the individuals of the province, but also include the people belonging to bordering parts of Afghanistan¹². Therefore, in the current review original articles highlighting the burden of CL published in last five years from various areas of Baluchistan will be discussed.

A study published in 2020 by Aisha Khan et al., reported the age and gender wise distribution of the CL along with the frequent site of the lesions. The study was carried out from 2018 -2019 throughout the province Baluchistan with 4072 included participants. According to their results they found 969/4072 (23.79%) cases in year 2018 (August to December) followed by 3103/4072 (76.20%) in year 2019 (January to December). Furthermore, children of age between 0-9 were seemed to be highly affected (39.6%) by the CL in both the years and male predominance was observed in both the years i.e. 55.7%. Additionally according to their analysis hands were the most specific site of infection (24.4%) followed by different areas of face¹³.

Darshna Kumari et al., in 2022 reported the infection from three coastal areas of province Baluchistan. They included 250 patients from Winder, Uthal and Lasbela areas, according to them there was male predominance and middle age group 11-40 years were suffered the disease more. The lesions were measured and the size of lesion were recorded to be approximately 3cm. Furthermore, they divided the lesion on the basis of wet ulcerative lesion and these were found to be highest in Winder (70%) and dry modular and popular lesion were recorded high in number in Uthal (50%)¹⁴.

Another study conducted by Farzad Khan et al., in Quetta Baluchistan revealed that there were 5032 positive cases of cutaneous leishmaniasis among them 2819 (56.02%) were male and 2213 (43.97%) were female from Quetta. Their results highlighted the trend of male predominance. As per their documentation 2,595,492 was the reported population of district Quetta according to census 2023 however, the cases were recruited from hospital and the sample was comprised of

only positive cases hence the findings are not reflecting the actual prevalence rather reporting the frequency of cases reported at various hospitals of district Quetta¹⁵.

A research on the prevalence and risk factors of cutaneous leishmaniasis was carried out in the district of Muslim Bagh (Killa-Saifullah), Baluchistan, Pakistan, among the 207 inhabitants of the various villages 192 people (92.75%) tested positive for CL, there were 64% male and 36% females (n = 69) infected individuals. Contrast to the findings of Aisha khan et., al. 2021 they reported high frequency of age group ranging from 20-29 years and the most frequent site of infection was seemed to be the legs^{13,16}.

In a case-control research published in 2021 data related to CL was evaluated they reported the same findings as of reported in following years by various researchers. The additional information that they highlighted was increase in cases during summer and decrease in winter season. They mentioned that the activity of sand fly becomes more frequent during monsoon compared to other seasons hence precautionary measures can be taken to reduce the disease burden^{10,17}.

The spread of CL is not limited to only Baluchistan different studies conducted in various areas of Pakistan has documented different pattern of spread. They have corelated it with low socio-economic status, poor life style, illiteracy and unawareness. Areas surrounded by grounds and less agricultural land have been discussed as one of the risk factors because majority of cases have been reported to have that habitat. Additionally, male predominance and school going age have been documented as population at risk which is showing that the sand fly hunts more in open spaces. That also emphasize on implementation of prevention and control program through special community based sanitation or through spray campaigns during the peak seasons¹⁸⁻²¹.

The studies reporting the various frequencies of CL in different parts of Baluchistan are highlighting the importance of the issue that the disease spread is not limited to coastal or border areas however the disease has penetrated in the cities and currently thousands of people are suffering the trauma of being infected and ill. In Pakistan considering leishmaniasis a vector born disease it has been included with malaria control program which is not justified as the sand fly share different morphological and pathological factors as well as their virulence is different therefore till date no any conclusive results have been obtained and disease is rising day by day^{22,23}. The rise in cases of these areas of Baluchistan has been attributed to poor life style,

unawareness about the disease course, expensive treatment, lost of followup due to commute issues in the population and lack of support from Government in prevention of this endemic²⁴⁻²⁶.

To reduce the prevalence of CL, it is crucial to educate the public and reduce stigma associated with the condition. Conduct monthly or quarterly community-based outreach programs in disease-endemic regions to educate populations on risk factors and infection prevention. Involving important influencers and elders in a community-focused intervention can significantly improve its success rate. Educating individuals on eliminating stigma around dermatological symptoms like CL might be included in such initiatives^{27, 28}. Lady health workers (LHWs) can assist women who are unable to attend community sessions. Local health workers (LHWs) provide health education to women and the public in their homes²⁹.

CL therapy should be straightforward and economical. Currently, therapy involves injecting medication directly into the lesion and typically lasts 4-8 weeks. More cheap and minimally intrusive therapeutic solutions are urgently required³⁰. To address the endemic nature of this disease in Balochistan, the government should acquire medications through the Provincial Medical Store Depot on an annual basis, ensuring that each district receives a fair portion based on the number of patients. This should also be included in the yearly budget.

Conclusion:

The reported study in last 5 years have demonstrated scattered prevalence of the CL in the province Baluchistan. Every district has shown diverse course of disease and different data. Though the consistent thing identified was the trend of increase in the incidence and prevalence across the province. The male gender and first 3 decades of life has been reported as the high-risk group. The most frequent sites of the lesions were reported to be the arms and legs. Furthermore, it was highlighted that the improvement in life style, education for the preventive measures and reduction in the stigma associated with CL needs to be emphasized through long term interventions and considering the emergence of new cases the CL should be dealt as a separate health issue and must not be merged with other health programs running in the country.

References:

1. Mock DJ, Hollenbaugh JA, Daddacha W, Overstreet MG, Lazarski CA, Fowell DJ, et al. Leishmania induces survival, proliferation and elevated cellular dNTP levels in human monocytes promoting acceleration of HIV co-infection. *PLoS pathogens*. 2012;8(4):e1002635.

2. Isenring E, Fehr J, Gültekin N, Schlagenhaut P. Infectious disease profiles of Syrian and Eritrean migrants presenting in Europe: a systematic review. *Travel medicine and infectious disease*. 2018;25:65-76.
3. Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R. Leishmaniasis: a review. *F1000Research*. 2017;6.
4. Gurel MS, Tekin B, Uzun S. Cutaneous leishmaniasis: A great imitator. *Clinics in dermatology*. 2020;38(2):140-51.
5. Kassi M, Kasi P. Diagnosis of cutaneous leishmaniasis by fine needle aspiration cytology: a report of 66 cases. *Tropical Doctor*. 2005;35(1):50-1.
6. Kassi M, Kassi M, Afghan AK, Rehman R, Kasi PM. Marring leishmaniasis: the stigmatization and the impact of cutaneous leishmaniasis in Pakistan and Afghanistan. *PLoS neglected tropical diseases*. 2008;2(10):e259.
7. Postigo JAR. Leishmaniasis in the world health organization eastern mediterranean region. *International journal of antimicrobial agents*. 2010;36:S62-S5.
8. Firdous R, Yasinzi M, Ranja K. Efficacy of glucantime in the treatment of Old World cutaneous leishmaniasis. *International journal of dermatology*. 2009;48(7):758-62.
9. Ejaz A, Raza N, Iftikhar N. Recurrent cutaneous leishmaniasis presenting as sporotrichoid abscesses: a rare presentation near Afghanistan border. *Dermatology Online Journal*. 2007;13(2).
10. Kayani B, Sadiq S, Rashid HB, Ahmed N, Mahmood A, Khaliq MS, et al. Cutaneous Leishmaniasis in Pakistan: a neglected disease needing one health strategy. *BMC Infectious Diseases*. 2021;21:1-10.
11. Katakura K. Molecular epidemiology of leishmaniasis in Asia (focus on cutaneous infections). *Current Opinion in Infectious Diseases*. 2009;22(2):126-30.
12. Afghan AK, Kassi M, Kasi PM, Ayub A, Kakar N, Marri SM. Clinical manifestations and distribution of cutaneous leishmaniasis in Pakistan. *Journal of tropical medicine*. 2011;2011(1):359145.
13. Khan A, Sajid R, Gul S, Hussain A, Zehri MT, Naz S, et al. Epidemiological and pathological characteristics of Cutaneous Leishmaniasis from Baluchistan Province of Pakistan. *Parasitology*. 2021;148(5):591-7.
14. Kumari D, Tariq H, Qadir MA, Imran I, Farooq G, Nasimullah M, et al. Incidence of Cutaneous Leishmaniasis in Coastal Areas (Winder, Uthal And Lasbela) of Balochistan Pakistan. *Pak-Euro Journal of Medical and Life Sciences*. 2022;5(2):405-10.
15. Kasi FK, Ahmed H, Khan N, Ali S, Ullah S, Kumar C, et al. Ratio of Incidence of Cutaneous Leishmaniasis and Site of Lesions in Different Age Groups of Patients of District Quetta, Balochistan, Pakistan. *Pak-Euro Journal of Medical and Life Sciences*. 2023;6(4):515-26.
16. Ullah Z, Ullah S, Muhammad B, Khan J, Shah G. Prevalence and Risk Factors of Cutaneous Leishmaniasis in Tehsil Muslim Bagh District Killa-Saifullah, Balochistan, Pakistan. *Pakistan's Multidisciplinary Journal for Arts & Science*. 2024:26-40.
17. Siriwardana Y, Zhou G, Deepachandi B, Akarawita J, Wickremarathne C, Warnasuriya W, et al. Trends in recently emerged *Leishmania donovani* induced cutaneous leishmaniasis, Sri Lanka, for the first 13 years. *BioMed Research International*. 2019;2019(1):4093603.
18. Ahmad S, Obaid MK, Taimur M, Shaheen H, Khan SN, Niaz S, et al. Knowledge, attitude, and practices towards cutaneous leishmaniasis in referral cases with cutaneous lesions: A cross-sectional survey in remote districts of southern Khyber Pakhtunkhwa, Pakistan. *Plos one*. 2022;17(5):e0268801.
19. Ayaz MM, Nazir MM, Ullah N, Zaman A, Akbar A, Zeeshan M, et al. Cutaneous leishmaniasis in the metropolitan City of Multan, Pakistan, a neglected tropical disease. *Journal of medical entomology*. 2018;55(4):1040-2.
20. Hussain M, Munir S, Ayaz S, Khattak BU, Khan TA, Muhammad N, et al. First report on molecular characterization of *Leishmania* species from cutaneous leishmaniasis patients in southern Khyber Pakhtunkhwa province of Pakistan. *Asian Pacific journal of tropical medicine*. 2017;10(7):718-21.

21. Mashhood AA, Malik SS, Haq MMU, Khan QU. Awareness and practice of preventive strategies in patients of cutaneous leishmaniasis. *Pakistan Armed Forces Medical Journal*. 2018;68(6):1683-85.
22. Haq RU, Saeed A, Lashari WA, Ahmed N, Ramzan T, Kannar A. Psychological and social burden of Cutaneous Leishmaniasis among the residents of Quetta city, Pakistan. *Cambridge Medicine Journal*. 2021.
23. Zeb I, Ali A, Nawab J, Khan MQ, Kamil A, Tsai K-H. Cutaneous leishmaniasis in male schoolchildren in the upper and lower Dir districts of Khyber Pakhtunkhwa, and a review of previous record in Pakistan. *ActaTropica*. 2020;209:105578.
24. Ahmad Z, Fatima SB, Masood N, Ullah F, Kamal S, Farooqi A. Spatial distribution of drinking and irrigation water quality in different climatic zones of Baluchistan, Pakistan. *Environmental Science and Pollution Research*. 2023;30(47):103801-22.
25. Irum S, Aftab M, Khan A, Naz S, Simsek S, Habib A, et al. Cutaneous leishmaniasis (CL): a cross-sectional community based survey on knowledge, attitude and practices in a highly endemic area of Waziristan (KPK Province), Pakistan. *ActaTropica*. 2021;213:105746.
26. Wijnant G-J, Dumetz F, Dirx L, Bulté D, Cuypers B, Van Bocxlaer K, et al. Tackling drug resistance and other causes of treatment failure in leishmaniasis. *Frontiers in Tropical Diseases*. 2022;3:837460.
27. Desjeux P. Global control and Leishmania HIV co-infection. *Clinics in dermatology*. 1999;17(3):317-25.
28. Bautista-Gomez MM, Doerfler J, del Mar Castro M. Barriers to cutaneous leishmaniasis care faced by indigenous communities of rural areas in Colombia: a qualitative study. *BMC Infectious Diseases*. 2022;22(1):302.
29. Polidano K, Parton L, Agampodi SB, Agampodi TC, Haileselassie BH, Lalani JM, et al. Community engagement in cutaneous leishmaniasis research in Brazil, Ethiopia, and Sri Lanka: a decolonial approach for global health. *Frontiers in Public Health*. 2022;10:823844.
30. Ramdas S. Cruel disease, cruel medicine: self-treatment of cutaneous leishmaniasis with harmful chemical substances in Suriname. *Social Science & Medicine*. 2012;75(6):1097-105.