

## **The interaction effect of nitrogen levels on growth characteristic of wheat crop (*Triticum aestivum* L.)**

### **ABSTRACT**

The present research work was carried at the Agronomy Farm, Department of Agronomy, Narendra Deva University of Agriculture and Technology, Narendra Nagar (Kumarganj) Faizabad (U.P.), during Rabi season of 2014-2015. This farm is located at a distance of 42 Km. from Faizabad district headquarters at Faizabad – Raibareilly road. The ANOVA of the research trail used a design: randomized block design (factorial) to evaluate growth traits, comprising of initial plant population, Plant height (cm), Number of shoots (m-2). The data indeed suggested that initial plant population was not significantly influenced by either variety levels or Nitrogen levels. Contrastingly, the data showed that Variety levels and Nitrogen levels significantly affected plant height at 30, 60, 90 days after sowing (DAS) and at harvest. The number of shoots 30 DAS was not affected significantly by Variety levels and Nitrogen levels in the data. In addition, Variety levels showed a significant effect on the shoot number at 60 and 90 DAS and also at harvest stage, while Nitrogen levels did not show significant effects in the data.

**Keys words:** Varieties, Nitrogen Levels, F-RBD, Wheat crop

### **INTRODUCTION**

Wheat, *Triticum aestivum* L., Family Poaceae, is one of the world's most important staple foods. Wheat is an exotic but ancient crop that has been grown in India since pre-historic times. Any crop has a social and cultural heritage, particularly so if it is a staple. Cultural heritage is transmitted from generation to generation. It has been constantly recreated by communities and groups in reaction to their environment, their relationship with nature, and their history. India occupies a prime position among the world's major wheat-producing and consuming countries. It is the second most important crop in Indian agriculture after rice. It is a dominant crop in north-western Indian states; the vast Indo-Gangetic plains are the most suitable providers of climate for it. This Indian region is described as „wheat bowl of India“. India is the 4<sup>th</sup> major producing country and also by area, it ranks 4<sup>th</sup> in the **world (FAO,1982)**. The cumulative impact of crop introduction in the long past has affected our habits to an extent that majority of our food sources we utilise today owe their origin too far off countries or regions outside India. About 55% of the world population uses wheat for intake of about 20% of food calories.

Wheat is cultivated worldwide on an area of 220.88 million hectares with a production of 725.47 million tonnes and productivity of 3.28 tonnes per hectares. In India, it is grown in an area of 30.47 million hectares with a production of 95.85 million tonnes and productivity of 3.15 tonnes per hectare. Six states, viz., U.P., Punjab, Haryana, M.P., Rajasthan and Bihar, contribute about 91% towards the total wheat production. Among them, Uttar Pradesh ranks first in respect of both area and production. It is followed by Punjab and Haryana. It covers an area of 9.2 mha with a production of 24.50 mt and productivity of 2.7 tonnes per hectare. However, its productivity is comparatively low as compared to the states of Punjab and Haryana. In the case of continuation of rice-wheat cropping system, ultimately with imbalanced fertilization, delay in wheat sowing acts as the main cause for low productivity.

Due to several self-evident reasons, wheat area sown in late has increased considerably in U.P. from past few years. In the state of U.P., rice-wheat is the most common cropping system followed by most of the farmers. By **Patel et.al.2012** Delayed transplanting of rice due to delayed monsoon, wider use of long duration varieties and heavy rainfall at the later stages of crop growth are the main causes responsible for wheat sown in late. Besides the proceeding crop like sugarcane, potato and toria also leave the fields very late. Under late sown condition wheat produces poor yield because of the lesser exploitation of the potentialities of the crop. Primary reductions in productivity occur due to delayed emergence of the seedling and assuredly the growth and development period of the crop by **annual report (2014-15)**. Delayed emergence of the crop followed by high temperature and hot desiccating winds during grain filling stages result in forcing of maturity in the case of late-sown wheat due to dehydration, finally resulting in a heavy reduction in the whole biomass and yield. The purpose of our research was to increase crop productivity without degrading soil fertility. varieties and nitrogen levels were used to restore soil fertility and nutrient availability, while also reducing environmental impact by **Maqsood et.at.(2012)**.

## **MATERIAL AND METHODS**

Field experiment was conducted at Agronomy Research Farm, Narendra Deva University of Agriculture and Technology, Narendra Nagar (Kumarganj) Faizabad (U.P.), during Rabi season of 2014-2015. This farm is located at an elevation of 42 Km. from Faizabad districts headquarter at Faizabad – Raibareilly road. It falls in the sub-tropical zone of Indogangatic plains. Experimental site lies between 26°47' North latitude, 82°12' East longitude at an attitude of about 113.0 meter from mean sea level and is subjected to extremes of weather conditions.

### **Experimental details:**

Material of experiment arranged in the randomized block design with four varieties PBW-373, HD-2327, and NW-1014 and six nitrogen levels (0, 40, 80, 120, 160, and 200 kg N ha<sup>-1</sup>) with three replications.

### **Details of Layout:**

Design: Randomized Block Design (Factorial), Replication : 3, Varieties : 3, Nitrogen levels: 6, Treatment combinations:18, Total no. of plots : 54 Gross plot size : 4 x5 = 20 m<sup>2</sup>, Net plot size : 3.2 x 4 = 12.8 m<sup>2</sup>, Row to row distance : 20 cm, Main irrigation channel : 1.5 m, Sub irrigation channel : 1 m, Replication border (path) : 1 m, Bunds: 0.5 m, Field border : 1 m. The traits monitored in the treatment were initial plant population (m<sup>-2</sup>), Plant height (cm), Number of shoots (m<sup>-2</sup>). On these traits, data recorded were statistically analyzed using the analysis of variance (ANOVA) technique as outlined by Gomez and Gomez (1992). The critical difference at 5 % level of probability was used to estimate the importance of variations between treatment means wherever F-test was significant (**Chandel et al., 1998**).

## **RESULTS AND DISCUSSION**

The initial plant population under different treatments based on varieties and nitrogen levels practices for the years 2014-2015 along with Table 1. and Fig.1.

**Varieties and Nitrogen levels:** For Varieties of the highest initial plant population Variety (V<sub>1</sub>) NW-1014 in the data was recorded at 140.00 this is followed by Variety (V<sub>2</sub>) HD-2327 reporting as 139.00 the lowest population was recorded in Variety (V<sub>3</sub>) PBW-373 reporting as 135.00. Nitrogen though, in the highest initiation plant population was seen in N<sub>4</sub> at 160kg/ha this was followed by 140.13, in N<sub>5</sub> that was at 200kg/ha with a recorded value of 138.69. The smallest

population was found in N<sub>0</sub>, which the control had a value of 138.14 closed result recorded by Prashanth *et.al.*(2022).

### **Plant height (cm) at 30 DAS**

#### **Varieties and Nitrogen levels:**

Table-1 and fig-2. Showed that Varieties that had the highest plant height at 30 days in (V<sub>3</sub>) NW-1014 data was that with 16.69 cm. (V<sub>2</sub>) HD-2327 had a plant height of 15.80 cm. The least plant height was recorded in (V<sub>3</sub>) PBW-373 with 15.54 cm this is as virtually the same as findings by Patel *et al.*, (2012) and Swelamet.*al.* (2008). The highest plant height at 30 days in data was recorded for N<sub>5</sub> (200kg/ha) with 18.25cm, while plant height of 17.17 cm was recorded for N<sub>4</sub> (160kg/ha). The lowest plant height was recorded in the control group N<sub>0</sub> with 14.40 cm by Fadhil *et.al.*(2022).

### **Plant height at 60 DAS**

Varieties and Nitrogen levels: showed that Variety level highest plant height at 60 days in data was for (V<sub>1</sub>) NW-1014 with 64.66 cm. (V<sub>2</sub>) PBW-373 had a plant height of 59.78 cm. The lowest plant height was recorded for (V<sub>3</sub>) HD-2327 with 58.56 cm. whereas in Nitrogen levels the highest plant height at 60 days data was for N<sub>5</sub> (200kg/ha) with 66.51 cm. N<sub>4</sub>(160 kg ha as a basal dose) had a recorded plant height at 65.88 cm. The lowest plant height was observed in the control N<sub>0</sub> at 53.68 cm, the same result was reported by Singh *et al.*, (2023) and Kumar *et.al.*(2023).

### **Plant height (cm) at 90 DAS**

**Varieties and Nitrogen levels:** Obtained the tallest plant at 90 days in (V<sub>3</sub>) NW-1014 data with 87.98 cm. The plant height in (V<sub>2</sub>) HD-2327 was 81.34 cm. The shortest plant was observed in (V<sub>3</sub>) PBW-373 with 79.68 cm, which was in line with Tripathi *et al.*,(2003), and Salanturet.*al.*(2006). In Nitrogen levels, the highest plant height at 90 days in data was for N<sub>5</sub> (200kg/ha) with 90.52cm. N<sub>4</sub> (160kg/ha) recorded a plant height of 89.64 cm. The minimum plant height was recorded from the control group N<sub>0</sub> with a plant height of 73.04 cm.

### **Plant height (cm) at Harvest Stage**

**Varieties and Nitrogen levels:** recorded Variety highest plant height at harvest stage in data was for V<sub>1</sub> NW- 1014 with 88.48 cm. (V<sub>2</sub>) PBW - 373 was recorded with a plant height of 81.84 cm. The lowest plant height was recorded for (V<sub>3</sub>) HD - 2327 with 80.18 cm. whereas in Nitrogen levels the highest plant height at harvest stage data was for N<sub>5</sub>(200kg/ha)with 91.20 cm. N<sub>4</sub>(160kg/ha as a basal dose) was observed with a plant height of 90.14 cm. The lowest plant height was obtained with control N<sub>0</sub>group, which was 73.54 cm similar result that was reported by Shekoofaet.*al.*(2008) and Maqsood *et.al.* (2012).

### **Number of shoots (m<sup>-2</sup>) at 30 DAS**

**Varieties and Nitrogen levels:** Table-1 and Fig 3. At the Variety, the highest Number of shoots (m<sup>-2</sup>) at 30 DAS stage in Varieties level table-1 indicated that the highest Number of shoots (m<sup>-2</sup>) in 30 DAS data was recorded for V<sub>1</sub> (NW-1014) with 182.00 m<sup>-2</sup>. V<sub>2</sub> (PBW-373) had a Number of shoots 180.70 m<sup>-2</sup>. The Number of shoots was in V<sub>3</sub> (HD-2327) with 175.50(m<sup>-2</sup>). Where recorded that the highest number of shoots (m<sup>-2</sup>) data was encountered for N<sub>5</sub> (200kg/ha) with 182.19 m<sup>-2</sup>, whereas N<sub>4</sub> (160 kg/ha as a basal dose) recorded the number of shoots with 182.17

m<sup>-2</sup>. The lowest number of shoots was found to be in the control group, N<sub>0</sub>, having 179.58 m<sup>-2</sup> closed finding reported by **Kumar *et.al.*(2020) andsatishet.al. (2022).**

#### **Number of shoots (m<sup>-2</sup>) at 60 DAS**

**Varieties and Nitrogen levels:**Reported that Variety the highest Number of shootsat60 DAS stage in data was for (V<sub>1</sub>) NW-1014 with 362.95 m<sup>-2</sup>.(V<sub>2</sub>) PBW-373 had a Number of shoots 356.59m<sup>-2</sup>.The lowest Number of shootswas recorded for (V<sub>3</sub>)HD-2327 with 329.26 m<sup>-2</sup>. Where as in Nitrogen levelsthe highest Number of shootsat 60 DAS stage data was for N<sub>5</sub> (200kg/ha) with 378.50 m<sup>-2</sup>.N<sub>4</sub> (160kg/ha as a basal dose) recorded a Number of shootsof 377.57 m<sup>-2</sup>.The lowest Number of shootswas observed in the control group N<sub>0</sub>with 307.65 m<sup>-2</sup>similar result reported by**Kumar *et.al.*(2023)and Dagashet.al (2014)**

#### **Number of shoots (m<sup>-2</sup>) at 90 DAS**

**Varieties and Nitrogen levels:** showed that Variety the highest Number of shootsat90 DAS stage in data was for (V<sub>1</sub>) NW-1014 with 387.30 m<sup>-2</sup>.(V<sub>2</sub>) PBW-373 had a Number of shoots 379.05 m<sup>-2</sup>.The lowest Number of shootswas recorded for (V<sub>3</sub>)HD-2327 with 350.06 m<sup>-2</sup>. Where as in Nitrogen levelsthe highest Number of shootsat 90 DAS stage data was for N<sub>5</sub> (200kg/ha) with 403.60 m<sup>-2</sup>.N<sub>4</sub> (160kg/ha as a basal dose) recorded a Number of shootsof 402.19 m<sup>-2</sup>.The lowest Number of shootswas observed in the control group N<sub>0</sub>with327.71m<sup>-2</sup>similar result reported by**Hussain *et.al.*(2017) and Sikarwar *et.al.* (2018).**

#### **Number of shoots (m<sup>-2</sup>) at harvest Stage**


**Varieties and Nitrogen levels:** estimated that Variety the highest Number of shoots at harvest stage in data was for (V<sub>1</sub>) NW-1014 with 383.42 m<sup>-2</sup>.As well as(V<sub>2</sub>) PBW-373 had a Number of shoots of 376.05 m<sup>-2</sup>.The lowest Number of shoots was recorded for (V<sub>3</sub>)HD-2327 with346.56 m<sup>-2</sup>. Where as in Nitrogen levelsthe highest Number of shoots at harvest stage data was for N<sub>5</sub> (200kg/ha) with 399.80 m<sup>-2</sup>.N<sub>4</sub> (160kg/ha as a basal dose) recorded a Number of shoots of 398.17 m<sup>-2</sup>.The lowest Number of shoots was observed in the control group N<sub>0</sub>with 324.43 m<sup>-2</sup>similar result reported by**Farooq *et.al.*(2018) and Prashanth *et al.*(2022)**


**Table 1: Effect of different treatments on initial plant population (m<sup>-2</sup>), Plant height (cm) and Number of shoots (m<sup>-2</sup>)**

Treatments	Initial plant population (m <sup>-2</sup> ) 20 DAS	Plant height (cm)				Number of shoots (m <sup>-2</sup> )			
		30 DAS	60 DAS	90 DAS	At harvest	30 DAS	60 DAS	90 DAS	At harvest
Varieties									


<b>PBW-373</b>	135.00	15.54	58.56	79.68	80.18	175.50	329.26	350.06	346.56
<b>HD-2327</b>	139.00	15.80	59.78	81.34	81.84	180.70	356.59	379.05	376.05
<b>NW-1014</b>	140.00	16.69	64.66	87.98	88.48	182.00	362.95	387.30	383.42
<b>SEm±</b>	1.80	0.21	0.86	1.13	1.07	2.45	4.93	5.11	5.06
<b>CD (P=0.05)</b>	NS	0.62	2.49	3.26	3.08	NS	14.18	14.69	14.54
Nitrogen levels( kg ha <sup>-1</sup> )									
<b>0</b>	138.14	14.40	53.68	73.04	73.54	179.58	307.65	327.71	324.43
<b>40</b>	136.62	15.07	57.34	78.02	78.52	177.61	328.62	351.16	347.46
<b>80</b>	136.10	15.77	61.00	83.00	83.50	176.93	349.60	372.40	368.68
<b>120</b>	138.32	16.70	63.44	86.32	86.82	179.82	363.58	388.30	385.42
<b>160</b>	140.13	17.17	65.88	89.64	90.14	182.17	377.57	402.19	398.17
<b>200</b>	138.69	18.25	66.51	90.52	91.20	182.19	378.50	403.60	399.80
<b>SEm±</b>	2.54	0.30	1.22	1.60	1.51	3.47	6.97	7.23	7.15
<b>CD (P=0.05)</b>	NS	0.88	3.53	4.62	4.35	NS	20.05	20.78	20.57

Whereas variety-(V<sub>1</sub>, V<sub>2</sub>, V<sub>3</sub>) and Nitrogen levels- N<sub>0</sub> N<sub>1</sub>, N<sub>2</sub>, N<sub>3</sub>, N<sub>4</sub>, N<sub>5</sub>





UNDER PEE



## CONCLUSION

The experiment concluded that for the initial plant population, plant height (cm) and Number of shoots /m<sup>2</sup> the best results were obtained under the Variety (V<sub>3</sub>) NW-1014. Among the Nitrogen levels(kgha<sup>-1</sup>) (N<sub>5</sub>) practices (200 kg/ha), recorded the highest values. The lowest values were observed under the Variety (V<sub>2</sub>)HD-2327 and Nitrogen level (N<sub>0</sub>) the control levels.

## REFERENCE

Maqsood, M., Shehzad, M. A., Asim, A., & Ahmad, W. (2012). Optimizing rate of nitrogen application for higher growth and yield of wheat (*Triticum aestivum* L.) cultivars. *Pakistan Journal of Agricultural Science*, 49(4), 491-496.

Shekoofa, A., & Emam, Y. (2008). Effects of nitrogen fertilization and plant growth regulators (PGRs) on yield of wheat (*Triticum aestivum* L.) cv. Shiraz. *Journal of Agricultural Science and Technology*, 10(2), 101-108.

Tripathi, S. C., Sayre, K. D., Kaul, J. N., & Narang, R. S. (2003). Growth and morphology of spring wheat (*Triticum aestivum* L.) culms and their association with lodging: effects of genotypes, N levels and ethephon. *Field crops research*, 84(3), 271-290.

Authority, V. B. (2015). Annual-Report 2014-15, Visva-Bharati, English Version.

Frevert, D. K., Hill, R. W., & Braaten, B. C. (1982). Estimation of FAO evapotranspiration coefficients. *Journal of Irrigation and Drainage Engineering*, 109(2), 265-270.

Salantur, A., Ozturk, A., & Akten, S. (2006). Growth and yield response of spring wheat (*Triticum aestivum* L.) to inoculation with rhizobacteria. *Plant Soil and Environment*, 52(3), 111.

Kumar, A., Yadav, D. D., Patel, V., Siddique, M. Z., Kumar, S., Singh, P. A., (2020) Combined effect of bio-fertilizer and micronutrients on fertility, growth and productivity of chickpea. *Int J Chem Stud*, 8(6), 2576-2579.

Kumar, A., Nayak, K., Kumar, K., Kumhare, A., & Tekam, Y. (2023). Effect on Yield, Attribute Character and Economics of Various Treatment in Wheat (*Triticum aestivum* L.) Crop. *International Journal of Plant & Soil Science*, 35(21), 823-826.

Kumar, A., Patel, V., Kumar, K., & Kumhare, A. (2023). Varietal Effect of Different Nitrogen Levels on Yield Attributing Characters and Quality Parameters of Wheat (*Triticum aestivum* L.) Crop. *International Journal of Environment and Climate Change*, 13(12), 246-249.

Satish, R., Dawson, J., Umesha, C., Reddy, S. M., Girish, M., & Prasanth, S. (2022). Performance of Varieties on Growth and Yield of Wheat (*Triticum aestivum* L.) Under Prayagraj Condition. *International Journal of Environment and Climate Change*, 12(10), 967-971.

Hussain, M. A., Dohuki, M. S., & Ameen, H. A. (2017). Response of some bread wheat (*Triticum aestivum* L.) cultivars to nitrogen levels. *Kufa Journal for Agricultural Sciences*, 9(4), 365-390.

Sikarwar, B. P. S., Gupta, D., & Singh, L. (2018). Effect of varying nitrogen levels and varieties on productivity and profitability of wheat (*triticum aestivum*) under rainfed condition. *Annals of Plant and Soil Research*, 20(1), 58-62.

Farooq, M., Khan, I., Ahmed, S., Ilyas, N., Saboor, A., Bakhtiar, M., ... & Khan, A. Y. (2018). Agronomical efficiency of two Wheat (*Triticum aestivum* L.) Varieties against different level of Nitrogen fertilizer in Subtropical region of Pakistan. *Int. J. Environ. Agric. Res*, 4(4), 28-36.

- Fadhil, A. H. (2020). Response of Wheat Cultivars of *Triticum aestivum* L to Nitrogen Fertilizer for Growth and Yield Traits. *Indian Journal of Ecology*, 47, 45-51.
- Patel, C. B., Singh, R. S., Yadav, M. K., Singh, S. K., Singh, M. K., Singh, K. K., & Mall, R. K. (2012). Response of different wheat (*Triticum aestivum* L. emend Fiori & Paol.) genotypes to various nitrogen levels under late sown conditions of Eastern Uttar Pradesh. *Environment & Ecology*, 30(3C), 1192-1196.
- Fernandez, G. C. (1992). Residual analysis and data transformations: important tools in statistical analysis. *HortScience*, 27(4), 297-300.
- SWELAM, A. A. A. (2008). Mean performance, correlation and path coefficient analysis for grain yield and its components of four bread wheat cultivars grown under four nitrogen levels. *Egyptian Journal of Agricultural Research*, 86(2), 561-573.
- Prashanth, S., Umesha, C., & Vikas, S. P. (2022). Performance of Wheat (*Triticum aestivum* L.) Genotypes their Growth and Yield under Irrigated Condition of Prayagraj. *International Journal of Environment and Climate Change*, 12(11), 1-7.
- Ullah, I., Ali, N., Durrani, S., Shabaz, M. A., Hafeez, A., Ameer, H., ... & Waheed, A. (2018). Effect of different nitrogen levels on growth, yield and yield contributing attributes of wheat. *Int J Sci Eng Res*, 9(9), 595-602.
- Sultana, S. R., Ahmad, A., Wajid, A., & Akhtar, J. (2013). Estimating growth and yield related traits of wheat genotypes under variable nitrogen application in semi-arid conditions. *Pakistan Journal of Life and Social Sciences*, 11(2), 118-125.
- Dagash, Y. M. I., Ahmed, I. S., & Khalil, N. A. (2014). Effect of nitrogen fertilization, sowing methods and sowing dates on yield and yield attributes of wheat (*Triticum aestivum* L). *Universal Journal of Plant Science*, 2(6), 108-113.
- Chandel, N. S., Maltepe, E., Goldwasser, E., Mathieu, C. E., Simon, M. C., & Schumacker, P. (1998). Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proceedings of the National Academy of Sciences*, 95(20), 11715-11720.