

## Effect of Solid State Fermentation on Phytochemical Composition of Rice Bran Meal

### ABSTRACT

In aquaculture, feed ingredients are the course of skyrocketing the cost of production which constitutes about 60 to 75% of the total operational costs of a fish production. Some of the feed ingredients used are stable food for man which are highly competitive and expensive. Consequently, farmers use locally available fish feed ingredients including agricultural by-products. These feed ingredients have led to increased pond productivity due to their nutrients constituent and bio-availability. This study evaluation the effect of solid state fermentation on phytochemical compositions of rice bran meal. Rice bran was collected in three different containers, ground into powder, then sieved and fermented using solid state fermentation procedure. Each sample was solidly fermented (wet at 10% moisture and keep in fermenter at ambient temperature) for 24 hours (1 day), 96 hours (4 days), and 168 hours (7 days). Afterward, the fermented samples were oven dried for one hour at 100°C as described. The fermented samples were taken to the laboratory for phytochemical analysis. The results obtained from this study revealed that fermenting rice bran for seven days (168hrs) using solid state fermentation method reduced the Alkaloid contain from  $12.61 \pm 1.04$  to  $8.13 \pm 0.16$ , Phytate from  $6.73 \pm 1.32$  to  $2.15 \pm 0.73$ , Tannins from  $4.60 \pm 0.13$  to  $1.52 \pm 0.24$ , Saponins from  $2.98 \pm 0.54$  to  $1.43 \pm 1.17$  and Oxalate from  $2.78 \pm 1.25$  to  $1.82 \pm 0.13$ . However, it increases Flavonoids from  $6.94 \pm 1.82$  to  $11.18 \pm 1.35$  and Phenols from  $3.58 \pm 1.65$  to  $1.71 \pm 0.32$ . This study demonstrates that, fermented rice bran meal can be incorporated as an essential part of the feed production in order to reduced cost of production thereby increasing the profit and enhances waste management.

**Keywords:** RiceBran Meal; Solid-State Fermentation and Phytochemical Composition.

**Comment [S1]:** Data for 1 day and 4 days are missing here. Compare the data to see the days of fermentation on phytochemical composition.

## 1. INTRODUCTION

The high cost of fish feed has been recognized as a major factor militating against rapid development of aquaculture in the developing countries due to the fact that most of the conventional feedstuffs being used in human foods and animal feeds hence leading to their exorbitant prices and scarcity [20]. However, problem of high cost of feeding in aquaculture is further exacerbated due to the scarce and expensive nature of some of the ingredients used in the formulation of fish feeds. In order to solve the problem of scarce and expensive feed ingredients, a number of non-conventional feedstuffs have been investigated most of which are alternative protein sources since this nutrient is considered as the most expensive nutrient [18].

Agro by-products are used as a fish feed ingredients which are less scarce and inexpensive and they are derived from the industry due to processing of main products in large quantity. They are less fibrous, more concentrated, highly nutritious, and less expensive as compared to crop residues which are crop-waste residues mainly consist of materials remaining after harvesting and processing, such as straw, husk or stubble from barley, wheat, rice, beans, oats, rye stalks, or stovers from corn, sorghum, cotton among others [8]. Maize has been a traditional energy source in formulated feed but rising cost and accompanying scarcity is making it increasingly uneconomical to include it in animal feeds. Therefore, there is need for the recruitment of other suitable ingredients that can be used as dietary energy source in the replacement of maize [13]. The need to solve the problem of feeding in aquaculture has already been demonstrated through various researches in the utilization of agricultural waste such as Cassava peel [15; 6], Sweet potato peel [14], Poultry offal [7], water hyacinth meal [19], Maize-cob meal [17], fermented sorghum [10] and fermented sorghum by-product [12] to mention a few.

Among the alternative of the local feed ingredients that can be used to feed fish are fermented agro by-products.

Rice is one of the cereals besides the major wheat, millet, and maize. Rice is a major food source for millions of people around the world. Rice is grown mostly in tropical areas under agricultural conditions in which major cereals fail to give substantial yields [1]. Rice is classified with maize, sorghum, and Coix (Job's tears) in the grass sub-family *Panicoideae*[1]. Rice is now getting popularity as it became a major source of energy and protein for millions of people in Africa.

This study investigate the effect of solid state fermentation on phytochemical composition of rice bran meal.

## **2. MATERIALS AND METHODS**

### **2.1 Experimental Site**

This research was conducted at the Nutrition lab of the Department of Fisheries Technology, School of Agriculture, Yobe State College of Agriculture, Science and Technology Gujba, Yobe State, Nigeria.

### **2.2 Source of Experimental Ingredients**

Rice bran was procured from feedstuffs market Damaturu, Yobe State.

### **2.3 Processing and Fermentation of Millet Bran**

Rice bran was collected in three different containers, ground into powder, then sieved and fermented using solid fermented procedure as described by [17]. Each sample was solidly fermented (wet at 10% moisture and keep in fermenter at ambient temperature) for 24 hours (1 day), 96 hours (4 days), and 168 hours (7 days). Afterward, the fermented samples were oven dried for one hour at 100°C as described. The fermented samples were taken to the laboratory for nutritional, phyto-chemical analysis.

### **2.4 Phytochemical Screening of Fermented Samples**

Two (2) grams each of the samples collected were used for quantitative phyto-chemical screening to assess Alkaloid, Cyanogens, Flavonoids, Glycosides, Phenols, Phytates, Oxalate, Saponin, Nitrite and Tannins on each of the treatments following the methods of [4;23].

### **2.5 Statistical Analysis**

All data generated are subjected to descriptive statistics to determine the mean values and then subjected to analysis of variance (ANOVA) at 95% probability level where the significant differences were detected. Means values were separated using Least Significant Difference (LSD). All data were analyzed using SPSS (Statistical Package for Social Sciences) version 20.0 Statistical Package.

## **3. RESULTS**

### **3.1 Phytochemical Composition of the Fermented Rice Bran Meal**

The quantitative investigation of the phytochemical constituents of the fermented Rice bran at various fermentation period shows that Alkaloid, Flavonoids, Phenols, Phytate, Oxalate, Saponin and Tannin are present in all the fermented bran meals at 24 hours (1 day), 96 hours (4 days) and

168 hours (7 days) fermentation are shown in table 1. The highest content of Alkaloids in the fermented rice bran was recorded in 0 hour (0 day) fermentation with  $12.61 \pm 1.04 \text{ mg/kg}$  and the lowest was recorded in 168 hours (7 day) fermentation with  $8.13 \pm 1.16$ . There is no significance ( $P < 0.05$ ) between 0 hour and day one. However, significance ( $P < 0.05$ ) was observed between day one and day seven. The highest content of Flavonoids was recorded in 168 hours (7 days) fermentation with  $11.18 \pm 1.35 \text{ mgRutin/kg}$  and the lowest was recorded in 0 hours (0 days) fermentation with  $6.94 \pm 1.82 \text{ mgRutin/kg}$ . There is no significance ( $P < 0.05$ ) between 0 hour and day one. However, there is significance ( $P < 0.05$ ) between 0 hour and day seven. The highest content of Phytate, Tannins, Saponins and phenols in rice bran were recorded in 0 hours (0 day) fermentation with  $6.73 \pm 1.32 \text{ mg/kg}$ ,  $4.60 \pm 0.13 \text{ mgAT/kg}$ ,  $2.98 \pm 0.54 \text{ g/kg}$  and  $3.58 \pm 1.65 \text{ gGAE/kg}$  respectively while the lowest were recorded in 168 hours (7 days) fermentation with  $2.15 \pm 0.73 \text{ mg/kg}$ ,  $1.52 \pm 0.24 \text{ mgAT/kg}$ ,  $1.43 \pm 1.17 \text{ g/kg}$  and  $1.71 \pm 0.32 \text{ gGAE/KG}$  respectively and highest content of oxalate was recorded in 0 hours (0 day) fermentation with  $2.78 \pm 1.25 \text{ mg}/100\text{g}$  and the lowest was recorded in 168 hours (7 days) fermentation with  $1.82 \pm 0.13 \text{ mg}/100\text{g}$ . There is significant difference ( $P < 0.05$ ) between day one day and day seven.

**Comment [S2]:** Write the result of day 1 and 4 also. You must include for each and every parameters in results for better understanding of reader.

Table 1: Quantitative Phytochemical Composition of Fermented Rice Bran Meal

Parameters	0Day Fermentation	1Day Fermentation	4Days Fermentation	7 Days Fermentation
Alkaloids (mg/kg)	12.61±1.04 <sup>a</sup>	10.24±1.52 <sup>b</sup>	9.18±0.11 <sup>c</sup>	8.12±1.16 <sup>c</sup>
Flavonoids (mgRutin/kg)	6.94±1.82 <sup>c</sup>	7.90±1.35 <sup>c</sup>	9.03±0.43 <sup>b</sup>	11.08±1.35 <sup>a</sup>
Phytate (mg/kg)	6.73±1.32 <sup>a</sup>	4.39±0.82 <sup>b</sup>	4.05±0.56 <sup>b</sup>	2.15±0.73 <sup>c</sup>
Tannins (mgTA/kg)	4.60±0.13 <sup>a</sup>	4.11±0.65 <sup>a</sup>	3.26±0.47 <sup>b</sup>	1.52±0.24 <sup>c</sup>
Saponins (g/kg)	2.98±0.54 <sup>a</sup>	2.50±0.17 <sup>a</sup>	1.88±1.54 <sup>a</sup>	1.43±1.17 <sup>a</sup>
Phenols (gGAE/kg)	3.58±1.65 <sup>a</sup>	2.91±1.53 <sup>a</sup>	2.14±0.12 <sup>a</sup>	1.71±0.32 <sup>b</sup>
Oxalate (mg∞/100g)	2.78±1.25 <sup>a</sup>	2.47±1.42 <sup>a</sup>	2.91±0.31 <sup>a</sup>	1.82±0.13 <sup>b</sup>

Means in the Same Row with Different Superscripts are Significantly Different (P<0.05)

#### 4. DISCUSSION

The phytochemical investigation of the fermented Rice bran at varying fermentation periods revealed that, the ingredients contains alkaloids, flavonoids, phenols, phytate, oxalate, saponins and tannins at varying concentrations as presented in the table [1](#). These chemical compounds are synthesized in natural food and / or feedstuffs by the normal metabolism of the plant species. The biochemical are also known as 'secondary metabolites' in plants and they have been shown to be highly biologically active [9]. The compounds which reduce feed intake and nutrients utilization of plants or plant products used for feeds and they play vital roles in determining the use of plants for fish feed [16]. More often than not, a single plant may contain two or more toxic compounds that add to the difficulties of detoxification. [9] who reported that, alkaloids are one of the largest groups of chemical compounds synthesized by plants and generally found as salts of plant acids such as oxalic, malic, tartaric or citric acid. They are small organic molecules, common to about 15 to 20 per cent of all vascular plants, usually comprising several carbon rings with side chains of one or more carbon atoms being replaced by nitrogen. They are considered to be anti-nutrients because of their action on the nervous system, disrupting or inappropriately augmenting electrochemical transmission. Consumption of high tropane alkaloids caused rapid heartbeat, paralysis and in severe case that lead to death. Uptake of high concentration of tryptamine alkaloids lead to staggering gate and death. Indeed, the physiological effects of alkaloids on fish are very evident. [8] who reported that flavonoids are diphenylpropanes that constitute one of the most characteristic classes of secondary metabolites in plants Flavonoids have been shown to be potent antioxidants, capable of scavenging hydroxyl radicals, superoxide anions and lipid peroxy radicals, and have been reported as having antibacterial, anti-inflammatory, antiallergic, antimutagenic, antiviral, antineoplastic, anti-thrombotic and vasodilatory actions [8;3]. Another one is oxalate which is an anti-nutrient which under normal conditions is confined to separate compartments. However, when it is processed and/or digested, it comes into contact with the nutrients in the gastrointestinal tract. When released, oxalic acid binds with nutrients, rendering them inaccessible to the body. Oxalic acid binds calcium and forms calcium oxalate which is insoluble. Calcium oxalate adversely affects the absorption and

utilization of calcium in the animal body [2]. If feed with excessive amounts of oxalic acid is consumed regularly, nutritional deficiencies are likely to occur, as well as severe irritation to the lining of the gut. Oxalic acid is of only minor significance as an anti-nutritive factor since microflora can readily metabolize soluble oxalates [9]. Phytate, is also a metabolites that is an inositol hexakisphosphate, containing compound that binds with minerals and inhibits mineral absorption. The cause of mineral deficiency is commonly due to its low bioavailability in the diet. The presence of phytate in fish feeds has been associated with reduced mineral absorption due to the structure of phytate which has high density of negatively charged phosphate groups which form very stable complexes with mineral ions causing non-availability for intestinal absorption [24]. Phytates are generally found in feed high in fibre especially in wheat bran, whole grains and legumes [21]. Phytic acid acts as a strong chelator, forming protein and mineral-phytic acid complexes; the net result being reduced protein and mineral bioavailability [11]. Phytic acid is reported to chelate metal ions such as calcium, magnesium, zinc, copper, iron and molybdenum to form insoluble complexes that are not readily absorbed from gastrointestinal tract. Phytic acid also inhibits the action of gastrointestinal tyrosinase, trypsin, pepsin, lipase and amylase [11]. This study is in contrast with [2] who reported decrease in tannins and phytate with increase fermentation period.

Similarly, Saponins reduce the uptake of glucose and cholesterol at the gut through intraluminal physicochemical interaction. Hence, it has been reported to have hypo cholesterolemic effects [22]. In fish saponins have been reported to reduce growth, feed efficiency and interferes with the absorption of dietary lipids and vitamins (A and E) [15]. It has been reported that saponins can affect animal performance and metabolism in a number of ways which include erythrocyte haemolysis, reduction of blood and liver cholesterol, depression of growth rate, enzyme inhibition and reduction in nutrient absorption [2]. Tannins may form a less digestible complex with dietary proteins and may bind and inhibit the endogenous protein such as digestive enzymes. The tannin-protein complexes are astringent and adversely affect feed intake and all plants contains phenolic compounds but their type and concentration may cause negative animal responses [12]. The concentration of condensed tannins above 4 per cent has been reported to be toxic to ruminants as they are more resistant to microbial attack and are harmful to a variety of microorganisms [24].

The results obtained in this study revealed that, the presence of alkaloids, flavonoids, phenols, phytates, oxalates, saponins and tannins in the fermented agro by-products. The concentrations of these metabolites were moderately present. [16] who reported that these secondary metabolites were present in varying concentrations and these variations can be explained by differences in agro-climatic conditions, age of plant, genotype, environmental conditions, post-harvest treatments, season of harvesting and maturation stage of the plants have strong influence on the phytochemical content of plants.

## 5. CONCLUSION

This study revealed the positive effects of solid state fermentation on phytochemicals composition of rice bran meal. The study revealed that, solid state fermentation of rice bran improved the flavonoids which serves as a flavour in the fish feed. The study also revealed that, solid state fermentation of rice bran reduces the alkaloids content of the rice bran meal which affect the nervous system thereby disrupting the electrochemical transmission in the fish.

## REFERENCES

1. Adekunle, A. A. (2012). Agricultural innovation in sub-saharan Africa: experiences from multiple stakeholder approaches. *Forum for Agricultural Research in Africa, Ghana*. Pp 21-29.
2. Akande K. E., Doma, U.D., Agu, H.O. and Adamu, H.M. (2010). Major Anti-nutrients Found in Plant Protein Sources: Their Effect on Nutrition. *Pakistan Journal of Nutrition*, 9 (8): 827-832.
3. Chakraborty, S. B., Peter, H. and Csaba, H. (2013) Application of phytochemicals as growth-promoters and endocrine modulators in fish culture. *Aquaculture*, 5: 1-19.
4. Edeoga, H.O., Okwu, D.E., Mbaebie, B.O. (2005). Phytochemical constituents of some Nigeria medicinal plants. *African Journal of Biotechnology*, 4(7):685-688.
5. Falaye, C. A., Chikwunde, A. G. and Micheal, B. J. (2014) Improving Nutritional Quality of Millet (*Pennisetum americanum*) by Solid State Fermentation and the Effects on the Growth Performance of African Catfish. *Journal of Natural Sciences Research*, 3 (6): 45-52.
6. Falaye, A.E. and Oloruntuyi, O.O. (1998) Nutritive Potential of Plantain Peel Meal and Replacement Value for Maize in Diets of African Catfish (*Clarias gariepinus*) Fingerlings *Journal of Tropical Agriculture (Trinidad)*, 75 (4): 488-492.
7. Fasakin, E.A. (2008). Fish as Food yesterday, today and forever *Inaugural Lecture*, series 48, The Federal University of Technology, Akure, pp 52.

8. Gao, Y., Lu, S., Wu, M., Yao, W., Jin, Z. and Wu, X. (2019). Effect of dietary protein levels on growth, feed utilization and expression of growth related genes of juvenile giant grouper (*epinephelus lanceolatus*). *Aquaculture*, 504, 369-374.
9. Habtamu, F. and Negussie, R. (2014). Anti-nutritional factors in plant foods: Potential health benefits and adverse effects. *International Journal of Nutrition and Food Sciences*, 3(4): 284-289.
10. Idowu1, T.A., Ishaku, A. H., Adedeji1, H. A., Olaniyi, I.J. and Sogbesan, O.A. (2016) Effect of Varying Levels of Fermented Sorghum (*Sorghum Bicolor*) On Growth and Haematology Profiles of *ClariasGariiepinus*(Burchell, 1822) Fingerlings. *Journal of Agriculture and Environment*, 12 (1): 89-96.
11. Khare, N. (2000). Phytic Acid Composition and its Effects. *Journal of Biochemistry*, 4(2): 321-336.
12. Mohammed, R. A., Jauro, I. A., Ishaku, A. H. and Usman, S. (2017) Growth Performance and Haematological Profiles of *Clariasgariiepinus*(Burchell, 1822) Fed Fermented Sorghum By-product Meal. *Journal of Agriculture and Environment*, 13(2): 137-147.
13. Obe, B.W. (2014). Growth Performance and Nutrient Utilization of Catfish Hybrid (*Heterobranchusbidorsalis*X *Clariasgariiepinus*) Fed Fermented Sorghum (*Sorghum bicolor*) Waste MealDiets. *International Journal of Applied Science and Technolog*, 4(3):156-160.
14. Olukunle, O. (2006). Nutritive Potential of Sweet Potato Peel Meal and Root Replacement Value for Maize in Diets of Africa Catfish (*Clariasgariiepinus*) Advanced Fry. *Journal of Food Technology*, 4(4): 289- 293.
15. Olurin, K.B., Olujo, E.A.A. and Olukoya, O.A. (2006) Growth of African catfish *Clariasgariiepinus*fingerlings, fed different levels of cassava. *World Journal of Zoology*, (1): 54-56.
16. Soetan K. O. and Oyewole, O. E. (2009).The need for adequate processing to reduce the anti-nutritional factors in plants used as human foods and animal feeds: A review. *African Journal of Food Science*, 3 (9): 223-232.
17. Sogbesan, O.A., K.N. Mohanta, P.K. Sahoo and P. Jayasankar (2012). *In: Application of solid state fermentation technology in aquaculture*, pp 76-83.
18. Sogbesan, A.O and Ugwumba A.A.A (2008) Nutritional Evaluation of Termite (*Macrotermes subhyalinus*) Meal as animal protein Supplements in the Diets of *Heterobranchuslongifilis*Fingerlings. *Turkish Journal of Fish and Aquatic Science*, 8:149-157.

19. Sotolu, A. O. (2008). Nutrient potential of water hyacinth as a feed supplement in sustainable aquaculture. *Obeche*, 26 (1): 45-51.
20. Suprayudi, M.A. (2010). Local Raw Materials: Challenges and Expectations of Future Aquaculture (Abstract). *Proceedings of the National Symposium on Biotechnology Aquaculture III* October 7, 2010. BDP, FPIK, IPB. 31p.
21. Thava, S. and James, M. (2001). Effects of Phytic Acid on Bioavailability in Fish Feed. *Int. Journal of Food Science*, 17(1): 419-431.
22. Umaru, H. A., Adamu, R. and Nadro, M.S. (2007). Effects of Anti-Nutritional Factors in Some Wild Edible Fruits of Northern Nigeria. *African journal of biotechnology*, 16 (1): 77-77.
23. Vaghasiya, Y., Dave, R. and Chanda, L. (2011). Phytochemical analysis of some medicinal plants from western region of India. *Resource Journal Medicinal Plants*, 5(5): 567-576.
24. Walter, H.L, Fanny, L., Charles, C., and Christian, R., (2002). Minerals and phytic acid interaction: is it a real problem for human nutrition. *International Journal of Food Science Technology*, (37) 727-739.
25. Yang, X., Wan, Z., Perry, H. Lu, Q., Wang, C., Hao, J., Li, F., Xie, J., Yu, T., Cui, T., Wang, M., Li and Ge, Q. H. (2012). Early millet use in northern China. *Proceeding of National. Academic Science USA* pp. 1-5.