

Genotypic × Environment Interaction and Stability Analysis for Bulb Yield and Yield Attributes in Onion (*Allium cepa* L.)

Formatted: Font color: Red

ABSTRACT

An experiment was conducted for estimating the Genotypic × Environment interaction and stability for bulb yield and yield attributes in onion. 36 elite genotypes were evaluated during winter season (*rabi*) 2022-23 at three different environments in 3 replications. Estimate of environment wise analysis of variance suggested the significant differences for all the characters in all environments. Significant mean squares due to environments were also observed for all the traits under study showed that environments selected for study were random and different in agro-climatic conditions. Genotypes x interactions were significant for all the characters. Joint consideration of mean performance and stability parameters revealed that parents RO-1 and RO-59 were above average stable for bulb yield and were considered suitable for general adaptation. The crosses RO-1 × RO-59, RO-1 × Pusa Shobha, RO-1 × Pusa Madhavi, Pusa Shobha × Pusa Madhavi and Pusa Madhavi × Pusa Red were considered suitable for better environmental condition for bulb yield, while, the crosses RO-1 × Bhima Kiran, RO-59 × Pusa Madhavi, RO-59 × Pusa Red, Bhima Kiran × Bhima Shakti, Bhima Kiran × Pusa Shobha, Bhima Kiran × Pusa Madhavi and Bhima Kiran × Kashi No. 1 were considered suitable for general adaptation.

Formatted: Font: Italic

Keywords: Stable genotype, ~~genotype~~ Genotype x ~~environment~~ Environment interaction, Onion, Stability

INTRODUCTION

Onion (*Allium cepa* L.) is one of the important bulb crop of the family Amaryllidaceae and grown widely all over the world and consumed in various forms. It has been in cultivation for more than 4000 years ago among vegetables. Onion is the third most important crop after potato and tomato. It is found in most market of the world throughout the year and can be grown under

Formatted: Font: Italic

wide range of Agro-climate condition. The maximum diversity of *Allium* species is found in a belt from Mediterranean basin to Iran and Afghanistan. Onion is cultivated mainly as annual for bulb production and biennial for seed production. The stability of performance of genotypes over environments is of great significance to the vegetable breeders in order to separate effect of environment. (Khar *et al.*, 2010) Many varieties of different crops do not show consistency in performance when grown over environments owing to the presence of genotype \times environment interaction. Such interactions change relative ranking of genotypes in different environments and also reduce magnitude of genetic differences among the genotypes. This poses a problem before the vegetable breeders in proper assessment of genotypes tested under different environmental conditions. The study of genotype \times environment interaction is important not only from the genetical and evolutionary point of view but also gains significance to tackle agricultural problem in general and with regards to breeding in particular (Tahir *et al.*, 2020). Stability study is not only important in finding out the stable genotypes for varied environmental conditions but also equally useful in screening the different genotypes for their comparative performance under a particular set of environments. Besides the high productivity of genotype, the stability of its performance is also necessary for stabilizing the production. Ceccarelli (1989) expressed that higher attention should be given to the assessment of yield stability. Eberhart and Russell (1966) discussed stability of genotypes in terms of three parameters – mean, regression or linear response and deviation from regression. They suggested that a stable variety is one with high mean, unit regression and least deviation from regression.

Formatted: Highlight

MATERIALS AND METHODS

The experiment was conducted at Horticulture Research Farm, Sri Karan Narendra College of Agriculture, Jobner Rajasthan to evaluate the stability of genotypes in different environmental conditions in terms of bulb yield and yield related characters. During *rabi* 2021-22 crossing block was constructed and direct F₁ crosses were produced following a half diallel mating design. The experimental material consisting of 8 parents and 28 F₁'s were planted during *rabi* 2022-23. All the genotypes were sown under three different environments namely; early, normal and late environment. The experiment was laid down in randomized block design

with three replications in all the environments. The plot for non-segregating generations (parents and F₁'s) represented two rows each. Rows were planted in 3m length spaced at 15cm apart and 10 cm interplant distance under all the environments. Non experimental rows were planted all around the experimental material to avoid any possible border effects. Sowing was done by hand drilling of the seeds in the soil. All other agricultural practices were done according to the recommendation and package of practices implemented at optimum period to establish good crop population in the field. Observations were recorded on five randomly selected competitive plants from each replication and environment for the bulb yield attributes and quality parameters namely, plant height, number of leaves, total chlorophyll content, number of days to 50% neckfall, average bulb weight, bulb volume, number of scales, neck thickness, equatorial diameter, polar diameter, polar : equatorial ratio, bulb yield, thrips incidence, purple blotch disease incidence, total soluble solids, allyl propyl disulphide. The data on each character for the genotypes were subjected to standard statistical analysis of variance for each environment separately [Panse and Sukhatme, 1985]. Later the data of each were subjected to pooled analysis of variance [Singh and Choudhary, 1985]. The stability analysis was done according to Eberhart and Russell (1966) model.

Results and Discussion

In the present investigation, the mean square due to environmental (E), genotype (G) × environmental (E), environmental (linear) and genotype × environmental (linear) were significant. The mean sum of squares due to pooled deviation (non linear) were significant indicating role of unpredictable causes affecting stability and prediction of these attributes would be difficult [Table-1]. [Khar *et al.* (2010) and Tahir *et al.* (2020) in onion, Dhaduktet *et al.* (2012), Singh *et al.* (2016) and Belay *et al.* (2019) in garlic, Tak *et al.* (2022) in cucumber also reported similar results.

For interpretation of results considering only bulb yield per plant, deviation from regression (unpredictable) was taken as a parameter of stability and regression coefficient as a parameter of responsiveness. The parents like RO-1 and RO-59 for average bulb weight and bulb yield, Pusa Madhavi for plant height and total chlorophyll content, Bhima Kiran, Pusa Shobha, Pusa Red for number of days to 50 % neckfall, Bhima Shakti for allyl propyl disulphide were

Comment [c1]: Use the same format of adding reference in the manuscript,

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

stable but their performance was exhibiting average stability and adaptability, hence, most stable. The parents like RO-1 for total soluble solids, Kashi No. 1 for allyl propyl disulphide, Pusa Madhavi for equatorial diameter exhibiting below average stability hence, were desirable for better environment. The Parents like RO-1 for neck thickness, bulb volume, number of scales, Bhima Kiran for number of leaves exhibiting above average stability hence, these crosses were suited for poor environmental condition [Table-2]. Among the F₁s RO-1 × Bhima Kiran, RO-59 × Pusa Red, Bhima Kiran × Bhima Shakti, Bhima Kiran × Pusa Shobha, Bhima Kiran × Pusa Madhavi and Bhima Kiran × Kashi No. 1 were most stable along with high mean value, hence were considered desirable. The crosses RO-1 × RO-59, RO-1 × Pusa Shobha, Pusa Shobha × Pusa Madhavi and Pusa Madhavi × Pusa Red had regression more than unity along with high mean value, exhibiting more responsiveness hence were considered desirable for better environmental conditions only. The crosses RO-1 × Kashi No. 1 had high mean value and regression coefficient more than unity, exhibiting less responsiveness, therefore were desirable for poor environmental condition [Table-2]. Khar *et al.* (2010) and Tahir *et al.* (2020) in onion, Dhaduktet *et al.* (2012), Singh *et al.* (2016) and Belay *et al.* (2019) in garlic, Somanath (2008), Soni (2009), Yadav and Ram (2010), Singh and Ram (2010), Khan and Sarolia (2019), Farag *et al.* (2019), Tak *et al.* (2022) in cucurbits also reported similar results.

Formatted: Highlight

Comment [c2]: Add recent reference in the support of the results.

Formatted: Highlight

Since *per se* performance and stability are two independent traits and are controlled by different sets of gene system (Yadav and Ram, 2010), It was expected that the stability of F₁s might be influenced by sets of genes for stability present in parents. The results indicated that sets of genes for stability were inherited from its parents but this relationship was not noticed in all the cases. On the basis of these results, it can be concluded that the inheritance of stability has differential ~~behaviour~~ behavior for genotypes. Therefore, further study is needed to learn more about inheritance of stability.

CONCLUSION

On the basis of phenotypic stability, parents *viz.* RO-1 and RO-59 were above average stable for bulb yield and were considered suitable for general adaptation. The crosses RO-1 X RO-59, RO-

Formatted: Font: Italic

1 X Pusa Shobha, RO-1 X Pusa Madhavi, Pusa Shobha X Pusa Madhavi and Pusa Madhavi X Pusa Red were considered suitable for better environmental condition for bulb yield, while, the crosses RO-1 X Bhima Kiran, RO-59 X Pusa Madhavi, RO-59 X Pusa Red, Bhima Kiran X Bhima Shakti, Bhima Kiran X Pusa Shobha, Bhima Kiran X Pusa Madhavi and Bhima Kiran X Kashi No. 1 were considered suitable for general adaptation and the cross RO-1 X Kashi No. 1 was considered suitable for poor environmental condition in the favour of bulb yield (q /ha).

Comment [c3]: Reframe conclusion part.

UNDER PEER REVIEW

Table - 1: Pooled analysis of variance showing mean squares of parents and F₁'s for yield and its contributing attributes

Particulars		Mean Squares													
Source of Variation	df	Plant height	Number of leaves	Chlorophyll content of leaves	Number of days to 50% neckfall	Average bulb weight (g)	Bulb volume (cc)	Number of scales	Neck thickness (cm)	Equatorial diameter (cm)	Polar diameter (cm)	Polar : Equatorial ratio	Bulb yield (q/ha)	Total soluble solids (%)	Allyl propyl disulphide (mg)
Genotypes (G)	35	104.58**	1.34**	0.011**	35.41**	77.47**	24.84**	1.05**	0.0153**	0.13**	0.12**	0.0021**	3313.48**	0.74**	353.23**
Environment (E)	2	1353.74**	26.36**	0.074**	813.29**	4510.98**	217.4**	6.53**	0.3003**	0.51**	1.08**	0.0129**	203060.96**	0.65**	2.23
G x E	70	22.42**	0.39**	0.01**	14.04**	11.86**	11.77**	0.45**	0.0048**	0.07**	0.05**	0.0024**	528.02**	0.36**	226.23**
E + (G x E)	72	59.4**	1.11**	0.011**	36.24**	136.83**	17.48**	0.62**	0.013**	0.08**	0.08**	0.0027**	6153.9**	0.37**	255.35**
E (linear)	1	2707.5**	52.72**	0.149**	1626.57**	9021.91**	434.8**	13.06**	0.6005**	1.02**	2.17**	0.0259**	406122.09**	1.31**	209.7**
G x E (linear)	35	17.36**	0.34**	0.014**	21.2**	9.15**	9.62**	0.53**	0.0044**	0.08**	0.05**	0.0028**	412.58**	0.28**	468.48**
Pooled deviation	36	26.72**	0.43**	0.005**	6.69**	14.16**	13.53**	0.36**	0.0051**	0.06**	0.04**	0.002**	625.59**	0.43**	17.29**
Pooled Error	216	4.653	0.097	0.001	2.653	4.033	2.507	0.04	0.001	0.017	0.017	0.0005	146.51	0.07	3.91

*, **Significant at 5 percent and 1 percent levels, respectively

Comment [c4]: Add unit of it.

Table-2: Estimates of stability parameters for Bulb yield (q/ha)

Parents/ cross	Bulb yield (q/ha)		
	\bar{X}	bi	S^2_{di}
Parents			
RO-1 (P1)	278.51	1.01	-139.4357
RO-59 (P2)	267.47	0.99	-36.3512
BhimaKiran (P3)	230.19	1.01	-145.5624
Bhima Shakti (P4)	232.64	1.01	5.876
PusaShobha (P5)	230.93	1.01	-145.8906
PusaMadhavi (P6)	223.30	1	-102.653
Kashi No. 1 (P7)	209.51	0.92	-127.4577
Pusa Red (P8)	206.92	0.95	0.2094
Crosses			
P1xP2	344.32	1.37**	-132.8259
P1xP3	282.51	1	-118.9602
P1xP4	232.86	0.6	665.8768*
P1xP5	320.75	1.45*	215.3321
P1xP6	325.79	1.44*	180.6978
P1xP7	286.96	0.71	143.4808
P1xP8	289.33	1.07	3270.9129**
P2xP3	237.16	0.98	-144.5181
P2xP4	217.44	0.54	515.4691*
P2xP5	239.53	0.99	-127.2998
P2xP6	253.76	0.98	-144.6507
P2xP7	242.94	1.02*	-145.2297
P2xP8	259.76	0.96	-103.0539
P3xP4	258.13	0.99	-77.0878
P3xP5	268.95	0.97	-131.0191
P3xP6	276.44	1.07**	-140.23402
P3xP7	273.32	0.98**	-145.9341
P3xP8	245.09	0.96	-26.1832
P4xP5	242.49	0.98	495.8368*
P4xP6	250.64	0.93	-93.7499
P4xP7	227.60	0.76	9322.33**
P4xP8	203.95	0.99	2858.816**
P5xP6	234.93	1	-6.5113
P5xP7	215.74	0.93	1160.4988**
P5xP8	271.47	1.37	253.6648
P6xP7	222.26	1.03	314.983
P6xP8	266.50	1.19	218.8018
P7xP8	251.09	0.83**	-141.3573
General mean	253.37	1.00	
S.E. (bi)		0.15	

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Reference

Belay, F., Tekle G. and Chernet S. 2019. Evaluation of genotype × environment interaction effect on performance of garlic (*Allium sativum* L.) genotypes in Tigray region, Northern Ethiopia using AMMI and GGE biplot analysis. *African Journal of Agricultural Research*, **16**(5): 691-701.

Formatted: Font: Bold

Dhadukt L. K., Patel A. G., Mehta D. R., Madariya R. B. 2012. Genotypes x environment interactions and stability analysis for bulb yield and its attributes in garlic (*Allium sativum* L.). *Progressive Agriculture*, **11**: 170- 173

Formatted: Font: Bold

Eberhart, S.A. and Russel, W.A. 1966. Stability parameters for comparing varieties. *Crop Science*, **6**: 34-40.

Farag, M. I., Behera, T. K., Munshi, A. D., Bharadwaj, C., Jat, G. S., Khanna M. and Chinnusamy, V. 2019. Physiological analysis of drought tolerance of cucumber (*Cucumis sativus*) genotypes. *Indian Journal of Agricultural Sciences*, **89**(9): 1445–50.

Khan R. and Sarolia D.K. 2019. Stability analysis in bitter gourd (*Momordica charantia* L.). *Agricultural Research Journal*, **56** (3): 392-400.

Formatted: Font: Bold

Khar, A., Devi, A., Mahajan V. and Lawande K.E. 2010. Stability analysis of some elite onion lines in late *kharif* season. *Indian J. Hort.*, **64**(4): 415-419.

Panse V.C., Sukhatme P.V. 1985. Statistical methods for agricultural workers. *Indian Council of Agricultural Research*, New Delhi. :87-89.

Comment [c5]: Add full details of reference

Singh R.K., Choudhary B.D. 1985. Biometrical methods in quantitative genetics analysis. *Kalyani Publishers*, New Delhi.

Singh, A. and Ram, H.H. 2010. Stability analysis for flowering and maturity traits in cucumber. *Veg. Sci.*, **39**: 31-34.

Singh, R. K., Dubey, B. K. & Gupta, R. P. 2016. Genotype x environment interaction and stability analysis for yield and its attributes in garlic (*Allium sativum* L.). *Journal of Spices and Aromatic Crops*, **25** (2) : 175–181.

Somanath, C.A. 2008. Stability analysis of some promising bitter gourd (*Momordica charantia* L.) hybrids. A thesis submitted to department of genetics and plant breeding, *University of Agriculture Sciences, Dharwad*.

Soni, K.K. 2009. Genetics and phenotypic stability of fruit yield and its components in bottle gourd [*Lagenaria siceraria* (Mol.) Standl]. Ph.D. Thesis submitted to *S.K.R.A.U, Bikaner, Rajasthan*.

Tahir, M., Aleem, S., Munawar, M., Parveen, N., Amin, E., Aslam, R., Saeed A. and Najeebullah, M. 2020. GGE biplot an effective tool to study genotype and genotype × environment interaction; a case study in onion (*Allium cepa* L). *Pak. J. Agri. Sci.*, Vol. **57**(6): 1565-1571.

Tak, S., Kaushik, R.A., Ameta, K.D., Dubey R.B. and Rathore R.S. 2022. Effect of GXE interaction on yield analysis of selected cucumber and Snapmelon breeding lines in Rajasthan. *The Pharma Innovation Journal*, **11**(2): 1978-1982.

Formatted: Font: Bold

Yadav, R.K. and Ram, H.H. 2010. Genotype-environment interaction and stability analysis for yield and yield attributing characters in musk melon. *Indian Journal of Horticulture*, **67** (Special Issue): 179-184.

Formatted: Font: Bold

UNDER PEPH