

1  
2  
3  
4  
5  
6  
7

# ASSESSMENT OF DIFFERENT GERMPLOSM OF *Aglaonema* FOR POT AND REMOVAL OF PARTICULATE MATTER IN AIR

## ABSTRACT

The aim of the experiment is to screen twelve germplasm of *Aglaonema* for their suitability as pot plants and removal of particulate matter in air. The experiment was designed using Completely Randomized Design (CRD) at the greenhouse complex of the Department of Floriculture and Landscape Architecture, ASPEE College of Horticulture, Navsari (Gujarat), during 2023-24. Morphological parameters were measured using various methods, including a measuring tape for plant height, a digital leaf area meter for leaf area, and manual counting for leaves and branches. Leaf color was evaluated on visual grading system and the RHS Colour Chart (2015), to assess visual appeal and characteristics and percent removal of particulate matter (PM) by the plant was measured using digital air quality index meter. *Aglaonema* 'Emerald Bay' exhibited maximum plant height (73.18 cm), leaf area (311.66 cm<sup>2</sup>), leaf length (39.28 cm), leaf width (15.26 cm), internodal length (3.43 cm) and visual leaf colour grade (4.88) and dark green leaf colour (137A) as per RHS colour chart followed by *Aglaonema commutatum* with the plant height (69.58 cm), leaf area (298.09 cm<sup>2</sup>), leaf length (34.28 cm), leaf width (11.10 cm), internodal length (3.34 cm) and visual leaf colour grade (4.80) and dark green leaf colour (139 A). Maximum number of leaves were recorded in *Aglaonema* 'Cutlass' (43.90), while maximum number of branches per plant were observed in *Aglaonema brevispathum* (13.56). *Aglaonema* 'Emerald Bay' showed the maximum removal rate of PM<sub>1</sub> per 24 hours (92.16%) followed by *Aglaonema commutatum* (85.97%). Among all the germplasms, *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* have been highly suitable as pot plant owing to its good plant height, leaf area and their efficiency in removal of PM<sub>1</sub>.

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

*Keywords:* *Aglaonema*, pot plant, morphological, visual leaf colour, particulate matter.

## 1. INTRODUCTION

Indoor gardening has been gaining high impetus with enhanced sense of emotional comfort, air quality and stress-free happy environment during the post COVID era [1]. Further, pot plant trading is growing with rapid urbanization and hasty life style [2], [3] and [4].

*Aglaonema*, commonly known as Chinese evergreens, is a genus of 21 species within the Araceae family, native to Southeast Asia, Northeast India, and Southern China, with a distribution extending to Malaysia, New Guinea, and the Philippines [5], [6]. This genus has been extensively studied for its adaptability to indoor environments and its potential uses [7], [8]. It is known for its striking foliage and adaptability, is a widely cultivated genus of ornamental plants, particularly valued as pot plants, it encompasses a diverse range of species and cultivars, each exhibiting unique morphological traits that influence their suitability for pot cultivation. *Aglaonema* is ideal for indoor environments due to its air-purifying capabilities, adaptability to low light, aesthetic diversity, low maintenance, making it

24 both functional and visually appealing[9],[10]. It is essential to evaluate and screen various  
25 germplasms for desirable characteristics as pot plant and its ability to curb indoor pollution.

26 *Aglaonema*'s contribution to indoor air quality is supported by its ability in removal of  
27 particulate matter and other pollutants, that makes it a valuable addition to indoor  
28 environments, promoting healthier and cleaner air [11].However, there are different species  
29 and varieties of *Aglaonema* that needs to be studied for air quality purification.

30 Hence this study focused on the screening of twelve *Aglaonema* germplasms,  
31 namely *Aglaonema siam aurora* (Lipstick), *Aglaonema* 'Emerald Bay', *Aglaonema*  
32 *commutatum*, *Aglaonema alumina*, *Aglaonema* 'Key Lime', *Aglaonema* cutlass, *Aglaonema*  
33 'White Rain', *Aglaonema brevispathum*, *Aglaonema* lime narrow, *Aglaonema commutatum*  
34 (variegated), *Aglaonema* white Chinese evergreen (Golden Bay), and *Aglaonema* Red  
35 Valentine. The evaluation is based on key morphological parameters and its ability to curb  
36 indoor pollution.

37

## 38 **2. MATERIAL AND METHODS**

39 The experiment was conducted during 2023-24 at the greenhouse complex of Department of  
40 Floriculture and Landscape Architecture, ASPEE College of Horticulture, Navsari (Gujarat),  
41 using a Completely Randomized Design (CRD) to screen twelve *Aglaonema* germplasms for  
42 their suitability as pot plants and removal of particulate matter in air. The plants were  
43 maintained under uniform growing conditions to ensure consistency in environmental factors  
44 such as light, water, and nutrients. Morphological parameters were recorded both at the start  
45 of the experiment and after two months. plant height was measured from the soil level to the  
46 growing tip using a measuring tape. Leaf area was determined using a digital leaf area  
47 meter, while leaf length and width were measured using a meter scale. The number of  
48 leaves and branches per plant were counted manually. The visual colour grade of  
49 *Aglaonema* plants was assessed using a standardized grading system [12]. This system  
50 evaluates leaf colour grade based on greenness with freshness and luster, assigning scores  
51 1= poor colour, 3= good, light green and 5= excellent dark green & silver contrast, and visual  
52 leaf color was evaluated using the RHS Colour Chart (2015). Internodal length was  
53 measured between nodes using a meter scale. Particulate matter (PM<sub>1</sub>) level in air was  
54 recorded by using enclosed chamber using digital air quality index meter [13] before and  
55 after 24 hours and the PM<sub>1</sub> removal rate by the plant was calculated on per cent basis. The  
56 data on all the parameters was collected twice at an interval of two months were pooled and  
57 statistically analyzed as per the method given by Panse and Sukhatme [14] and the results  
58 were evaluated at 5% level of significance.

## 59 **3. RESULTS AND DISCUSSION**

60 Observations on morphological parameters viz., plant height (cm), leaf area (cm<sup>2</sup>), leaf  
61 length (cm), leaf width (cm) and internodal length (cm) of twelve germplasm of *Aglaonema* are  
62 represented in the table 1. Maximum plant height exhibited by *Aglaonema* 'Emerald  
63 Bay' (73.18 cm), closely followed by *Aglaonema commutatum* (69.58 cm). In contrast,  
64 *Aglaonema siam aurora* (lipstick) displayed the minimum plant height (30.68 cm). Maximum  
65 leaf area exhibited by *Aglaonema* 'Emerald Bay' (311.66 cm<sup>2</sup>), closely followed by  
66 *Aglaonema commutatum* (298.09 cm<sup>2</sup>). In contrast, *Aglaonema siam aurora* (lipstick)  
67 displayed the smallest leaf area (111.55 cm<sup>2</sup>). *Aglaonema* 'Emerald Bay' showed the  
68 maximum leaf length (39.28 cm) followed by *Aglaonema commutatum* (34.28 cm) while it was  
69 minimum in *Aglaonema siam aurora* (lipstick) (15.25 cm). *Aglaonema* 'Emerald Bay' exhibited

70 maximum leaf width (15.26 cm), closely followed by *Aglaonema* 'white Rain' (13.30 cm),  
 71 whereas *Aglaonema* 'cutlass' displayed the narrowest leaves (4.83 cm). Maximum internodal  
 72 length was recorded in *Aglaonema* 'Emerald Bay' (3.43 cm), closely followed by *Aglaonema*  
 73 *commutatum* (3.34 cm). In contrast, *Aglaonema* 'lime narrow' exhibited the shortest  
 74 internodal length (1.52 cm).

75 Observations on number of leaves, number of branches per plant, visual leaf color grade of  
 76 twelve germplasm of *Aglaonema* are represented in the table 2. Maximum leaf count was  
 77 observed in *Aglaonema* 'Cutlass' (43.90), followed by *Aglaonema* 'Alumina' (25.08), while  
 78 *Aglaonema* 'White Rain' had the minimum (11.16). *Aglaonema brevispathum* exhibited  
 79 maximum number of branches per plant, with an average of 13.56, closely followed by  
 80 *Aglaonema* 'Cutlass' (12.24 branches per plant). In contrast, *Aglaonema siamauror*'  
 81 (Lipstick) displayed the lowest branching frequency, with an average of 4.20 branches per  
 82 plant. *Aglaonema* 'Emerald Bay' exhibited the highest visual leaf color grade (4.88), closely  
 83 followed by *Aglaonema* 'Alumina' (4.80) on the basis of 5 point visual grade for freshness,  
 84 greenness and luster. Variation with regard to colour as per RHS chart among different  
 85 germplasms was observed, wherein *Aglaonema* 'Emerald Bay', *Aglaonema*  
 86 *commutatum* and *Aglaonema alumina* exhibited dark green colour, *Aglaonema siam aurora*  
 87 (Lipstick) exhibited dark purple red, *Aglaonema* 'Key Lime' and *Aglaonema* lime narrow  
 88 exhibited medium yellow green, *Aglaonema* cutlass exhibited medium green, *Aglaonema*  
 89 'White Rain', *Aglaonema commutatum* (variegated) and *Aglaonema* white Chinese  
 90 evergreen (Golden Bay) exhibited light green and *Aglaonema* Red Valentine exhibited light red  
 91 pink colour (Fig.1).

92 Observations on removal rate of Particulate Matter per 24 h (PM1) (%) is represented in the  
 93 table 3. *Aglaonema* 'Emerald Bay' showed the maximum removal rate of PM1 (%) in 24  
 94 hours (92.16%) followed by *Aglaonema commutatum* (85.97%) while it was minimum  
 95 in *Aglaonema* Red Valentine (63.06%). The results suggest that the morphological  
 96 characteristics of plants play a crucial role in reducing PM concentration. Leaf traits,  
 97 including shape, arrangement, and the leaf area and hair structure on the surface, are  
 98 particularly important for PM accumulation. Plants remove PM through stomatal uptake and  
 99 deposition on the leaf surface. PM particles can penetrate the stomata and accumulate  
 100 beneath the epidermis or in the palisade layer of leaves [15],[16],[17],[18],[19] and [20].  
 101 Higher leaf area in *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* contributed to  
 102 higher PM removal rate.

103  
 104  
 105  
 106  
 107  
 108

**Table 1: Morphological characters of twelve germplasm of *Aglaonema***

<b>Morphological characters</b>					
<b>Treatments Pooled data of two months</b>					
	<b>Plant Height(cm)</b>	<b>Leaf area (cm<sup>2</sup>)</b>	<b>Leaf length (cm)</b>	<b>Leaf width (cm)</b>	<b>Internodal length (cm)</b>
<b>G<sub>1</sub></b>	30.68	111.55	15.25	6.28	1.92
<b>G<sub>2</sub></b>	73.18	311.66	39.28	15.26	3.43

<b>G<sub>3</sub></b>	69.58	298.09	34.28	11.10	3.34
<b>G<sub>4</sub></b>	40.24	212.43	21.71	10.53	2.91
<b>G<sub>5</sub></b>	54.74	212.26	23.22	11.69	3.08
<b>G<sub>6</sub></b>	45.07	113.70	27.82	4.83	2.23
<b>G<sub>7</sub></b>	62.75	236.36	33.94	13.30	2.06
<b>G<sub>8</sub></b>	56.48	271.19	29.14	12.57	3.00
<b>G<sub>9</sub></b>	56.15	269.81	25.40	7.57	1.52
<b>G<sub>10</sub></b>	66.49	154.72	26.09	8.30	2.25
<b>G<sub>11</sub></b>	35.42	252.93	25.58	12.21	2.80
<b>G<sub>12</sub></b>	37.05	119.65	20.61	6.35	1.81
<b>S.Em ±</b>	<b>0.39</b>	<b>1.76</b>	<b>0.45</b>	<b>0.15</b>	<b>0.04</b>
<b>CD at 5%</b>	<b>1.13</b>	<b>5.03</b>	<b>1.30</b>	<b>0.38</b>	<b>0.10</b>

109

110

**Table 2: Morphological characters of twelve germplasm of *Aglaonema***

<b>Morphological characters</b>					
<b>Treatments</b>	<b>Pooled data of two months</b>				
	<b>Number of leaves</b>	<b>Number of branches per plant</b>	<b>visual leaf colour grade</b>		
			<b>(5-point scale)</b>	<b>RHS Colour group</b>	<b>colour code</b>
<b>G<sub>1</sub></b>	13.43	4.20	4.21	Dark purple red	46 A
<b>G<sub>2</sub></b>	14.16	4.89	4.88	Dark green	137 A
<b>G<sub>3</sub></b>	11.18	8.36	4.80	Dark green	139 A
<b>G<sub>4</sub></b>	25.08	9.35	4.79	Dark green	131B
<b>G<sub>5</sub></b>	19.78	6.58	4.34	Medium yellow green	151 D
<b>G<sub>6</sub></b>	43.90	12.24	4.61	Medium green	144 A
<b>G<sub>7</sub></b>	11.16	10.83	4.30	Light green	141D
<b>G<sub>8</sub></b>	13.63	13.56	4.71	Medium grey green	191 C
<b>G<sub>9</sub></b>	12.86	11.56	4.71	Medium yellow green	154 D
<b>G<sub>10</sub></b>	15.85	9.11	4.68	Light green	144 B
<b>G<sub>11</sub></b>	11.79	5.11	4.69	Light green	145 D
<b>G<sub>12</sub></b>	12.93	5.00	4.06	Light red pink	49 D
<b>S.Em ±</b>	<b>0.34</b>	<b>0.13</b>	<b>0.03</b>		
<b>CD at 5%</b>	<b>0.98</b>	<b>0.36</b>	<b>0.10</b>		


111 **Table .1 and 2. G<sub>1</sub>:** *Aglaonema siam aurora* (Lipstick), **G<sub>2</sub>:** *Aglaonema* 'Emerald Bay', **G<sub>3</sub>:** *Aglaonema*  
112 *commutatum*, **G<sub>4</sub>:** *Aglaonema alumina*, **G<sub>5</sub>:** *Aglaonema* 'Key Lime', **G<sub>6</sub>:** *Aglaonema cutlass*, **G<sub>7</sub>:**  
113 *Aglaonema* 'White Rain', **G<sub>8</sub>:** *Aglaonema brevispathum*, **G<sub>9</sub>:** *Aglaonema lime narrow*, **G<sub>10</sub>:** *Aglaonema*  
114 *commutatum* (variegated), **G<sub>11</sub>:** *Aglaonema white Chinese evergreen* (Golden Bay), **G<sub>12</sub>:** *Aglaonema*  
115 *Red Valentine*

116 **Table 3:Effect of different germplasms of *Aglaonema* on Removal rate of PM1 (%) in**  
117 **24 h**

<b>Removal rate of PM<sub>1</sub>(%)/ 24 h</b>	
<b>Treatments</b>	<b>Mean (T)</b>
<b>G<sub>1</sub>:</b> <i>Aglaonema siam aurora</i> (Lipstick)	<b>64.96</b>

<b>G<sub>2</sub>: Aglaonema 'Emerald Bay'</b>	<b>92.16</b>
<b>G<sub>3</sub>: Aglaonema commutatum</b>	<b>85.97</b>
<b>G<sub>4</sub>: Aglaonema alumina</b>	<b>80.05</b>
<b>G<sub>5</sub>: Aglaonema 'Key Lime'</b>	<b>81.09</b>
<b>G<sub>6</sub>: Aglaonema cutlass</b>	<b>70.03</b>
<b>G<sub>7</sub>: Aglaonema 'White Rain'</b>	<b>85.11</b>
<b>G<sub>8</sub>: Aglaonema brevispathum</b>	<b>81.13</b>
<b>G<sub>9</sub>: Aglaonema lime narrow</b>	<b>81.15</b>
<b>G<sub>10</sub>: Aglaonema commutatum (variegated)</b>	<b>72.15</b>
<b>G<sub>11</sub>: Aglaonema white Chinese evergreen (Golden Bay)</b>	<b>72.15</b>
<b>G<sub>12</sub>: Aglaonema Red Valentine</b>	<b>63.06</b>
<b>S.Em ±</b>	<b>0.12</b>
<b>CD at 5%</b>	<b>0.36</b>

118



119

120

121

**Fig. 1. Effect of different germplasms of *Aglaonema* on percent removal rate of PM<sub>1</sub>(%)/24h**

122

123

124

125

126

127

#### **4. CONCLUSION**

128

129

130

131

132

133

134

135

Based on the findings of the present investigation, *Aglaonema* 'Emerald Bay' and

136

137

*Aglaonema commutatum* exhibited superior morphological characteristics with maximum

values for plant height, leaf area, leaf length, leaf width, visual leaf colour grade with dark

green and lustrous leaf colour and along with efficient removal rate of particulate matter and

thus, making them the more suitable germplasms for pot plants. Overall, *Aglaonema*

'Emerald Bay' and *Aglaonema commutatum* are promising germplasms for pot plant

production and indoor air purification.

#### **Disclaimer (Artificial intelligence)**

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language

138 Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the  
139 writing or editing of this manuscript.

#### 140 **ACKNOWLEDGEMENTS**

141 We acknowledge ASPEE college of Horticulture (NAU, Navsari, Gujrat) for providing the  
142 best platform and necessary infrastructure and support in conduction of this research  
143 endeavor.

#### 144 **COMPETING INTERESTS**

145 Authorshavedeclaredthatnocompetinginterestsexits.

#### 146 **AUTHORS' CONTRIBUTIONS**

147 Thisworkwascarriedoutincollaborationamongallauthors.

148 Allauthorsreadandapprovedthefinalmanuscript.

#### 149 **REFERENCES**

- 150 [1] Singh A. Indoor Gardening for Clean Air. *Nursery Today*. 2020; July-October: 39-42.
- 151 [2] Singh A, Ahlawat TR, Chavan S and Patel AI. Wonder World of Adeniums. *Floriculture*  
152 *Today*.2016;20(9): 44-49.
- 153 [3]GuddadM, Singh A, Shah HP, Chaudhari P and Ahlawat TR . Effect of Foliar Application  
154 of Chemicals on Plant Architecture in Potted *Ixora chinensis* var. 'Mini Double'.*Curr J Appl*  
155 *Sci Tech*.2022;41(32): 9–15.
- 156 [4] Sindhuja M, Singh A, Bhandari AJ,ShahHP,Patel Aland Parekh VB.  
157 Morphologicalcharacterization of different genotypes of adenium.*Indian J Hort*. 2022;79(3):  
158 296-304.
- 159 [5]Nicolson DH. A revision of the genus *Aglaonema* (araceae). *Smithsonian contributions to*  
160 *botany*. 1969;1-66.
- 161 [6]Govaerts R and Frodin DG. World checklist and bibliography of Araceae (and  
162 Acoraceae).*Royal Botanic Gardens, Kew*.2002;1–560.
- 163 [7]HennyRJ,ChenJ,MellichTA andBrennan MS. 'MoonlightBay'*Aglaonema*.  
164 *HortScience*.2008(5);43:1598–1599.
- 165 [8] LiDM, Zhu GF, Yu Band Huang D.Comparative chloroplast genomes and  
166 phylogenetic relationshipsof *Aglaonema modestum* and five variegated cultivars of  
167 *Aglaonema*.*PLoS One*.2022;17(9): e0274067.
- 168 [9] Chen J, Li Q and Dai X. Effect of different growing media on the growth of  
169 *Aglaonema*.*Acta Hort*.2002; 572:265–270.
- 170 [10] Wolverson BC and Wolverson JD. Plants and soil microorganisms: removal of  
171 formaldehyde, xylene, and ammonia from the indoor environment. *J*  
172 *MissiAcadSci*.1993;38(2):11–15.
- 173 [11]KimKJ, KilMJ, Song JS, Yoo EH, Son KC and Kays SJ. Efficiency of volatile  
174 formaldehyde removal by indoor plants: Contribution of aerial plant parts versus the root  
175 zone. *HortScience*.2010; 45(10):1489–1495.
- 176 [12] HennyRJ, Poole RT and Conover CA. Visible leaf color grade.  
177 *HortScience*.1983;18(3):374.
- 178 [13] YooMH, Kwon YJ, Son KC,Kays SJ. Efficacy of indoor plants for the removal of single  
179 and mixed volatile organic pollutants and physiological effects of the volatiles on the  
180 plants.*JASHS*.2006; 131(4):452–458.
- 181 [14] Panse, V.G. and Sukhatme, P.V. *Statistical Methods for Agricultural Workers*. 4th ed.  
182 New Delhi: Indian Council of Agricultural Research; 1985.
- 183 [15] Chen H, Wang Z, Li X. Investigation of photo regulation in *Hedera helix* for improving  
184 indoor air quality.*JEnviron Manage*.2017;203: 981–988.

- 185 [16] ChenQ,Huang S, Shen L, Chen X. Mitigating indoor air pollutants using plant-based  
186 methods: A phytoremediation approach.Air QualAtmos Health.2017;10(4):473–485.  
187 [17] ChenW, Liu Y, Tong D.Photoregulation of potted Hedera helix to remove volatile  
188 formaldehyde for improving indoor air quality.Build Environ.2017;177: 245–253.  
189 [18] JangJ, Park S, LeeJ, KimJ. Effect of different houseplants on indoor air quality  
190 improvement. EnvironRes.2021; 194:110627.  
191 [19] RyuY, Jeon H, Lee J, Kim S. Evaluation of indoor air quality improvement by different  
192 types of houseplants.Indoor Built Environ.2013; 22(2):284–295.  
193 [20] Han J.A study on the effectiveness of various indoor plants in improving indoor air  
194 quality. J Environ Manage.2019;240: 285–293.  
195

## 196 **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

197 **Particulate matter 1 (PM1):** A type of air pollution that refers to particles that are less than 1  
198 micron in diameter.