

## “Heterosis and Inbreeding Depression for Yield and Yield Attributing Traits of

Maize(*Zea Mays*L.)”

Comment [I1]: Maize (Zea mays L.)

### Abstract

In present investigation45 single cross hybrids were evaluated along with their parental lines and checks heterosis and inbreeding depression. The standard heterosis, the hybrid combinations DI- 123 x DI- 111(91.11), DI- 121 x DI- 117 (80.82), DI- 121 x DI- 111(80.33), DI- 121 x D- 56 (78.33), DI- 123 x DI- 117 (74.10), DI- 122 x D- 52(70.23), DI- 121 x D- 52 (67.04) and DI- 117 x D- 56 (62.24) were identified as best for grainyield and some of the yield contributing traits. Maximum inbreeding depression was obtained from DI- 111 x DI- 118 (61.50) for grain yield per plant (g).

Comment [I2]: Investigation 45

Comment [I3]: Space between grain and yield

Comment [I4]: space after full stop

**Keywords:** Maize, Yield, Heterosis, Inbreeding depression

### Introduction

Maize (*Zea mays* L.) is one of the most versatilecrop having wider adaptability under varied agro-climatic conditions. Globally, maize is known as queen of cereals because it has the highest genetic yield potential among the cereals. Maize is a principal food crop for human consumption as well as animal feed around the world and as a model plant for understanding evolution, domestication, and heterosis (Romay *et al.*, 2013) it ranks third after wheat and rice (Gerpacio and Pingali, 2007).The kernel protein of maize is made up of five different fractions, viz., albumin 7%, globulin 5%, non-protein nitrogen 6%, prolamine 52% and glutelin 25% and left over 5% residual nitrogen.

Comment [I5]: versatile crop

In India its production has been recorded about 33.60mt from 10.00mha area with average productivity of 3.3 tons per ha. (Anonymous, 2022).

For improving of yield and quality parameters insight knowledge of genetic mechanism involved in the inheritance of various quantitative and quality traits is essential for effective manipulation. Hence, present investigation has been taken up to gather the information on the heterosisand selection of parents for productivity in maize. Diallel mating heterosisand design provides an opportunity to mate the given set of parents in all possible combinations (Griffing, 1956) and thus helps in the selection of desirable parents for utilization in the hybridization programme, as well as in the choice of appropriate breeding procedure for the genetic improvement of traits in maize. Grafius (1959) and Kebbleand Pellew (1910) concluded that the genetic basis of heterosis for a complex trait like yield could be explained by the multiplicative interaction on the phenotype level for component trait.

Comment [I6]: heterosis and

Comment [I7]: heterosis and

Comment [I8]: space

Comment [I9]: reference are missing

The information's on heritability and genetic advances are also useful for plant breeder. Genetic advance means improvement in the genotypic value of new population over base population. The systematic exploitation of heterosis for yield per plot and quality can be achieved through heterotic grouping only (**Melchinger and Gumber 1998**).

### **Materials and Methods**

The present investigation was carried out during *Rabi* 2022- 23 at Student's instructional farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur(U.P.).

The experimental material for present investigation comprised of 10 parental lines *viz.*, DI- 122, DI-121, DI-124, DI-123, DI-109, DI-111, DI-118, DI-117, D-52 and D-56. These parental lines were crossed to develop 45 F<sub>1</sub>s and F<sub>2</sub>s using half diallel mating design. A total of 100 treatments (10 parents + 45 F<sub>1</sub>s + 45 F<sub>2</sub>s) were evaluated for the study of genetical analysis for fifteen quantitative characters in maize.

The individual plant data for the following characters were recorded in each treatment of all replications on 10 randomly tagged plants in each parent, F<sub>1</sub>s and F<sub>2</sub>s for traits *viz.*, Days to 50% tasseling, Days to 50% silking, Plant height (cm), Days to 75 % dry husk, Number of cobs per plants, Cob length (cm), Cob diameter (cm), Cob weight (g), Number of grains per cobs, 100 Kernel weight (g), Shelling percentage, Protein content (%), Lysine content (%), Tryptophan content (%) and Grain yield per plant (g).

Heterosis was estimated as per cent increase or decrease in the mean values of F<sub>1</sub> hybrid over the standard commercial check variety value by formula heterosis (%) = [(F<sub>1</sub> - CV)/CV] x 100. where, F<sub>1</sub>s and CV represented the mean performance of hybrid and standard check variety. The significance test for heterosis was done by using standard error of the value of check variety.

The coefficient of inbreeding depression was calculated by the following formula:

$$\text{Inbreeding depression value (\%)} = \frac{\bar{F}_1 - \bar{F}_2}{\bar{F}_1} \times 100$$

Where,

$\bar{F}_1$  = Mean of F<sub>1</sub> generation

$\bar{F}_2$  = Mean of F<sub>2</sub> generation

The combining ability analysis was worked out by the procedure suggested by **Griffings (1956)** Method 2. Model I.

### **Result and Discussion**

According to **Griffing and Lindstrom (1954)** the expression of heterosis was due to accumulation of desirable genes in a hybrid plant through the crossing of parents in their genetic makeup and it had very often been related to the magnitude of genetic diversity. **Robinson (1960)** and **Moll et al., (1964)** were of the view that heterosis was mostly dependent on genetic diversity present in the parental stock.

Heterosis unless it is economical, is of no use. In the present investigation heterosis in F<sub>1</sub>'s over commercial variety (D- 52) and inbreeding depression were estimated and presented in Table 1.

It is evident from the result DI- 123 x D- 52 (-12.61), DI- 109 x DI- 118 (-12.18), DI- 109 x D- 56 (-10.88), DI- 111 x D- 56 (-10.88) and DI- 118 x DI- 117 (-10.88) the crosses combinations showed significant negative heterosis over economic parent for days to 50% tasseling in desirable direction out of which DI- 123 x D- 52 (-12.61) showed maximum while minimum in DI- 122 x D- 56 (-1.75). Significant negative heterosis over economic parent for days to 50% silking was ranged from -8.87 (DI- 121 x DI- 111) to -1.54 (DI- 122 x D- 56), and maximum from DI- 121 x DI- 111 (-8.87) followed by DI- 121 x DI- 118 (-6.95), DI- 122 x DI- 109 (-6.95) and DI- 122 x DI- 109 (-6.56). Cross combinations viz., DI- 123 x DI- 109 (-9.49), DI- 123 x DI- 117 (-8.54), DI- 123 x D- 52 (-8.22) and DI- 124 x D- 52 (-8.22) revealed significant negative heterosis over economic parent for days to 75% dry husk. These hybrids may be commercialized as early maturing maize hybrids after confirmation of their suitability for commercial cultivation.

Significant positive heterosis over economic parent was revealed by two cross combinations for plant height (cm), four for number of cobs per plant and cob diameter (cm), five for cob weight (g), six for cob length (cm), number of grains per cob and 100- kernel weight (g), seven for shelling (%), tryptophan content (%) and protein content (%), eight for lysine content (%). Cross combinations viz., DI- 123 x DI- 111 (91.11), DI- 121 x DI- 117 (80.82), DI- 121 x DI- 111 (80.33), DI- 121 x D- 56 (78.33), DI- 123 x DI- 117 (74.10), DI- 122 x D- 52 (70.23), DI- 121 x D- 52 (67.04) and DI- 117 x D- 56 (62.24) revealed significant positive heterosis over economic parent for grain yield per plant (g). Similar results were obtained by **Singh et al., (2010)**, **Kiani et al., (2015)**, **Annor and Badu (2019)**, **Chandan et al., (2018)**, **Abdulla et al., (2019)**, **Chaurasia, et al., (2020)**, **Lone et al., (2021)**, **Mekasha et al., (2022)**, **Subba et al., (2022)** and **Lalet et al., (2023)**. These hybrids may be commercialized as high yielding, bold seeded and quality rich maize hybrids after confirmation of their suitability for commercial cultivation.

Comment [I10]: and inbreeding

Comment [I11]: which DI

Comment [I12]: silking was ranged

Comment [I13]: revealed significant

Comment [I14]: four for

Comment [I15]: six for cob

Comment [I16]: for shelling

Comment [I17]: revealed significant

Comment [I18]: reference is missing

Comment [I19]: ref. missing

Comment [I20]: Mekasha et

Comment [I21]: and Lal et al.,

Comment [I22]: bold seeded and

Comment [I23]: suitability for

One of the characteristics of heterosis is that the increase in vigour is confined to  $F_1$  generation. There is considerable depression from  $F_1$  to  $F_2$  and later generation. Shull (1914) reported that high inbreeding depression (positive) is the reflection of higher heterosis especially in cross-pollinated crop like maize. It may be seen from the present study that hybrid combinations that showed higher estimates of heterosis in general found to show substantial inbreeding depression (Table 2). In maize, inbreeding is accompanied by a reduction in the mean phenotypic value of most of the traits of economic importance simply because of reduction in fitness.

Maximum inbreeding depression for days to 50% tasseling were observed from DI-122 x DI-109 (4.92), DI-109 x D-56 (4.00) for days to 50% silking, DI-122 x D-56 (2.26) for days to 75% dry husk, DI-109 x DI-117 (-4.78) for plant height, DI-121 x DI-118 (-37.30) for number of cobs per plant, DI-123 x D-52 (-10.62) for cob length (cm), DI-124 x DI-118 (-10.00) for cob diameter (cm), DI-122 x DI-124 (-44.47) for cob weight (g), DI-124 x D-56 (4.90) for number of grains per cob, DI-124 x DI-111 (4.01) for 100-kernel weight (g), DI-123 x DI-109 (23.08) for shelling (%), DI-111 x DI-118 (4.43) for lysine content (%), DI-123 x DI-109 (23.08) for tryptophan content (%), DI-122 x DI-109 (10.15) for protein content (%), and DI-111 x DI-118 (61.50) for grain yield per plant (g).

Comment [124]: for plant

These findings were reported earlier by Talukder *et al.*, (2016), Ilyaset *et al.*, (2020), Raj *et al.*, (2020) and Lone *et al.*, (2021).

### Conclusion

Among 45 crosses which generated from all possible combinations, eight revealed significant positive heterosis over economic parent for grain yield per plant (g) viz., DI-123 x DI-111 (91.11), DI-121 x DI-117 (80.82), DI-121 x DI-111 (80.33), DI-121 x D-56 (78.33), DI-123 x DI-117 (74.10), DI-122 x D-52 (70.23), DI-121 x D-52 (67.04) and DI-117 x D-56 (62.24). These crosses may be further used in maize improvement programme for the development of superior inbred lines.

### References

- Abdulla, S. S., Abdulkhaliq, D. A., and Towfiq, S. I. (2022). partial diallel analysis of maize inbred lines for kernel yield and its components in sulaimani-iraq. *The Iraqi Journal of Agricultural Science*, 53(5), 1190-1202.
- Annor, B., Badu-Apraku, B., Nyadanu, D., Akromah, R., and Fakorede, M. A. (2019). Testcross performance and combining ability of early maturing maize inbreds under multiple-stress environments. *Scientific Reports*, 9(1), 13809.

**Anonymous, (2022).** Agriculture – Statistical year book India (Ministry of statistics and programme implementation), [mospi.nic.in/statistical-year-book-india/2022](http://mospi.nic.in/statistical-year-book-india/2022).

**Chaurasia, N. K., Nirala, R. B. P., and Singh, B. (2020).** Combining ability and heterosis studies in maize (*Zea mays* L.) under kharif season. *International Journal of Current Microbiology and Applied Sciences*, 9(11): 2576-2586.

**Gerpacio, R. V., & Pingali, P. L. (2007).** Tropical and subtropical maize in Asia: production systems, constraints, and research priorities. *CIMMYT*.

**Griffing, B., (1956).** Concept of general combining ability and specific combining ability in relation to diallel crossing system. *Australian Journal of Biological Sciences* 9: 463-493.

**Griffing, B., and E. W. Lindstrom. (1954).** Study of the combining abilities of corn inbreds having varying properties of corn belt and non-corn belt germplasm. *Agron. J.* 46: 545–552.

**Ilyas, M., Khan S. A., Awan S. I., Rehman S., Khan M. R and Hafeez S. (2020)** Study of heterosis and inbreeding depression under natural and water stress conditions in diverse maize hybrids, *Sarhad Journal of Agriculture*, 36(1): 324-332.

**Keeble F, Pellew C, 1910.** The mode of inheritance of stature and of time of flowering. *J Genet* 1:47-56.

**Kiani, T. T., Hussain, M., and Rahman, H. (2015).** Heterosis and inbreeding depression for grain yield variables in indigenous maize germplasm. *Sarhad Journal of Agriculture*, 31(4), 217-223.

**Lone, B. A., Hamid, A., Ali, G., Fayaz, A., and Kumar, S. (2021).** Standard heterosis for grain yield and its attributing traits in early maturing maize hybrids, *International Journal of Plant & Soil Science*, 33(24) 418-421.

**Mekasha, G. M., Chere, A. T., and Woreti, D. N. (2022).** Mid-parent and better parent heterosis study on highland quality protein maize hybrids in Ethiopia. *International Journal of Plant & Soil Science*, 34(21), 226-248.

**Melchinger A.E. and Gumber R. K. (1998).** Overview of heterosis and heterotic groups in agronomic crops, In: Concepts and Breeding of Heterosis in Crop Plants, (ed. Lamkey and Staub). *CSSA Special Publications*, Madison, USA: 29-44.

**Raj, G. S., Marker, S., Scaria, S., Mohd, S., and Ansari, S. (2020).** Quantitative studies on heterosis and inbreeding depression in quality protein maize for terminal heat tolerance. *Journal of Pharmacognosy and Phytochemistry*, 9(2), 784-788.

**Rani, N., Prasad Nirala, R. B., Pal, A. K., and Ranjan, T. (2021).** Combining ability for quantitative traits under normal and heat stress conditions in maize (*Zea mays* L.). *Bangladesh Journal of Botany*, 50(3), 659–669.

**Romay, M. C., Millard, M. J., Glaubitz, J. C., Peiffer, J. A, and Swarts, K. L, (2013).** Comprehensive genotyping of the USA national maize inbred seed bank. *Genome Biol.* 14: R55.

**Singh, S. B., Gupta, B. B., and Singh, A. K. (2010).** Heterotic expression and combining ability analysis for yield and its components in maize (*Zea mays* L.) inbreds. *Progressive Agriculture*, 10(2), 275-281.

**Subba, V., Nath, A., Kundagrami, S., and Ghosh, A. (2022).** Study of combining ability and heterosis in quality protein maize using line x tester mating design. *Agricultural Science Digest-A Research Journal*, 42(2), 159-164.

**Talukder, M. Z. A., Karim, A. S., Ahmed, S., and Amiruzzaman, M. (2016).** Combining ability and heterosis on yield and its component traits in maize (*Zea mays* L.). *Bangladesh Journal of Agricultural Research*, 41(3), 565-577.

**Yadav, I., Sharma, V., Kumar, M., Yadav, L.P., Mishra, A., Singh, V., Singh Dhanda. P., Yadav, A., Yadav, M., Singh, S.K. and Kamaluddin, (2023).** Assessment of Gene Action and Identification of Heterotic Hybrids for Enhancing Yield in Field Pea. *Horticulturae*, 9(9):997.

UNDER PEER REVIEW

**Table 1: Heterosis and inbreeding depression for all 15 traits over check D 52 (p9)**

S.N	Hybrids	Days to 50% tasseling		Days to 50% silking		Days to 75% dry husk		Plant height (cm)		Number of cobs per plant		Cob length (cm)	
		EH	ID	EH	ID	EH	ID	EH	ID	EH	ID	EH	ID
1	DI-122 x DI- 121	-2.18	2.23	-4.24 *	-1.21	-4.42 *	-1.99	0.94	-0.76	39.85 **	10.75**	14.24 **	-1.56
2	DI-122 x DI- 124	-3.91	0.46	-5.79 **	-0.42	-6.01 **	-4.37	2.28	1.30	35.34 **	-0.17	21.29 **	2.55
3	DI-122 x DI- 123	-4.36 *	4.09	-5.79 **	-2.46	-3.8	-0.66	3.98	1.64	9.77 **	0.01	13.38 **	1.34
4	DI-122 x DI- 109	-3.05	4.92*	-6.56 **	-4.96*	-3.8	-0.99	0.94	2.06	39.85 **	3.23	10.48 **	2.33
5	DI-122 x DI- 111	-6.96 **	-1.40	-4.63 *	-3.25	-2.85	1.62	0.00	2.28	0.00	24.81**	14.50 **	5.99*
6	DI-122 x DI- 118	-5.22 *	2.30	-6.95**	-2.91	-2.53	-0.32	-1.90	1.74	0.00	5.26	18.85 **	6.43*
7	DI-122 x DI- 117	-3.91	2.71	-5.02 *	-2.04	-2.53	1.31	1.71	2.80	39.85 **	3.23	15.56 **	1.88
8	DI-122 x D- 52	-3.48	1.35	-4.24 *	-1.21	-1.9	-0.33	1.51	0.93	30.08 **	30.64**	21.29 **	11.25**
9	DI-122 x D- 56	-1.75	3.98*	-1.54	0.39	-2.21	2.26	3.22	7.72	24.81 **	0.01	14.50 **	3.45
10	DI-121 x DI-124	-9.13 **	-3.82	-2.32	-0.40	-1.9	1.94	-0.76	2.29	9.77 **	4.11	6.33 **	-0.43
11	DI-121 x DI-123	-8.27 **	0.94	-5.40 **	0.42	-6.65 **	-5.08*	-0.19	1.71	24.81 **	32.53**	14.90 **	4.02
12	DI-121 x DI-109	-6.52 **	0.001	-6.95 **	-1.24	-4.75 *	-5.95*	2.28	3.34	35.34 **	3.89	9.23 **	-1.99
13	DI-121 x DI-111	-9.13 **	-3.34	-8.87 **	-4.65*	-4.42 *	-5.97*	2.28	1.67	45.11 **	31.09**	12.26 **	0.01
14	DI-121 x DI-118	-6.52 **	0.47	-5.79 **	0.001	-3.8	-1.32	2.85	1.66	-5.26 *	-37.30**	15.38 **	2.86
15	DI-121 x DI-117	-7.83 **	-1.42	-5.79 **	0.81	-6.01 **	-2.70	-0.19	0.95	39.85 **	32.26**	18.00 **	4.47
16	DI-121 x D- 52	-8.27 **	-2.84	-5.40 **	-0.81	-4.11	-2.97	0.00	1.33	35.59 **	11.11**	19.12 **	4.26
17	DI-121 x D- 56	-9.13 **	-1.44	-2.7	0.39	-6.96**	-3.74	3.22	0.001	24.81 **	-8.43*	10.55 **	4.00
18	DI-124 x DI-123	-10.00 **	-4.83*	-3.48	-1.20	-6.01 **	-1.01	0.57	1.13	5.26 *	5.00	13.84 **	-1.91
19	DI-124 x DI-109	-10.88 **	-5.37**	-5.40 **	-0.81	-4.75 *	-2.33	-0.19	0.95	24.81 **	3.61	10.42 **	2.27
20	DI-124 x DI-111	-8.27 **	0.47	-6.56 **	-2.89	-4.42 *	-1.99	1.51	-2.24	35.59 **	18.89**	21.49 **	6.84
21	DI-124 x DI-118	-10.88 **	-8.30**	-3.48	1.20	-5.70 **	-2.36	3.79	0.001	35.34 **	7.78	10.94 **	-2.61
22	DI-124 x DI-117	-6.52 **	-2.32	-4.63 *	-1.63	-4.11	-0.99	3.98	1.64	9.77 **	-23.29**	14.50 **	3.68
23	DI-124 x D- 52	-9.13 **	-3.82	-5.79**	-2.88	-8.22**	-4.48*	2.65	-0.19	15.04 **	-17.65**	8.57 **	-5.65*
24	DI-124 x D- 56	-9.13 **	-1.44	-3.48	0.40	-6.65 **	-2.72	3.22	2.76	24.81 **	12.05**	1.32	-8.20**
25	DI- 123 X DI-109	-9.57 **	-0.49	-4.63 *	-2.84	-9.49 **	-6.29**	3.79	0.55	5.26 *	-28.57**	4.81 *	-2.33
26	DI- 123 X DI-111	-10.43 **	-6.31**	-4.24 *	-1.21	-7.60 **	-2.74	4.55 *	1.63	39.85 **	13.98**	4.61	0.88
27	DI- 123 X DI-118	-10.00**	-6.77**	-5.40 **	0.42	-7.91 **	-3.78	3.98	0.73	35.34 **	18.89**	7.91 **	4.28
28	DI- 123 X DI-117	-9.57 **	-1.44	-5.79 **	0.00	-8.54 **	-3.11	2.08	3.16	24.81 **	-8.43*	8.31 **	3.23
29	DI- 123 X D- 52	-12.61 **	-4.97*	-3.08	-1.20	-8.22 **	-3.79	3.79	0.001	35.34 **	0.00	-0.07	-10.62**
30	DI- 123 X D- 56	-10.43 **	0.001	-5.79 **	-0.42	-6.01 **	-0.33	-1.33	-1.93	24.81 **	19.88**	12.26 **	2.11

31	DI-109 X DI-111	-9.57 **	-2.88	-4.63 *	-1.63	-6.32 **	-2.70	3.22	4.41*	-9.77 **	5.83	4.35	0.63
32	DI-109 X DI-118	-12.18 **	-9.42**	-1.92	2.76	-2.85	1.30	4.36	2.00	39.85 **	10.75**	4.35	0.63
33	DI-109 X DI-117	-9.57 **	-4.33	-6.56 **	-2.48	-3.8	-0.66	-0.76	-4.78*	0.01	0.01	2.83	0.00
34	DI-109 X D-52	-9.13 **	-1.44	-5.79 **	1.64	-4.75 *	-1.99	2.65	6.65**	0.01	0.01	2.37	-0.64
35	DI-109 X D- 56	-10.88 **	-6.83**	-3.48	4.00	-4.75 *	-1.00	0.00	0.76	39.85 **	10.75**	3.03	1.47
36	DI-111 X DI- 118	-9.57 **	-1.93	-6.17 **	-0.41	-5.70 **	-2.69	-1.14	-0.96	35.34 **	26.11**	5.47 *	3.13
37	DI-111 X DI- 117	-12.18 **	-4.96*	-5.79 **	-2.46	-4.75 *	-1.66	-1.71	0.78	20.30 **	-8.12*	6.59 **	5.81*
38	DI-111 X D- 52	-10.00 **	-3.99	-4.63 *	2.43	-5.70 **	-1.35	1.71	3.92	9.77 **	-9.59**	1.98	0.45
39	DI-111 X D- 56	-10.88 **	-6.35**	-6.17 **	-1.23	-3.8	0.33	2.85	-0.37	5.26 *	5.00	8.57 **	5.10
40	DI-118 X DI- 117	-10.88 **	-4.39	-5.40 **	1.22	-2.85	-0.65	1.14	-2.81	35.34 **	11.11**	10.55 **	0.42
41	DI-118 X D- 52	-9.57 **	-2.88	-5.79 **	-2.46	-2.85	-2.61	3.42	4.03	39.85 **	6.99*	13.84 **	1.39
42	DI-118 X D- 56	-10.43 **	-1.46	-6.56 **	-4.13	-1.58	-0.64	3.98	5.83*	35.59 **	18.89**	14.70 **	5.75*
43	DI-117 X D-52	-7.83 **	-7.08**	-2.32	0.001	-3.47	-0.32	2.85	0.001	35.34 **	11.11**	17.53 **	2.58
44	DI-117 X D-56	-6.96 **	-4.68*	-3.08	-1.98	-3.47	0.001	4.55 *	1.27	39.85 **	3.23	16.48 **	2.49
45	D-52 X D -56	-10.00 **	-6.77**	-5.40 **	-4.08	-0.63	-1.59	4.36	0.55	39.85 **	10.75**	23.07 **	10.18**
	SE ±		1.24		0.99		1.55		2.19		0.11		0.45
	CD at 5%		2.46		1.96		3.08		4.34		0.22		0.89
	CD at 1%		3.03		2.42		3.79		5.35		0.27		1.10

Continue....

S.N	Hybrids	Cob diameter		Cob weight (g)		Number of grains per cobs		100 kernel weight (g)		Shelling (%)		Lycine content (%)	
		EH	ID	EH	ID	EH	ID	EH	ID	EH	ID	EH	ID
1	DI-122 × DI- 121	70.66 **	28.07**	-13.70 **	5.06*	-29.68 **	10.12**	7.43 **	2.56	5.39 *	3.80	10.33 **	4.68*
2	DI-122 × DI- 124	74.55 **	14.24**	-18.93 **	-44.47**	-28.82 **	18.90**	8.39 **	1.65	4.34	1.33	31.00 **	12.96**
3	DI-122 × DI- 123	46.71 **	10.82**	12.03 **	25.88**	9.26 **	10.63**	12.40 **	4.59*	6.97 **	3.02	11.44 **	5.96*
4	DI-122 × DI- 109	52.69 **	19.61**	-10.70 **	-18.66**	-33.83 **	10.45**	6.95 **	5.73*	2.90	0.89	21.03 **	5.49*
5	DI-122 × DI- 111	36.83 **	12.69**	-18.10 **	20.94**	-29.57 **	6.22*	4.20	1.25	8.71 **	5.29*	7.75 **	7.88**
6	DI-122 × DI- 118	29.94 **	2.53	-2.21	11.35**	9.77 **	12.70**	6.46 *	3.49	4.34	1.78	12.92 **	2.61
7	DI-122 × DI- 117	42.81 **	6.29*	2.36	13.10**	-28.29 **	1.84	4.05	5.29*	6.14 **	1.75	1.11	6.57*
8	DI-122 × D- 52	55.69 **	1.92	38.23 **	26.42**	2.75	19.44**	13.84 **	9.87**	10.65 **	1.25	11.81 **	3.96
9	DI-122 × D- 56	9.88 **	11.44**	13.73 **	9.35**	-25.36 **	0.35	13.99 **	4.91*	8.57 **	0.84	8.86 **	3.73
10	DI-121 × DI-124	34.73 **	8.89**	42.15 **	6.83*	21.65 **	2.15	2.75	4.37*	4.48	0.89	17.71 **	8.78**
11	DI-121 × DI-123	17.07 **	12.28**	48.25 **	12.86**	19.82 **	22.75**	6.46 *	1.54	5.59 *	1.75	4.06	2.13
12	DI-121 × DI-109	27.84 **	2.81	34.50 **	6.67*	21.82 **	12.82**	5.16 *	5.37*	2.90	0.01	11.07 **	2.33
13	DI-121 × DI-111	41.62 **	6.13*	58.31 **	14.22**	33.98 **	28.52**	5.64 *	3.20	4.98 *	0.01	14.76 **	4.82*

14	DI-121 × DI-118	37.72 **	6.96*	51.48 **	2.62	50.70 **	22.15**	3.71	3.58	4.42	0.01	14.02 **	2.27
15	DI-121 × DI-117	54.79 **	19.92**	53.44 **	9.53**	31.04 **	1.35	5.64 *	1.37	7.60 **	2.99	16.97 **	5.99*
16	DI-121 × D- 52	44.61 **	20.50**	65.66 **	9.98**	2.36	3.70	9.99 **	6.45*	5.81 *	0.01	18.82 **	11.18**
17	DI-121 × D- 56	24.85 **	5.28*	74.11 **	10.90**	8.60 **	24.97**	11.29 **	1.60	7.66 **	0.01	16.97 **	8.20**
18	DI-124 x DI-123	46.71 **	22.86**	22.17 **	11.29**	-1.00	16.63**	6.95 **	3.61	10.09 **	3.34	16.24 **	9.84**
19	DI-124 x DI-109	48.80 **	16.10**	14.98 **	14.99**	-32.61 **	10.83**	5.16 *	3.21	5.45 *	0.01	14.02 **	-1.94
20	DI-124 x DI-111	36.23 **	4.40	19.42 **	9.66**	-22.51 **	1.08	4.68	4.01	5.94 *	0.01	27.31 **	20.58**
21	DI-124 x DI-118	19.76 **	-10.00**	5.93 *	10.11**	-19.59 **	4.85	5.16 *	2.43	7.19 **	3.43	16.61 **	8.23**
22	DI-124 x DI-117	34.73 **	11.56**	20.73 **	14.82**	-24.03 **	7.32*	9.99 **	7.02**	11.25 **	1.24	0.37	-11.76**
23	DI-124 x D- 52	27.84 **	2.34	31.18 **	8.56**	-20.60 **	8.04*	12.25 **	5.59*	8.85 **	0.01	14.76 **	2.89
24	DI-124 x D- 56	51.20 **	27.13**	21.43 **	10.94**	-38.92 **	4.90	12.25 **	3.44	9.40 **	1.68	13.65 **	6.17*
25	DI-123 X DI-109	47.60 **	11.36**	37.23 **	11.52**	20.66 **	26.59**	7.57 **	2.11	2.77	0.01	22.88 **	6.31*
26	DI-123 X DI-111	55.09 **	14.48**	84.55 **	35.95**	6.74 **	14.61**	14.18 **	7.18**	4.29	0.01	36.53 **	21.89**
27	DI-123 X DI-118	58.68 **	22.64**	28.17 **	11.31**	14.68 **	10.96**	11.58 **	6.05*	2.21	0.01	13.28 **	6.19*
28	DI-123 X DI-117	42.81 **	13.42**	54.21 **	16.88**	20.33 **	26.70**	17.08 **	10.71**	3.73	0.01	22.51 **	9.04**
29	DI-123 X D- 52	31.74 **	6.82*	37.53 **	9.36**	-21.13 **	2.30	10.47 **	6.42*	3.87	0.01	15.13 **	6.41**
30	DI-123 X D- 56	27.25 **	4.24	41.41 **	11.39**	26.13 **	37.48**	15.63 **	6.01*	10.65 **	4.16	-18.08 **	-31.08**
31	DI-109 X DI-111	23.65 **	8.72**	40.06 **	20.70**	23.89 **	10.21**	16.11 **	5.15	5.03 *	3.08	19.56 **	4.63
32	DI-109 X DI-118	40.72 **	7.87*	54.87 **	17.73**	-23.58 **	9.40**	10.32 **	3.67	4.01	2.21	3.69	-0.36
33	DI-109 X DI-117	27.84 **	2.11	-5.08	-3.03	-3.42	12.99**	10.81 **	9.58**	11.12 **	8.01**	14.76 **	4.18
34	DI-109 X D-52	30.84 **	10.76**	24.31 **	-11.06**	-4.63 *	-7.43*	6.27 *	5.13*	5.59 *	2.62	7.38 **	3.09
35	DI-109 X D- 56	62.57 **	26.89**	71.41 **	6.55*	-10.09 **	2.06	7.91 **	3.13	2.77	0.01	23.62 **	11.04**
36	DI-111 X DI- 118	33.83 **	5.37*	-11.93 **	12.35**	-41.21 **	10.79**	10.81 **	5.22*	5.03 *	1.76	16.61 **	4.43
37	DI-111 X DI- 117	37.82 **	6.52*	-34.39 **	18.12**	-33.79 **	25.83**	8.06 **	8.04**	10.51 **	7.83**	4.43	2.83
38	DI-111 X D- 52	24.25 **	16.87**	-41.48 **	30.07**	-51.32 **	10.07**	4.82	3.68	7.19 **	1.75	14.76 **	8.04**
39	DI-111 X D- 56	41.62 **	18.60**	-34.91 **	18.72**	-34.50 **	18.11**	9.65 **	5.98*	8.85 **	1.27	12.55 **	6.89*
40	DI-118 X DI- 117	51.80 **	16.57**	65.32 **	56.31**	-5.40 *	34.61**	15.15 **	10.77**	4.48	0.01	18.82 **	11.18**
41	DI-118 X D- 52	55.69 **	19.23**	54.26 **	30.37**	-14.26 **	13.16**	6.27 *	3.63	7.33 **	2.58	4.80	4.58
42	DI-118 X D- 56	31.74 **	12.95**	57.52 **	47.84**	-10.15 **	42.03**	9.50 **	5.73*	7.66 **	3.52	15.13 **	6.09*
43	DI-117 X D-52	51.80 **	17.16**	35.16 **	8.34**	38.12 **	27.22**	7.91 **	2.24	9.59 **	1.69	19.56 **	10.19**
44	DI-117 X D-56	34.73 **	13.56**	40.96 **	9.27**	-3.74	20.87**	14.95 **	8.81**	8.57 **	2.55	13.28 **	4.56
45	D-52 X D -56	44.01 **	5.41*	38.05 **	8.45**	-10.46 **	11.54**	17.22 **	6.17*	10.65 **	3.32	4.06	4.96
	<b>SE ±</b>		<b>0.11</b>		<b>2.70</b>		<b>4.84</b>		<b>0.55</b>		<b>1.51</b>		<b>0.09</b>
	<b>CD at 5%</b>		<b>0.22</b>		<b>5.36</b>		<b>9.60</b>		<b>1.09</b>		<b>3.00</b>		<b>0.18</b>
	<b>CD at 1%</b>		<b>0.27</b>		<b>6.60</b>		<b>11.83</b>		<b>1.34</b>		<b>3.69</b>		<b>0.22</b>

Continue...

SN	Hybrids	Tryptophan content (%)		Protein content (%)		Grain yield/plant (g)	
		EH	ID	EH	ID	EH	ID
1	DI-122 × DI- 121	-8.02 **	11.76**	7.78 **	5.73*	-15.90 **	17.11**
2	DI-122 × DI- 124	35.14 **	4.00	7.04 **	4.61	-17.37 **	18.11**
3	DI-122 × DI- 123	16.22 **	6.98*	6.17 **	0.81	-0.31	9.74**
4	DI-122 × DI- 109	29.73 **	4.17	5.80 **	10.15**	-12.73 **	14.14**
5	DI-122 × DI- 111	13.51 **	7.14**	4.57 *	8.62**	-34.91 **	33.04**
6	DI-122 × DI- 118	10.81 **	7.32**	5.80 **	6.65*	-4.68 *	6.61
7	DI-122 × DI- 117	5.41 *	7.69**	4.07	5.93*	12.81 **	13.36**
8	DI-122 × D- 52	16.22 **	4.65*	4.94 *	0.82	70.23 **	59.94**
9	DI-122 × D- 56	13.51 **	4.76*	3.70	2.38	20.69 **	14.42**
10	DI-121 × DI-124	24.32 **	0.00	3.33	0.48	58.96 **	32.13**
11	DI-121 × DI-123	8.11 **	7.50**	3.33	8.36**	40.75 **	26.92**
12	DI-121 × DI-109	16.22 **	6.98**	4.07	1.30	2.53	-27.79**
13	DI-121 × DI-111	21.62 **	8.89**	3.70	6.79*	80.33 **	44.63**
14	DI-121 × DI-118	18.92 **	9.09**	4.57 *	0.59	59.50 **	20.55**
15	DI-121 × DI-117	10.81 **	14.63**	4.07	0.59	80.82 **	30.30**
16	DI-121 × D- 52	24.32 **	13.04**	2.47	0.36	67.04 **	7.27*
17	DI-121 × D- 56	27.03 **	12.77**	4.07	2.49	78.33 **	17.39**
18	DI-124 x DI-123	-13.51 **	6.25*	3.70	2.74	46.28 **	28.04**
19	DI-124 x DI-109	18.92 **	4.55	3.33	2.63	17.29 **	1.87
20	DI-124 x DI-111	32.43 **	14.29**	4.57 *	1.65	12.57 **	6.21
21	DI-124 x DI-118	10.81 **	17.07**	2.84	1.56	17.29 **	20.56**
22	DI-124 x DI-117	5.41 *	17.95**	3.33	2.87	5.69 **	5.88
23	DI-124 x D- 52	18.92 **	11.36**	4.07	1.19	1.28	10.35**
24	DI-124 x D- 56	2.70	18.42**	3.70	1.55	9.61 **	10.46**
25	DI- 123 X DI-109	5.41 *	23.08**	4.07	0.95	27.18 **	16.85**
26	DI- 123 X DI-111	10.81 **	14.63**	5.80 **	0.70	94.11 **	38.61**
27	DI- 123 X DI-118	18.92 **	6.82**	2.84	1.32	41.86 **	12.81**
28	DI- 123 X DI-117	27.03 **	8.51**	2.84	8.40**	74.10 **	32.70**
29	DI- 123 X D- 52	21.62 **	13.33**	1.60	0.73	38.89 **	11.03**
30	DI- 123 X D- 56	-13.51 **	18.75**	5.31 *	9.73**	49.67 **	42.03**
31	DI-109 X DI-111	24.41 **	4.35	6.17 **	0.35	29.43 **	39.07**
32	DI-109 X DI-118	8.11 **	5.00	3.33	1.67	14.80 **	18.26**
33	DI-109 X DI-117	18.92 **	4.55	4.94 *	0.82	-28.38 **	5.22
34	DI- 109 X D-52	13.60 **	9.52**	3.70	2.26	12.16 **	9.65**
35	DI-109 X D- 56	2.70	7.89**	4.20 *	1.66	44.52 **	6.92
36	DI-111 X DI- 118	13.51 **	11.90**	4.07	2.14	0.56	61.50**
37	DI-111 X DI- 117	8.11 **	22.50**	3.83	2.26	-29.58 **	25.93**
38	DI-111 X D- 52	18.92 **	9.09**	3.70	1.55	-41.27 **	8.77*
39	DI-111 X D- 56	18.92 **	13.64**	4.94 *	1.18	-32.91 **	12.59**
40	DI-118 X DI- 117	21.62 **	11.11**	4.32 *	0.59	32.22 **	51.24**
41	DI-118 X D- 52	10.81 **	9.76**	4.69 *	3.30	6.40 **	7.65
42	DI-118 X D- 56	21.62 **	8.89**	3.95	0.24	23.97 **	57.30**
43	DI-117 X D-52	10.81 **	12.20**	4.94 *	0.82	44.80 **	16.36**
44	DI-117 X D-56	18.92 **	11.36**	6.17 **	0.81	62.24 **	19.28**
45	D-52 X D -56	13.51 **	11.90**	4.57 *	4.01	24.84 **	14.69**
	SE ±		0.0110		0.084		2.44
	CD at 5%		0.0218		0.17		4.84
	CD at 1%		0.0269		0.21		5.97

EH= Economic Heterosis, ID= Inbreeding Depression