

# DETERMINATION OF NOVEL PLANT GROWTH PROMOTING BACTERIA *Stenotrophomonas maltophilia* ISOLATED FROM RHIZOSPHERIC SOILS OF SOUTH GUJARAT FOREST, INDIA

## ABSTRACT:

Forest regions have good weather conditions for agricultural practices. Crop growth and productivity are affected by the presence of novel bio organisms in these areas. The present study sought to identify the location of the forest rhizosphere soil of some field crops. Screening procedures were used to assess their growth capabilities and evaluate their potential as PGPB. Various features of PGPB, like phosphate and potassium solubilization, nitrogen fixation, indole-3-acetic acid (IAA) production, ammonia (NH<sub>3</sub>) synthesis, siderophore generation, and hydrogen cyanide (HCN) synthesis, were analyzed morphologically, biochemically, and physiologically during the screening process. One isolate out of the nine only produced good results for every attribute studied that promoted plant development. By sequencing their 16S rRNA, the promising isolates were determined to be *Stenotrophomonas maltophilia*. To assess the efficacy of PGPB and its liquid and powder formulation as a biofertilizer, a field trial had been conducted. In comparison to the uninoculated control, all tested inoculants considerably enhanced the yield components of several crops, the amount of N and P that plants could absorb from the soil, the activity of some soil enzymes, and the overall number of bacteria. The application of *Stenotrophomonas maltophilia* as a PGPR demonstrated significant promise in addressing the obstacles associated with sustainable agriculture in a variety of environmental circumstances.

**Keywords:** PGPB, soil enzymes, 16S rRNA sequencing, *Stenotrophomonas maltophilia*

## 1. INTRODUCTION:

Plant growth-promoting rhizobacteria (PGPR) are bacteria that inhabit the plant root surface and enhance plant growth. (Adedeji, Häggblom et al. 2020). The use of these microorganisms can diminish reliance on artificial pesticides and fertilizers, thus fostering sustainable agriculture (Patel, Kumar et al. 2021). PGPR can boost plant growth either directly or indirectly. Direct plant growth promotion involves mechanisms that supply plants with nitrogen (N), phosphorus (P), iron (Fe), and IAA, whereas indirect promotion involves the suppression of phytopathogens through antimicrobial metabolites as well as extracellular enzymes. (Chouyia, Romano et al. 2020, Wang, Solanki et al. 2023). A multitude of advantageous microorganisms, that includes bacteria as well as fungi, synthesize antibiotics as secondary metabolites that impede the growth as well as proliferation of phytopathogens by

altering their cell walls, membranes, or metabolic pathways. Beneficial bacteria can compete with phytopathogens for vital nutrients in the root zone. Plants exhibit heightened resistance to pathogens that induce systemic resistance. The proliferation and dissemination of infections can be halted by breaking the cell walls. The cell can undergo hydrolysis due to the secretion of extracellular proteases and lipases. (Kumari, Mallick et al. 2019, de Andrade, Santos et al. 2023).

Gram-negative *Stenotrophomonas* species are linked to a variety of environments, including both plant and animal hosts. (Hayward, Fegan, et al. 2010). Furthermore, this bacterium is global and omnipresent, existing in a wide range of environmental habitats, including harsh ones. It is fundamentally associated with the rhizosphere of plants, where it predominantly facilitates the elemental cycling of nitrogen and sulphur. It decomposes complex molecules and contaminants, promoting the growth and health of plants. (An and Berg 2018, Pérez-Pérez, Oudot et al. 2020).

The advantages of the genus *Stenotrophomonas* for plant systems have been documented by numerous researchers. The potential utility of *Stenotrophomonas* bacteria as efficient bio inoculants for enhancing plant development and controlling a number of food crop diseases is prompting more research into this bacterium. Biotechnological interest in this area is expanding (Ulrich, Kube et al. 2021). It also has the capability to protect itself from bacteria and fungi that induce plant diseases, endure biotic as well as abiotic stress, and demonstrate anti-quorum sensing as well as anti-biofilm capabilities. (Ulrich, Kube et al. 2021). *S. rhizophilia*, closely related, presents a viable alternative for biotechnological uses without compromising human health. (An and Berg 2018).

In the accumulation zone of the Dediypada forests in the Bharuch district of Ankleshwar, Gujarat, India, a new strain of bacteria was discovered. We carried out tests for plant growth capacity, 16s rRNA sequencing analysis (Sherpa, Das et al. 2021), and morphological and biochemical characterisation of the *Stenotrophomonas maltophilia* strain (John and Thangavel 2017).

## **2. MATERIALS AND METHODS:**

### **2.1 SITE DESCRIPTION**

Soil samples had been collected in the winter months from the Dediypada jungle in South Gujarat (Dediypada: 21.60452°N, 73.74450°E). Samples of soil were taken at a depth of 4 inches from the root zones of the soils. The adherent soil sample was shipped in less than

24 hours to Arihant Bio Science (India) Pvt Ltd.'s microbiological lab in Ankleshwar, Gujarat, India, where it was kept at a regulated 4°C. Despite this, it was found that the soil samples taken from the jungle environment had a pH of 7.3.

## **2.2 ISOLATION OF BACTERIA**

A forest plant's rhizosphere yielded a bacterium that was isolated and cultured for 48 hours at 37°C, pH 7, and standard nutritional agar media. The bacterial isolate was further described by its biochemical properties (Bergey's Manual of Systematic Bacteriology), morphological characteristics (Gram's staining), and cultural conditions.

## **2.3 MORPHOLOGICAL AND MICROSCOPIC OBSERVATION:**

The strain was cultivated and purified on various specialized media, such as Jensen's medium, nutritional medium, manual CRYEM medium, Azotobacter specific medium, PKVK medium, and Aleksandrow medium, at  $37 \pm 2$  °C for 48–72 hours in a BOD incubator. By using the plate streak approach, a single colony was produced. Under a microscope, the single colony's morphological features were examined.

Gram's staining for microscopic identification of bacterial isolates Each clean glass slide received a thick layer of strain, which was then heat dried for a short while by passing over a burner. Following the addition of one to three drops of crystal violet stain, the smear was allowed to rest at room temperature for one minute. After that, a mild tap water wash was used to get rid of the crystal violet stains. After applying one to three drops of gram's iodine, the smear was left to incubate for thirty to sixty seconds at room temperature. The smear was then gently cleaned with tap water after being decolorized with Gram's decolourizer and counter-stained with one to three drops of safranin. It was then incubated for thirty to sixty seconds at room temperature. After an overnight period at room temperature, the slide was examined under a light microscope.

## **2.4 BIOCHEMICAL CHARACTERIZATION**

The strain was screened for extra cellular enzymes such as lipase, oxidase, protease (Artha, Pramono et al. 2019), nitrate reductase (Kim and Seo 2018), starch hydrolysis (Hamza, Hussein et al. 2017), amylase (Artha, Pramono et al. 2019), urease, cellulose (Artha, Pramono et al. 2019), methyl red, Voges proskauer, gelatin hydrolysis, casein, hydrogen sulphide, citrate utilization (Hamza, Hussein et al. 2017) and catalase (Pervin, Jannat et al. 2017) was carried out in accordance with the guidelines (Reddy, Pradhan et al. 2010). According to the

procedures described by Liu, Tian et al. (2020), the BIOLOG system verified each and every biochemical test.

## 2.5 MOLECULAR IDENTIFICATION AND PHYLOGENY

The phenol-chloroform approach was utilized to recover the whole genomic DNA from the bacterium. Primer 27F/1492R was used to amplify a portion of the 16S rRNA gene. (Jo, Tagele et al. 2020). The isolated strain's 16S rRNA gene sequence was found using BLAST alignment, which was deduced from the NCBI public database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), and entered into the NCBI GenBank database. The study's homologous 16S rRNA gene sequences were obtained using NCBI BLAST searches. The evolutionary distances are measured in base substitutions per site and had been calculated by employing the Maximum Composite Likelihood technique. Eleven nucleotide sequences were examined. The 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and noncoding codon locations had been examined. All positions with incomplete data and gap were removed. The completed dataset had a total of 758 locations. Evolutionary analyses were conducted in MEGA6. (Saitou and Nei 1987, Dopazo 1994). All of the nucleotide sequences produced from this research were added to GenBank (NCBI) with the GeneBank accession number PQ192266.

## 2.6 PLANT GROWTH PROMOTING TRAITS


The recently isolated strain was evaluated in relation to several traits that promote plant growth, including the production of indole acetic acid (Khan, Ali et al. 2023), phosphate (Chen and Liu 2019), potassium (Hu, Chen et al. 2006), nitrogen fixation (Arsita, Karim et al. 2020), zinc (Fasim, Ahmed et al. 2002), organic acid (John and Thangavel 2017), ammonia (Sherman 1986), HCN (Castric and Castric 1983), siderophore (Xu, Wang et al. 2019), and ACC deaminase (John and Thangavel 2017, Ayaz, Ali et al. 2024, Saadaoui, Faize et al. 2024).

## 3. RESULTS AND DISCUSSION

### 3.1 ISOLATION AND BIOCHEMICAL CHARACTERIZATION OF BACTERIA

PGPR promotes plant development directly or indirectly by colonizing plant roots and lowering the presence of harmful bacteria. The rhizosphere is an ecological niche in which plant roots and soil microorganisms coexist and exchange nutrients. Thus, rhizobacteria help to improve soil fertility and sustainability. In this research, we investigated bacterial ecology in jungle plant rhizosphere. Soil samples were gathered from several fields in Dediypada forest. So many isolates were obtained using the serial dilution procedure. To evaluate the plant

growth promoting ability of the isolated strain, we streaked the plates in different specific mediums such as Nitrogen agar medium, Jensen agar medium, CRYEM agar medium, Aleksandrow agar medium, PKVK agar medium, GYC agar medium, Nutrient agar medium, and Calcium agar medium, as shown in **Figure 1**. Positive growth on all of the agar mediums listed above indicates that the isolated strain is capable of fixing atmospheric nitrogen, and solubilizing potassium, phosphate, and calcium.



**Figure 1:** primary screening for plant growth promoting traits A) Nitrogen free agar medium, B) Jensen agar medium, C) CRYEM agar medium, D) PKVK agar medium, E) GYC agar medium, F) Nutrient agar medium, G) Aleksandrow agar medium, H) Calcium agar medium, I) Species-specific medium.

**Table 1** presents the results of various biochemical tests performed in vitro conditions. The result shows that the isolated strain is gram's negative and shows positive results towards different enzyme production tests.

**Table 1: Biochemical characterization of bacteria**

| Sr. No. | Biochemical Test          | Strain   |
|---------|---------------------------|----------|
| 1       | Gram's stain              | Negative |
| 2       | Amylase production test   | +        |
| 3       | Protease production test  | +        |
| 4       | Lipase production test    | +        |
| 5       | Chitinase production test | +        |

### 3.2 MOLECULAR IDENTIFICATION BASED ON 16S rRNA GENE SEQUENCE

16S rRNA gene sequencing was used to identify the organisms based on their appearance, biochemical properties, and microscopy. Based on an identity criterion of at least 97% for the 16S rRNA gene sequencing result, it was concluded that the strain IAM 12423 isolate was closely related to the *Stenotrophomonas maltophilia* strain, with 99.21% ANI.

**Table 2** shows the identification of bacterial strains using 16S rRNA gene sequences.

**Table 2:** Identification of bacteria based on 16S rRNA gene sequencing

| Description  | Scientific Name                                    | Max Score | Total Score | Query Cover | E value | Per. ident | Acc. Len | Accession   |
|--|--|-----------|-------------|-------------|---------|------------|----------|-------------|
| Stenotrophomonas maltophilia strain IAM 12423 16S ribosomal RNA, partial sequence      | Stenotrophomonas maltophilia                       | 1369      | 1369        | 100%        | 0       | 99.21      | 1538     | NR_041577.1 |
| Stenotrophomonas maltophilia strain NBRC 14161 16S ribosomal RNA, partial sequence     | Stenotrophomonas maltophilia                       | 1365      | 1365        | 100%        | 0       | 99.08      | 1470     | NR_113648.1 |
| Stenotrophomonas maltophilia strain LMG 958 16S ribosomal RNA, partial sequence        | Stenotrophomonas maltophilia                       | 1365      | 1365        | 100%        | 0       | 99.08      | 1500     | NR_119220.1 |
| Stenotrophomonas maltophilia strain ATCC 13637 16S ribosomal RNA, partial sequence     | Stenotrophomonas maltophilia                       | 1363      | 1363        | 100%        | 0       | 99.08      | 1467     | NR_112030.1 |
| Stenotrophomonas pavanii strain LMG 25348 16S ribosomal RNA, partial sequence          | Stenotrophomonas pavanii                           | 1349      | 1349        | 100%        | 0       | 98.68      | 1497     | NR_118008.1 |
| Stenotrophomonas pavanii strain ICB 89 16S ribosomal RNA, partial sequence             | Stenotrophomonas pavanii                           | 1347      | 1347        | 100%        | 0       | 98.68      | 1483     | NR_116793.1 |
| Stenotrophomonas maltophilia strain ATCC 19861 16S ribosomal RNA, partial sequence     | Stenotrophomonas maltophilia                       | 1341      | 1341        | 100%        | 0       | 98.55      | 1517     | NR_040804.1 |
| [Pseudomonas] hibiscicola strain ATCC 19867 16S ribosomal RNA, partial sequence        | [Pseudomonas] hibiscicola                          | 1341      | 1341        | 100%        | 0       | 98.42      | 1519     | NR_024709.1 |
| Stenotrophomonas geniculata ATCC 19374 = JCM 13324 16S ribosomal RNA, partial sequence | Stenotrophomonas geniculata ATCC 19374 = JCM 13324 | 1319      | 1319        | 100%        | 0       | 97.89      | 1497     | NR_024708.1 |
| Stenotrophomonas cyclobalanopsidis strain TPQG1-4 16S ribosomal RNA, partial sequence  | Stenotrophomonas cyclobalanopsidis                 | 1301      | 1301        | 99%         | 0       | 97.63      | 1439     | NR_180613.1 |

The gene sequences of the isolated bacterial strain were submitted to NCBI GenBank under the accession code PQ192266. A phylogenetic tree of the isolated strain and its relatives was created (**Figure 2**).





Fig.2. Phylogenetic tree of the isolated strain and its relatives

**Figure 2** shows a phylogenetic tree of strains among *Stenotrophomonas* species and related taxa. The Neighbour-Joining approach was used to infer evolutionary history. The bootstrap consensus tree constructed from 500 repetitions (Felsenstein 1985) is used to illustrate the taxa's evolutionary history. Branches associated with partitions occurring in fewer than 50% of bootstrap replicates are collapsed. The percentage of duplicate trees in which the relevant taxa clustered together during the bootstrap test (500 repetitions) is given next to the branches (Felsenstein 1985). The evolutionary distances had been calculated by employing the Maximum Composite Likelihood technique and are shown as the number of base substitutions per site (Tamura, Nei et al. 2004). This study comprised eleven nucleotide sequences. All not known positions have been eliminated from each sequence pair employing the pairwise deletion method. The final dataset comprised 762 locations. MEGA11 was employed to do evolutionary analysis. (Tamura, Stecher et al. 2021). The findings are consistent with previous research and indicate that the genus *Stenotrophomonas* has the ability to promote plant development.

### 3.3 PLANT GROWTH-PROMOTING TRAITS

Figure 3 depicts the findings of PGP traits for the *Stenotrophomonas maltophilia* strain. *Stenotrophomonas maltophilia* strain produced ammonia ( $\text{NH}_3$ ), IAA, hydrogen cyanide (HCN), siderophore, fluorescent, and organic acid. Masclaux-Daubresse, Daniel-Vedele et al. (2010) state that ammonia build-up in the soil may result in alkaline conditions that inhibit the growth of a number of plant diseases. Since ammonia production encourages the growth of roots, shoots, and biomass, which speeds up plant growth, it is crucial for the accumulation of nitrogen. The PGPB produces a lot of indole acetic acid because it facilitates nutrient absorption and root growth. Keswani, Singh et al. (2020) state that IAA, which is generated by bacteria, is one of the most significant auxins in the regulation of plant vegetative development. By synthesizing some antibiotics or enzymes that break down cell walls, the production of hydrogen cyanide (HCN) as a secondary metabolite inhibits the growth of the majority of fungal phytopathogens, according to Kumar, Bahadur et al. (2015). Through its effect on the mobilization of elements from rock-forming minerals and its indirect promotion of plant development, hydrogen cyanide also improved the availability of nutrients. According to research, some plant growth-promoting bacteria (PGPB) may remove iron from the soil by producing low-molecular-weight compounds (400–1500 Da) (Shanmugaiah, Nithya et al. 2015). Siderophores are chemicals that bind to iron and enhance their bioavailability in plants

(Goswami, Thakker et al. 2016). Good bacteria in the soil and on plants produce siderophores, which are a crucial biological control mechanism because they prevent detrimental plant diseases from getting iron supplies by out-competing them (Khoshru, Mitra et al. 2020).



**Fig. 3.** PGP traits for the *Stenotrophomonas maltophilia* strain

**Figure 3:** Shows the Plant growth-promoting activities like I)  $\text{NH}_3$  production II) IAA production III) HCN production IV) Siderophore production V) Fluorescent production VI) Organic acid production VII) Bacterial growth curve VIII) PCR of strain.

Diverse plant growth-promoting bacteria (PGPB) frequently have numerous growth-stimulating characteristics, each demonstrating distinct activities under varying environmental and soil conditions. (Olanrewaju, Glick et al. 2017). Phosphate and zinc solubilization, together with IAA, HCN, and exopolysaccharide production, are but a subset of the several characteristics exhibited by PGPB that facilitate plant growth. (Kalam, Basu et al. 2020). The research conducted by Zhao, Ding et al. (2024) indicates that *Stenotrophomonas* strains exhibit remarkable environmental adaptability and have great promise for use as plant growth-promoting bacteria. Alexander, Singh et al. (2019) suggest that *S. maltophilia* requires more investigation as a PGPR for crop growth in  $\text{N}_2$  deficient environments. *Stenotrophomonas maltophilia* AVP 27 is a potential plant growth-

promoting rhizobacteria with a wide range of pathways, according to [Kumar and Audipudi \(2015\)](#). Our isolated strain may be used in the field to enhance plant development and replace nutrients since it exhibits several actions that promote plant growth.

#### 4. CONCLUSION

We isolated *Stenotrophomonas maltophilia* from the rhizospheric soils of south Gujarat forest for this study. 16s rRNA sequencing and bioinformatics analysis were used to confirm the isolated species. The ability of the isolated species to develop plants was examined in vitro using a variety of biochemical assays and plant growth activators. The findings of this investigation suggests that the isolated strain in the present study should be evaluated for its capacity to promote plant growth in unsterilized soil as well as in the field. Although beneficial rhizobacteria have a well-established effect on improving plant development, more research is still needed to fully understand the physiological and molecular processes underlying these microbes' improved utilization. However, mechanistic studies involving PGPR and plants are still required to explore the mechanisms by which PGPR promotes plant growth.

#### COMPETING INTERESTS

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

#### Disclaimer (Artificial intelligence)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. QuillBot free version for editing of manuscript
- 2.
- 3.

#### REFERENCES:

- Adedeji, A. A., et al. (2020). "Sustainable agriculture in Africa: Plant growth-promoting rhizobacteria (PGPR) to the rescue." *Scientific African* 9: e00492.
- Alexander, A., et al. (2019). "Plant growth promoting rhizobacterium *Stenotrophomonas maltophilia* BJ01 augments endurance against N<sub>2</sub> starvation by modulating physiology and biochemical activities of *Arachis hypogea*." *Plos one* 14(9): e0222405.
- An, S.-q. and G. Berg (2018). "*Stenotrophomonas maltophilia*." *Trends in microbiology* 26(7): 637-638.
- Arsita, R., et al. (2020). Isolation and identification of nitrogen-fixing bacteria in the corn rhizosphere (*Zea mays* L.) originating from Jeneponto Regency, South Sulawesi. IOP Conference Series: Earth and Environmental Science, IOP Publishing.
- Artha, O., et al. (2019). Identification of extracellular enzyme-producing bacteria (proteolytic, cellulolytic, and amylolytic) in the sediment of extensive ponds in Tanggulejo, Gresik. IOP Conference Series: Earth and Environmental Science, IOP Publishing.
- Ayaz, M., et al. (2024). "Exploring plant growth promoting traits and biocontrol potential of new isolated *Bacillus subtilis* BS-2301 strain in suppressing *Sclerotinia sclerotiorum* through various mechanisms." *Frontiers in Plant Science* 15: 1444328.
- Castric, K. F. and P. A. Castric (1983). "Method for rapid detection of cyanogenic bacteria." *Applied and Environmental Microbiology* 45(2): 701-702.
- Chen, Q. and S. Liu (2019). "Identification and characterization of the phosphate-solubilizing bacterium *Pantoea* sp. S32 in reclamation soil in Shanxi, China." *Frontiers in microbiology* 10: 2171.
- Chouyia, F. E., et al. (2020). "P-solubilizing *Streptomyces roseocinereus* MS1B15 with multiple plant growth-promoting traits enhance barley development and regulate rhizosphere microbial population." *Frontiers in Plant Science* 11: 1137.

- De Andrade, L. A., et al. (2023). "Plant growth-promoting rhizobacteria for sustainable agricultural production." *Microorganisms* 11(4): 1088.
- Dopazo, J. (1994). "Estimating errors and confidence intervals for branch lengths in phylogenetic trees by a bootstrap approach." *Journal of Molecular Evolution* 38: 300-304.
- Fasim, F., et al. (2002). "Solubilization of zinc salts by a bacterium isolated from the air environment of a tannery." *FEMS microbiology letters* 213(1): 1-6.
- Felsenstein, J. (1985). "Phylogenies and the comparative method." *The American Naturalist* 125(1): 1-15.
- Goswami, D., et al. (2016). "Portraying mechanics of plant growth promoting rhizobacteria (PGPR): a review." *Cogent Food & Agriculture* 2(1): 1127500.
- Hamza, T. A., et al. (2017). "Isolation and characterization of nitrogen fixing bacteria from rhizosphere soil collected from Shell Mele agricultural center, southern Ethiopia." *J Agric Sci Food Technol* 3(7): 117-124.
- Hayward, A., et al. (2010). "Stenotrophomonas and Lysobacter: ubiquitous plant-associated gamma-proteobacteria of developing significance in applied microbiology." *Journal of applied microbiology* 108(3): 756-770.
- Hu, X., et al. (2006). "Two phosphate-and potassium-solubilizing bacteria isolated from Tianmu Mountain, Zhejiang, China." *World journal of Microbiology and Biotechnology* 22(9): 983-990.
- Jo, H., et al. (2020). "Response of soil bacterial community and pepper plant growth to application of *Bacillus thuringiensis* KNU-07." *Agronomy* 10(4): 551.
- John, N. and M. Thangavel (2017). "*Stenotrophomonas maltophilia*: a novel plant growth promoter and biocontrol agent from marine environment." *Int. J. Adv. Res* 5: 207-214.

- Kalam, S., et al. (2020). "Functional and molecular characterization of plant growth promoting *Bacillus* isolates from tomato rhizosphere." *Heliyon* 6(8).
- Keswani, C., et al. (2020). "Auxins of microbial origin and their use in agriculture." *Applied Microbiology and Biotechnology* 104: 8549-8565.
- Khan, A. R., et al. (2023). "Plant–Microbes Interaction: Exploring the Impact of Cold-Tolerant *Bacillus* Strains RJGP41 and GBAC46 Volatiles on Tomato Growth Promotion through Different Mechanisms." *Biology* 12(7): 940.
- Khoshru, B., et al. (2020). "Current scenario and future prospects of plant growth-promoting rhizobacteria: an economic valuable resource for the agriculture revival under stressful conditions." *Journal of Plant Nutrition* 43(20): 3062-3092.
- Kim, J. Y. and H. S. Seo (2018). "In vitro nitrate reductase activity assay from *Arabidopsis* crude extracts." *Bio-protocol* 8(7): e2785-e2785.
- Kumar, A., et al. (2015). "Does a plant growth-promoting rhizobacteria enhance agricultural sustainability." *J Pure Appl Microbiol* 9(1): 715-724.
- Kumar, N. P. and A. V. Audipudi (2015). "Exploration of a novel plant growth promoting bacteria *Stenotrophomonas maltophilia* AVP27 isolated from the chilli rhizosphere soil." *International Journal of Engineering Research and General Science* 3(1): 265-273.
- Kumari, B., et al. (2019). "Plant growth promoting rhizobacteria (PGPR): modern prospects for sustainable agriculture." *Plant Health Under Biotic Stress: Volume 2: Microbial Interactions*: 109-127.
- Liu, Q., et al. (2020). "Cryobacterium ruanii sp. nov. and Cryobacterium breve sp. nov., isolated from glaciers." *International Journal of Systematic and Evolutionary Microbiology* 70(3): 1918-1923.

- Masclaux-Daubresse, C., et al. (2010). "Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture." *Annals of botany* 105(7): 1141-1157.
- Olanrewaju, O. S., et al. (2017). "Mechanisms of action of plant growth promoting bacteria." *World journal of Microbiology and Biotechnology* 33: 1-16.
- Patel, J. S., et al. (2021). PGPR formulations and application in the management of pulse crop health. *Biofertilizers*, Elsevier: 239-251.
- Pérez-Pérez, R., et al. (2020). "Isolation and characterization of *Stenotrophomonas* associated to maize (*Zea Mays* L.) rhizosphere."
- Pervin, S., et al. (2017). "Characterization of Rhizobia from Root Nodule and Rhizosphere of *Lablab purpureus* and *Vigna sinensis* in Bangladesh." *Turkish Journal of Agriculture-Food Science and Technology* 5(1): 14-17.
- Reddy, G., et al. (2010). "Cryobacterium *Pindariense* sp. nov., a psychrophilic bacterium from a Himalayan glacier." *Int J Syst Evol Microbiol* 60: 866-870.
- Saadaoui, M., et al. (2024). "Evaluation of Tunisian wheat endophytes as plant growth promoting bacteria and biological control agents against *Fusarium culmorum*." *Plos one* 19(5): e0300791.
- Saitou, N. and M. Nei (1987). "The neighbor-joining method: a new method for reconstructing phylogenetic trees." *Molecular biology and evolution* 4(4): 406-425.
- Shanmugiah, V., et al. (2015). "Biocontrol mechanisms of siderophores against bacterial plant pathogens." *Sustainable approaches to controlling plant pathogenic bacteria*: 167-190.
- Sherman, N. (1986). *Microbiology, a Laboratory Manual*, Reading, Mass.: Addison-Wesley Publishing Company.

- Sherpa, M. T., et al. (2021). "Draft genome sequence of *Stenotrophomonas maltophilia* strain P13 gives insight into its protease production and assessment of sulfur and nitrogen metabolism." *Current Research in Microbial Sciences* 2: 100012.
- Tamura, K., et al. (2004). "Prospects for inferring very large phylogenies by using the neighbor-joining method." *Proceedings of the National Academy of Sciences* 101(30): 11030-11035.
- Tamura, K., et al. (2021). "MEGA11: molecular evolutionary genetics analysis version 11." *Molecular biology and evolution* 38(7): 3022-3027.
- Ulrich, K., et al. (2021). "Genomic analysis of the endophytic *Stenotrophomonas* strain 169 reveals features related to plant-growth promotion and stress tolerance." *Frontiers in microbiology* 12: 687463.
- Wang, Z., et al. (2023). "Promoting plant resilience against stress by engineering root microenvironment with *Streptomyces* inoculants." *Microbiological Research* 277: 127509.
- Xu, W., et al. (2019). "Diversity of cultivable endophytic bacteria in mulberry and their potential for antimicrobial and plant growth-promoting activities." *Microbiological Research* 229: 126328.
- Zhao, Y., et al. (2024). "A comprehensive comparative genomic analysis revealed that plant growth promoting traits are ubiquitous in strains of *Stenotrophomonas*." *Frontiers in microbiology* 15: 1395477.