

Original Research Article

***In vitro* bioactivity guided Isolation and Partial Characterization of Stigmasterol from *Plectranthus esculentus*. N.E.Br. Crude Extract.**

ABSTRACT

Aim: *The study was aimed at the in-vitro antibacterial activity evaluation and isolation of phytocompound(s) from *Plectranthus esculentus* crude extract.*

Place and Duration: *This research work was carried out at Gombe State University in the Department of Chemistry for a period of one and half year which involved sample collection, preparation, purification and analysis of isolate.*

Methodology: *The coarsely powdered plant tuber was extracted with methanol. The filtrate obtained was concentrated on a rotavapor. The crude extract was re-dissolved in water: methanol (9:1) and partitioned with n-hexane, ethyl acetate and n-butanol respectively. The crude fractions were tested against a panel of microbes namely *Helicobacter pylori*, *Shigella flexneri*, and *Salmonella typhi*, using macro-dilution technique. The most active and sufficient n-hexane fraction was purified on column chromatography.*

Results: *The crude extract (≈172 g) was obtained from 3.1 kg of the stem bark sample. Fractionation of the crude extract yielded 18.7g, 13.6 g and 26.4 g for n-hexane, ethylacetate and n-butanol respectively. All the crude fractions displayed moderate to strong activity against the microbes. The n-hexane and ethyl acetate fractions were the most active fractions with MICs ranging from 62.5 - 150 µg/mL where *Salmonella typhi* and *Helicobacter pylori* were the most susceptible with MIC of 62.50 µg/mL. The n-butanol fraction was weaker with MIC ranging from 125 - 250 µg/mL. Gradient elution of the n-hexane fraction yielded a phytocompound coded C5 with R_f value of 0.56 in n-hexane: ethyl acetate (6:4) and melting point of 136-138°C.*

Characterization of C5 using spectroscopic data from FT-IR, GC-MS, ¹H NMR and compared with literature identified C5 as stigmasterol.

Conclusion: The study had shown that *Plectranthus esculentus* contains bioactive principles and may serve as a potential source of antibacterial agents.

Key words: *Plectranthus esculentus*, antibacterial agent, column chromatography, characterization, stigmasterol, phytocompound

1. INTRODUCTION

Natural products of plant origin had been used by man for centuries as a source of medicines for healthcare [1]. It is believed that more than eighty (80%) percent of the World population depends on medicinal plants for cultural reasons, affordability and the renewed interest in natural products out of safety concerns for synthetic drugs [2, 3]. These plants contain bioactive principles such as alkaloids, flavonoids, glycosides, saponins, terpenoids, polyphenols, polyketides and tannins with potential pharmacological properties such as antimicrobial, anticancer, antioxidant, anti-inflammation, antinociceptive and anthelmintic to mention just a few [4, 5].

The adoption of interdisciplinary research and technological advances has facilitated the successful isolation and characterization of single chemical entities (SCEs) with defined pharmacological activities at tolerable doses from complex bio-matrices [6, 7]. Many medicines and drugs contain active ingredients that originated from plant sources. These include vinblastine and vincristine from *Catharanthus roseus* used in the treatment of cancer [8, 9], artemisinin from

Artemisia annua used in the treatment of malaria and morphine from *Papaver somniferum* used as a pain reliever [10].

Plectranthus esculentus is a dicotyledonous plant from the family Lamiaceae commonly known as living-stone potato or "rizga" (kaffir) in Hausa can be found in the Nigerian middle belt region, Southern Africa and Zimbabwe [11, 12]. It is used ethnomedicine to treat stomachache, cancer, backache, worms and skin disorders [13].

The FT-IR analysis of *Plectranthus esculentus* tuberous root acetone extract revealed the presence of organic chromophores such as organic hydroxyl (-OH), amines, carbonyl, alkyl, and aromatic functional groups while the HPLC analysis revealed the presence of kaempferol and vanillic acids. These results confirm the presence of medicinally important phytochemical substances in *Plectranthus esculentus* extracts. Most of these phytochemicals have demonstrated pharmacological properties such as antifungal, antidiabetic anticancer and antimicrobial activities [13, 14]. Balogun *et al.* [15] reported the isolation and characterization of β -sitosterol and oleanolic acids from *Plectranthus esculentus* leaf extract while 19-dehydrousolic acid, yarumic acid and β -sitosterol were obtained from its tuberous root extract. Despite the widespread use of *Plectranthus esculentus* as food and phytomedicine, reports on isolation and characterization of its phytocompounds are quite scarce. Hence, we report for the first time the isolation and characterization of stigmasterol from *Plectranthus esculentus* root tuber methanol crude extract based on available literature.

2. MATERIAS AND METHODS

2.1. Sample collection and identification

The plant *Plectranthus esculentus* (tubers) was collected from Zawan Du Jos South LGA of Plateau State Nigeria in December, 2021. The plant was identified by Dr. Zainab Abubakar Department of Botany Gombe State University, Nigeria.

2.2 Preparation and Extraction plant material

The tubers were sliced; air-dried and coarsely powdered using a motorized miller. The powdered *Plectranthus esculentus* tubers (3.1 kg) was extracted with 10 L of methanol at room temperature using the maceration method with occasional shaking for seven (7) days. Methanol is quite useful as a solvent because it facilitates the extraction of both polar and non-polar organic constituents and can be easily removed from the crude extract due to its low molecular weight; also methanol containing water below 20% acts as preservative and does not allow the growth of mold during extended extraction periods [16].

The extract was filtered using Whatman No. 1 filter paper and then concentrated under reduced pressure with the aid of a rotary evaporator (BÜCHI R-100, Switzerland) at 40°C. The crude extract obtained was dried to a constant weight in a desiccator and kept in a cool and dry container until required for use.

2.3 Antibacterial activity

The antibacterial activity of *Plectranthus esculentus* crude extract fractions was evaluated using the broth dilution method reported by Kwaji *et al.* [17] with slight modifications. The bacterial strains employed were *Helicobacter pylori*, *Salmonella typhi* and *Shigella flexneri*. Nine test tubes containing two-fold serial dilutions of extracts or commercial antibiotic (Gentamicin) were made from the stock concentration to obtain final concentrations in the range of 500 µg/mL to 3.9 µg/mL. The ninth test tube (No. 9) served as negative control (broth + bacterium inoculum). Each test tube got culture medium (2 mL) + plant

extract or Gentamicin (1.8 mL) + 0.2 mL standardized bacterial inoculums (1×10^6 cfu/mL). The test tubes were covered with sterile aluminium foil and incubated at 37°C for 24 hours. Test tubes were observed for turbidity or growth after 18-24 hrs period. The lowest concentration that showed no turbidity or growth in the test tube was recorded as the minimum inhibitory concentration (MIC). Tests were carried out in triplicate.

2.4 Isolation of compounds from *Plectranthus esculentus* crude extract

Plectranthus esculentus n-hexane crude extract fraction (10 g) initially dissolved in methanol sufficient to form solution was adsorbed to 20 g of activated silica gel 60 (70-230 mesh) and then allowed to dry. Column packing was effected using the wet slurry method. Gradient elution was performed with hexane/ethyl acetate and ethyl acetate/methanol at 5% increase in volume (100:00), (95:05), (90:10), (85:15), (80:20), (75:25), etc. Eluents of column fractions were collected in batches of 100 mL. Each column fraction was collected in a separate conical flask and numbered serially. All column fractions were concentrated and combined based on their thin layer chromatography (TLC) profile. Finally, the isolate C5 which showed a single spot was washed and recrystallized from methanol. This involves the use of suitable solvent in which the isolate is insoluble while the impurities are soluble at room temperature [17].

3. RESULTS AND DISCUSSIONS

The crude extract of *Plectranthus esculentus* displayed significant to moderate activity against the tested bacterial isolates (Table 1). According to Teke et al. [18] and Demgne et al. (20), MIC values of 24.4 to 78.2 $\mu\text{g/mL}$ are reckoned as significant (i.e. $\text{MIC} < 100 \mu\text{g/mL}$), while $100 < \text{MIC} = 625 \mu\text{g/mL}$ are moderate or inactive for $\text{MIC} > 625 \mu\text{g/mL}$ for both bacterial and fungal pathogens. Based on this classification and also relative to the gentamicin standard, the *Plectranthus esculentus* crude extract fractions displayed significant activity against *H. pylori*

and *Salmonella typhi* and hence could be used to control infections associated with these organisms. The hexane and ethyl acetate fractions were identified as the most active fractions with MICs ranging from 62.5 - 125 µg/mL on all the microbes. This may partly justify the use of *Plectranthus esculentus* in ethnomedicine [19].

Table 1: Showing the minimum inhibitory concentration (MIC) of *Plectranthus esculentus* crude methanol extract fractions and the gentamicin antibiotic as control..

Bacteria	n-Hexane MIC (µg/mL)	Ethyl acetate MIC (µg/mL)	n-Butanol MIC (µg/mL)	Gentamicin MIC (µg/mL)
<i>H. pylori</i>	62.5	62.5	125	31.25
<i>Shigella flexneri</i>	125	125	250	31.25
<i>Salmonella typhi</i>	62.5	62.5	125	15.5

3.1 Characterization of *Plectranthus esculentus* Isolate (C5).

The FT-IR spectrum of C5 (Fig.1) revealed the presence several functional groups as presented in Table 2. These absorption frequencies align with those of sterol based on literature reports [21].




Fig. 1: FT-IR spectrum of C5

Table 2 shows the peaks or the functional groups frequencies of vibration

Table 2: FT-IR Spectrum frequencies of C5

SN	Frequency (cm ⁻¹)	Type of Vibration
1	3458	O-H stretching
2	2938	C-H stretching.(CH ₃)
4	2867	C-H stretching (CH ₂)
5	1639	C=C stretching
6	1465	CH ₃ bending
7	1383	CH ₂ bending
8	1133	CH ₂ bending
9	800	CH bending




Fig. 3: GC-MS spectrum of C5

The GC-MS analysis indicated that the isolated compound C5 is Stigmasterol as suggested by GC-MS library (Fig.3) with 99% hit. The fragmentation pattern is in agreement with literature [18]. The mass spectrum of the isolated compound showed a parent molecular ion [M⁺] peak at m/z 412 amu (Fig.3). The base peak m/z is 55(100). The percentage fragment of other peaks relative to the base peak are m/z - 412(12), 381(6), 351(6), 329(10), 300(7), 271(12), 255(16), 206(12), 145.1(20), 83(40) with the molecular formula C₂₉H₄₈O (Fig. 3).

The activity guided isolation and characterization of *Plectranthus esculentus* root tuber crude extract led to the identification of stigmasterol (Fig. 4). The root tubers of *Plectranthus esculentus* can be considered as a natural reservoir for stigmasterol in addition to other plant sources. Stigmasterol isolated from *Psychotria sycophylla* and tested against *Enterobacter aerogenes* EA27 yielded an MIC value of 64 µg/mL. The antibacterial was attributed to the

presence of hydroxyl (-OH) at position 3 on the chemical structure of stigmasterol (Fig.4) which in solution provides an acidic environment that is both inhibitory and bactericidal to pathogens depending on the concentration [20]. It is important to note that glycosylation of the stigmasterol hydroxyl group led to loss of antimicrobial activity [22]. The pharmacological activities of stigmasterol such as antimicrobial, antidiabetic, anticancer, anti-inflammatory, antioxidant, anti-osteoarthritis and neuroprotective activities are well documented [23, 24].




Fig. 4: Chemical Structure of Stigmasterol.

3.2: CONCLUSION

The evaluation of the antibacterial activity of *Plectranthus esculentus* revealed that the n-hexane and ethylacetate fractions exhibited strong activity against *Salmonella typhi* and *Helicobacter pylori* with MICs of 62.50 $\mu\text{g/mL}$. The observed strong antibacterial activities could be as a result of the ability of the nonpolar constituents of n-hexane and ethylacetate fractions to diffuse faster across the bacterial lipophilic cell membrane than the predominantly polar constituents of

n-butanol fraction. The n-hexane fraction which was of higher amount than either the ethylacetate and the n-butanol fraction was purified on column chromatography to obtain stigmasterol. The isolation and characterization of stigmasterol demonstrates the presence of bioactive compounds in the root tubers of *Plectranthus esculentus* and validates its use in ethnomedicine. It is recommended that the ethylacetate which demonstrated similar activity to that of the n-hexane fraction should also be investigated for the presence of bioactive compounds.

ACKNOWLEDGEMENT

The authors wish to acknowledge the support of the Technologists in the Department of Chemistry and the Gombe State University Management for creating a Research enabling environment.

CONFLICT OF INTEREST

Authors wish to declare no conflict of exists.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

- 1). Ahmad, S. R. and Karmakar, S. The Role of Medicinal Plants in Drug Discovery across the World. *Indian Journal of Pure and Applied Biosciences*. 2023; 11(2): 30-41
<http://doi.org/10.187.82/2582-2845.8995>.
- 2). Miraldi, E. and Baini, G. Medicinal Plants and Health in Human History; From Empirical Use to Modern Phytotherapy. *Journal of Siena Academy of Sciences*. 2018; 10:7-12 DOI: 10.4081/jsas.2018.8529.
- 3). Rasul, M. G. Extraction, Isolation and Characterization of Natural Products from Medicinal Plants. *International Journal of of Basic Sciences and Applied Computing*. 2018; 2(6): 1-6.
- 4). Dias DA., Urban S., Roessner U. A Historical Overview of Natural products in Drug Discovery. *Metabolites* 2012; 2: 303-336. Doi: 10.3390/metabo2020303
- 5). Mahre MB, Umaru B, Ojo NA, Saidu AS, Yahi D, Ibrahim RM., Mshelbwala PP Acute Toxicity, Phytochemistry and Antidiarrheal Effects of *Celtis integrifolia* Lam. Aqueous Leaf Extract in Wistar Albino Rats. *British Journal of Pharmaceutical Research*. 2016; 14(5):1-7.
- 6). Najmi, A., Javed, SA., AL Bratty, M. and Alhazmi, HA. Modern Approaches in the discovery and Development of Plant-Based Natural Products and their Analogues as Potential Therapeutic Agents. *Molecules*, 2022; 27: 349-364.
<https://doi.org/10.3390/molecules2702049>.

- 7). Queiroz, E. F., Guillarme, D. and Wolfender, J. Advanced high Resolution Chromatographic Strategies for efficient Isolation of Natural Product from Complex Biological Matrices; From metabolite Profiling to Pure Chemical Entities. *Phytochem. Rev.* 2024; <https://doi.org/10.1007/s1101-024-09928-w>.
- 8). Hisiger, S. and Jolicoeur, M. Analysis of *Catharanthus roseus* Alkaloids by HPLC. *Phytochem. Rev.* 2007; 6: 207-234 Doi: 10.1007/s1101.006-9036-y.
- 9). Dhyani P., Quispe C., Sharma E., Bhahukhandi A., Sati P., Attri DC. et al. Anticancer Potential of Alkaloids; a key emphasis to Colchicine, Vinblastine, Vincristine, Vindesine, Vinorelbine and Vincamine. *BMC Cancer Cell International.* 2022; 22:206-225. <https://doi.org/s12935-022-02624-9>.
- 10). Wani KI, Choudhary S., Zehra A., Naeem M., Weathers P. and Aftab T. Enhancing Artemisinin content in and delivery from *Artemisia annua*: a review of alternative, classical, and transgenic approaches, *Planta.* 2021; 254: 29-43. <https://doi.org/10.1007/s00425-021-03676-3>.
- 11). Schippers, R.R. African indigenous vegetables: An overview of the cultivated species. University of Greenwich, Natural Resources Institute: London, UK. 2000; pp. 81- 83.
- 12). Mwanja, Y., Goler, E. and Gugu, F. Flowering and seed-setting studies in Livingstone potato (*Plectranthus esculentus* NE Br.) in Jos-Plateau, Nigeria. *International Journal of Plant Breeding and Genetics.* 2015; 9: 275-279.

- 13). Eleazu, C.O., Eleazu, K.C., Ironkwe, A., Iroaganachi M.A. Effect of Livingstone potato (*Plectranthus esculentus* N.E.Br) on diabetes and its complications in Streptozotocin induced diabetes in rats. *Diabetes & Metabolism Journal*. 2014; 38:366-374.
- 14). Tapera M., Gitore SA., Shumba T., Male NP., Kaseke S. Phytochemical profile of *Plectranthus esculentus* N.E.Br obtained from Zimbabwe authenticates its medicinal uses *International Journal of Pharmacognosy and Phytochemical Research*, 2019; 11(1); 27-36
- 15). Balogun, OK., Alemika T., Ojerinde O. Phytochemical, Antimicrobial and Antioxidant Screening of Leaf and Tuber Extracts of Three Varieties of *Plectranthus esculentus* Grown by Berom People of Nigeria. *International Journal of Pharmacognosy and Phytochemical Research*, 2019, 11. 27-36.
- 16). Abubakar AR. and Haque M. Preparation of Medicinal Plants: Basic Extraction and Fractionation Procedures for Experimental Purposes. *J Pharm Bioall Sci*, 2020; 1-10. Doi: 10.4103/jpbs.JPBS_175_19.
- 17). Kwaji, A., Adamu, H. M. and Chindo I. Y. Isolation, characterization and biological properties of betulin from *Entada africana* Gull. and Perr. (Mimosaceae). *Journal of Applied and Advanced Research*, 2018; 2(3): 28-31.
- 18). Teke, N. G., Lunga, P. K., Wabo, K. H., Kuate, J., Vilarem, G., Giacinti, G. et al. Antimicrobial and antioxidant properties of methanol extract fractions and compounds from the stem bark of *Entada abyssinica* Stend ex A. Satabie. *BMC Complementary and Alternative Medicine*. 2011; 11 (57): 1-8.

- 19). Njolke MA, Yafeh AV, Sechang M, Hillary J, Ndeki TP. The Antibacterial and Antifungal Activities of *Plectranthus esculentus* (Mambilla Gwe-Rizgah). *Mambilla Journal of Science and Environment*. 2024; 5(1):25-35.
- 20). Demgne OMF, Mbougna JFT, Seukep AJ, Mbaveng AT, Tene M, Nayim P. et al. Antibacterial Phytocomplexes and Compounds from *Psychotria sycophylla* (Rubiaceae) Against drug Resistant Bacteria. *Advances in Traditional Medicines*. <https://doi.org/10.1007/s13596-021-00608-0>.
- 21). Jiang, K., Gachumi, G., Poudel, A., Shurmer, B., Bashi, Z., El-Aneed, A. The establishment of tandem mass spectrometric fingerprints of phytosterols and tocopherols and the development of targeted profiling strategies in vegetable oils. *American Society for Mass Spectrometry*. 2019; 30 (1): 1700-1712.
- 22). Correa GM, Abreu VGC, Martin DAB, Takahashi JA, Fontoura HD, Cara DC, et al. Antiinflammatory and Antimicrobial activities of Steroids and Triterpenes Isolated from Aerial Parts of *Justicia acuminatissima* (Acanthaceae). *International Journal of Pharmacy and Pharmaceutical Sciences*, 2014; 6(6):75-81.
- 23). Zhang X., Wang J., Zhu L., Wang X., Meng F., Xia L., et al. Advances in Stigmasterol on its Anti-cancer effect and Mechanism of Action. *Frontiers in Oncology*, doi: 10.3389/fonc.2022.1101289 .
- 24). Backrim S., Benkhaira N., Bourais I., Benali T., Lee L-H., ElOmri N., et al. Health benefits and Pharmacological Properties of Stigmasterol. *Antioxidants* 2022; 11, 1912. <https://doi.org/10.3390/antiox11101912>.

UNDER PEER REVIEW