

A study on Extended Spectrum β -lactamase Producing (ESBL) and Fluoroquinolone-resistant *Escherichia coli* in the Japanese quails (*Coturnix japonica*) from Puducherry (India)

ABSTRACT

Antimicrobial resistance (AMR) is a growing public health threat worldwide, and members of the Enterobacteriaceae family are among the clinically important bacteria rapidly developing resistance to available antibacterial agents. The most important mechanism of resistance to third-generation cephalosporins among members of the Enterobacteriaceae is the production of extended-spectrum β -lactamase (ESBL) enzymes. Quails also represent a source of animal protein deficiency in developing countries including India. This study aimed to detect the presence of extended-spectrum β -lactamases (ESBLs) and fluoroquinolone-resistant *Escherichia coli* from fecal samples of quails in Puducherry. A total of 36 fecal samples of quails from six different retail outlets were collected. *E. coli* could be isolated from 27 fecal samples and confirmed as *E. coli* by polymerase chain reaction by targeting the *uspA* gene. The isolates were preliminarily screened for ESBL production by indicator antimicrobial disc and phenotypic ESBL production was confirmed by combination disc method. The isolates were subjected to genotypic detection of ESBL (*bla* CTX-M, *bla*TEM, and *bla* SHV) by PCR. Out of 27 isolates, 16 isolates were positive for the presence of ESBL-producing genes, in which, 9 harbored *bla*CTX-M, 2 harbored *bla*TEM, and 5 harbored *bla*SHV genes. Similarly, out of 27 isolates 13 isolates were found positive for fluoroquinolone resistance in which 13 (48.18%) isolates harbored the *qnrS* gene, and 11 (40.74%) isolates harbored the *qnrB* gene. Antimicrobial susceptibility test showed that the isolates exhibited a high level of resistance against Cefpodoxime (90%), Amoxicillin/clavulanic acid (70%), Ceftriaxone (60%), Cefotaxime (55%), Ceftazidime (45%), Ciprofloxacin (40%) and Aztreonam (30%).

Keywords: AMR, ESBL, Fluoroquinolone resistant, *E. coli*, *uspA*

1. INTRODUCTION

Over the past five decades, there has been a significant increase in poultry meat production, which now constitutes nearly one-third of global meat production (1,2). The rearing of birds other than chickens as quails has been developed in recent years worldwide to supply human food requirements (3). Quail farming is increasing globally daily as it requires low-rearing investment and has fast

financial returns. In addition, quails differ from other poultry species in that they use less feed, overgrow, reach sexual maturity early, have short generation intervals, have short incubation periods, have highly nutritious meat, and are less susceptible to viral diseases. (4,5,6). Common bacteria in the poultry gut include *Escherichia coli*, which causes various diseases such as omphalitis, cellulitis, swollen head syndrome, coli granuloma, and colibacillosis (7). The primary cause of the emergence, persistence, and dissemination of resistant bacteria is the overuse of antibiotics in intensive animal agriculture, particularly in raising pigs and poultry (8). Swine, cattle, poultry, and turkey have all been found to harbor ESBL-producing bacteria (9). The emergence of plasmid-mediated quinolone resistance was first reported in strains of *Klebsiella pneumoniae* in the United States in 1998. In the following years, several distantly related plasmids-mediated Qnr determinants were described in Enterobacteriaceae (*qnrA*, *qnrB*, *qnrC*, *qnrD*, and *qnrS*). It is becoming more common to find ESBLs and additional plasmid-mediated antibiotic-resistance genes such as *qnr* (10). Humans may acquire multidrug-resistant bacteria through zoonotic transmission from broiler farms and agricultural settings, or indirectly by food-borne diseases (11).

Puducherry, a state with intensive quail farming, lacks data regarding antibiotic use and antibiotic resistance in poultry farms. This study was therefore done to determine the antibiotic-resistant *E. coli*, study their antibiotic susceptibility pattern, and identify the resistance genes in quails in Puducherry.

2. MATERIAL AND METHODS

2.1 Collection and Processing of Samples

A total of six retail outlets were chosen for sample collection and from each retail outlet six cloacal swabs were collected using a sterile swab aseptically from healthy quails randomly in and around Puducherry. The collected 36 cloacal swabs were transported to the Department of Veterinary Microbiology Laboratory, Rajiv Gandhi Institute of Veterinary Education and Research (RIVER), Puducherry for further processing.

2.2 Isolation and Identification of *E. coli*

Each collected cloacal swab was inoculated into a Luria broth and subjected to incubation overnight at 37°C. A loopful of the culture was streaked onto MacConkey agar and incubated. The pink colonies that were lactose fermenting were subjected to Gram staining. Bergey's Manual of Systematic Bacteriology (1984) was used to identify the gram-negative bacteria as *E. coli* up to the species level based on their biochemical tests and cultural characteristics.

PCR was employed for genotypic confirmation of the *E. coli* isolates, utilizing primers targeting the *uspA* gene (12). PCR amplification was conducted using an automated thermal cycler (Eppendorf Mastercycler, Germany), following the PCR reaction conditions outlined in the protocol (12). The PCR products were analyzed by agarose gel electrophoresis with the expected product size is 884 bp for the *uspA* gene. Following PCR confirmation of *E. coli* isolates, the resulting PCR products underwent analysis through agarose gel electrophoresis. Subsequently, all confirmed *E. coli* isolates were

subjected to further detection of ESBL through both phenotypic and genotypic methods, along with genotypic detection of fluoroquinolone resistance.

2.3 Phenotypic detection of ESBL production

The preliminary phenotypic screening of ESBL production was carried out by using five indicator antimicrobial discs: ceftriaxone (30 µg), cefotaxime (30 µg), ceftazidime (30 µg), cefpodoxime (10 µg) and aztreonam (30 µg) discs. Among these five indicator discs isolates show resistance to at least one antibiotic was considered as positive for ESBL production. Further ESBL production was confirmed by following the combination disc method using cefotaxime and cefotaxime/ clavulanic acid, in the plate of Mueller-Hinton agar at a distance of 25 mm apart as per the recommendation of CLSI guidelines (2019). Plates were incubated for 16-18 hours at 37°C. The isolates were confirmed as ESBL producers if the inhibition zone diameter around the combination disc was ≥ 5 mm when compared to the disc containing respective cephalosporin alone (13).

2.4 Genotypic Detection Antimicrobial Resistance Genes

Following identification as *E. coli*, all isolates underwent PCR analysis using primers targeting *bla* TEM, *bla* SHV, and *bla* CTX-M genes for ESBL confirmation and also confirmation of fluoroquinolone-resistant genes such as *qnrS*, and *qnrB* by PCR. Template DNA preparation from *E. coli* strains followed the method described by Zhang *et al.* (14).

Table 1: Primers details used for PCR in this study and their amplicon size

Target	Primer sequence (5'-3')	Size (bp)	Annealing Temperature	References
<i>uspA</i>	CTGGAAGAGGCTAGCCTGGACGAG AAAATCGGCACCGGTGGAGCGATC	844	60°C	Chen <i>et al.</i> , 1998 (12)
<i>bla</i> TEM	ATGAGTATTCAACATTTCCG CTGACAGTTACCAATGCTTA	867	56°C	Rasheed <i>et al.</i> , 2003 (15)
<i>bla</i> SHV	AGGATTGACTGCCTTTTTG ATTTGCTGATTTGCTCG	393	56°C	Colom <i>et al.</i> , 2003 (16)
<i>bla</i> CTX-M	GAAGGTCATCAAGAAGGTGCG GCATTGCCACGCTTTTCATAG	560	59°C	Sharma <i>et al.</i> , 2013 (17)
<i>qnrS</i>	GCAAGTTCATTGAACAGGGT TCTAAACCGTCGAGTTCGGCG	428	52°C	Cattoir <i>et al.</i> , 2007 (18)
<i>qnrB</i>	GGMATHGAAATTCGCCACTG TTTGCGYGYCGCCAGTCGAA	264	50°C	

2.5 Antibiotic susceptibility testing

Antimicrobial susceptibility test was done on Mueller-Hinton agar (HiMedia, Mumbai) plate as per the recommendation of Clinical Laboratory Standard Institute (2019) using the following commercially (HiMedia, Mumbai) available antimicrobial discs (HiMedia) used in this study were

Amoxicillin/clavulanic acid (30 µg), Cefpodoxime (30 µg), Cefotaxime (30 µg), Aztreonam (30 µg), Ceftazidime (30 µg), Enrofloxacin (30 µg), and Ceftriaxone (30 µg).

3. RESULTS AND DISCUSSION

In this study, among 36 cloacal swabs examined, a total of 27 (75%) *E. coli* isolates were obtained, using cultural and biochemical characterization followed by genotypic confirmation of PCR using *uspA* gene specific for *E. coli* (844 bp) (Fig.1). The higher occurrence rate of *E. coli* is about 65%, and a lower rate of *E. coli* 3.5 % reported from quail cloacal samples (3,19).




Fig 1. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 844 bp for the *uspA* gene in *E. coli* isolates. Lane 1: 100 bp ladder; Lane 2: Negative control; Lane 3: Positive control; Lane 4,5,6,7,8: Field isolates for *uspA* gene in *E. coli*

In this study, 20 (74.07%) isolates were found preliminarily positive for ESBL production by indicator antimicrobial agents. Out of 20, phenotypically 17 (62.9%) isolates were confirmed as ESBL producers by combination disc method (Fig.2).



Fig 2. Phenotypic confirmation of ESBL production of *E. coli* isolates

In Kerala, 6/22 (27.27%) isolates were phenotypically positive for ESBL production in quails were reported (20). The genotypic method confirms the presence of genes associated with ESBL production in *E. coli* isolates. In this study, out of 27 isolates, 16 (59.25%) isolates were found positive for ESBL-producing genes in which 9 (33.33%) isolates harbored CTX-M (Fig. 3), 2 (7.4%) isolates harbored *bla* TEM (Fig. 4) and 5 (18.51%) isolates harbored *bla* SHV gene (Fig. 5).




Fig 3. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 540 bp for the CTX-M gene in *E. coli* isolates. Lane 1: 100 bp ladder; Lane 4 to 11: Field isolates for CTX-M gene in *E. coli*; Lane 2: Negative control; Lane 3: Positive control




Fig 4. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 867 bp for the TEM gene in *E. coli* isolates. Lane 1: 100 bp ladder; Lane 2&3,4,5,6: Field isolates for TEM gene in *E. coli*; Lane 7: Positive control; Lane 8: Negative control




Fig 5. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 393 bp for the SHV gene in *E. coli* isolates. Lane 1: 100 bp ladder; Lane 4,5,6,7,8,9: Field isolates for SHV gene in *E. coli*; Lane 3: Positive control; Lane 2: Negative control

Two isolates (7.4%) harbored both *bla* TEM and *bla* SHV. 20 isolates (90.9%) harbored at least one ESBL gene, 10 isolates (45.45%) harbored *bla* CTX-M gene, and 15 isolates (68.18%) harbored *bla* TEM gene in quails reported in India (20). In this study, out of 27 isolates 13 isolates were found positive for fluoroquinolone resistance in which 13 (48.18%) isolates harbored the *qnrS* gene (Fig. 6), and 11 (40.74%) isolates harbored the *qnrB* gene (Fig. 7).




Fig 6. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 428 bp for the *qnrS* gene in *E. coli* isolates. Lane 3: 100 bp ladder; Lane 1,5,6,7,8,9,10: Field isolates for *qnrS* gene in *E. coli*; Lane 2: Negative control; Lane 4: Positive control




Fig 7. Agarose gel electrophoresis showing the results of polymerase chain reaction amplified product of 264 bp for the *qnrB* gene in *E. coli* isolates. Lane 1: 100 bp ladder; Lane 2&3,4,5: Field isolates for *qnrB* gene in *E. coli*; Lane 6: Positive control; Lane 7: Negative control

No study has reported the presence of fluoroquinolone resistance genes in Japanese quails in India. In poultry farms of Southern Karnataka, a study was conducted and found results that the *qnrB* gene 38% was commonly detected followed by the *qnrS* gene 27% and *qnrA* gene 21.5% in *Escherichia coli* in collected fecal samples. From North India, the prevalence of *qnrS*, and *qnrB* were 37.50% and 5.83% respectively in from the cloacal swabs of commercial chickens reported (23). This is the first study conducted on Japanese quails of Puducherry and detected the presence of ESBL-producing and fluoroquinolone-resistant *E. coli* in fecal samples of quails in Puducherry. Antimicrobial susceptibility test showed that the majority of the isolates exhibited a high level of resistance against Amoxicillin/clavulanic acid (100%), Cefpodoxime (100%), Cefotaxime (80%), Aztreonam (70%), Ceftazidime (70%), Enrofloxacin (70%) and Ceftriaxone (40%).

4. CONCLUSION

The present study's findings indicate that Japanese quails may serve as a reservoir for ESBL and fluoroquinolone resistance *E. coli*. Similarly, the presence of ESBL-producing and fluoroquinolone-resistant *E. coli* in quails was a major concern indicating infection at the farm level. The present study showed the increased occurrence of ESBL-producing *E. coli* in quails in the Puducherry region and their resistant antibiogram pattern among quails.

References

1. Ritchie H, Roser M. Meat and dairy production (2017). Our World in Data. <https://ourworldindata.org/meat-production>
2. FAO. Meat market review, an overview of global meat market developments in 2019.
3. Khoshbakht R, Seifi S, Karimi A and Khosravi M. Molecular identification of *Campylobacter*, *Arcobacter*, and *Salmonella* in Japanese quail (*Coturnix japonica*) reared in farms of Northern Iran. *J. Food Qual. Hazards Control*. 2017;4:58-62.
4. Santos T, Murakami A, Fanhani J and Oliveira C. Production and reproduction of egg-and meat-type quails reared in different group sizes. *Revista Brasileira de Ciencia Avicola*. 2011;13: 9-14.
5. Yusuf M. S, El Nabtiti A.S. and Cui H. Effects of NENP vs LELP diets on some laying and reproductive performance parameters of Japanese quail's hens. *J. Ad. Agric. Technol* .2016;3 (2): 132-135
6. Hassan A.M, Mohammed D.A, Hussein K.N and Hussen S.H. Comparison among three lines of quail for egg quality characters. *Science Journal of University of Zakho*. 2017; 5 (4): 296-300.
7. Kabir S. M. L. Avian colibacillosis and salmonellosis: A closer look at epidemiology, pathogenesis, diagnosis, control, and public health concerns. *International Journal of Environmental Research and Public Health/International Journal of Environmental Research and Public Health*. 2010;7(1):89–114. <https://doi.org/10.3390/ijerph7010089>
8. World Health Organization Antimicrobial resistance: global report on surveillance 2014. Geneva, Switzerland.
9. Mughini-Gras L, Dorado-García A, Van Duijkeren E, Van Den Bunt G, Dierikx C. M, Bonten M. J. M et al. Attributable sources of community-acquired carriage of *Escherichia coli* containing β -lactam antibiotic resistance genes: a population-based modeling study. *The Lancet Planetary Health*. 2019; 3(8): e357–e369. doi:10.1016/S2542-5196(19)30130-5.
10. Kar D, Bandyopadhyay S, Bhattacharyya D, Samanta I, Mahanti A, Nanda P et al. Molecular and phylogenetic characterization of multidrug resistant extended spectrum beta-lactamase producing *Escherichia coli* isolated from poultry and cattle in Odisha, India. *Infection, genetics and evolution: journal of molecular epidemiology and evolutionary genetics in infectious diseases*. 2015;29:82–90. <https://doi.org/10.1016/j.meegid.2014.11.003>
11. Agyare C, Etsiapa Boamah V, Ngofi Zumbi C, Boateng Osei F. Antibiotic Use in Poultry Production and Its Effects on Bacterial Resistance. *IntechOpen*. 2019; doi: 10.5772/intechopen.79371
12. Chen J, Griffiths MW. PCR differentiation of *Escherichia coli* from other gram-negative bacteria using primers derived from the nucleotide sequences flanking the gene encoding the universal stress protein. *Lett Appl Microbiol*. 1998;27(6):369-71. doi:10.1046/j.1472-765x.1998.00445.x.

13. Clinical and Laboratory Standards institute. Performance Standards for Antimicrobial Susceptibility Testing: Twenty- Fourth Informational Supplement. M100-S24. Clinical and Laboratory Standards Institute, Wayne, PA, USA.2019.
14. Zhang QQ, Ying GG, Pan CG, Liu YS, Zhao JL. Comprehensive evaluation of antibiotics emission and fate in the river basins of China: source analysis, multimedia modeling, and linkage to bacterial resistance. *Environ Sci Technol*. 2015;49(11):6772-82. doi: 10.1021/acs.est.5b00729.
15. Rasheed JK, Jay C, Metchock B, Berkowitz F, Weigel L, Crellin J et al. Evolution of extended-spectrum beta-lactam resistance (SHV-8) in a strain of *Escherichia coli* during multiple episodes of bacteremia. *Antimicrob Agents Chemother*. 1997;41(3):647-53.doi:10.1128/AAC.41.3.647.
16. Colom K, Pérez J. J, Alonso R.A.F, Lariño E & Cisterna R. Simple and reliable multiplex PCR assay for detection of blaTEM, blaSHV and blaOXA-1 genes in Enterobacteriaceae. *Fems Microbiology Letters*. 2003;223: 147–151. doi:10.1016/S0378-1097(03)00306-9.
17. Sharma M, Pathak S, Srivastava P. (). Prevalence and antibiogram of Extended Spectrum β -Lactamase (ESBL) producing Gram negative bacilli and further molecular characterization of ESBL producing *Escherichia coli* and *Klebsiella* spp. *Journal of clinical and diagnostic research*: 2013; 7(10): 2173–2177. <https://doi.org/10.7860/JCDR/2013/6460.3462>
18. Cattoir V, Poirel L, Rotimi V, Soussy CJ, and Nordmann P. Multiplex PCR for detection of plasmid-mediated quinolone resistance qnr genes in ESBL-producing Enterobacterial isolates. *J Antimicrob Chemother*. 2007; 60:394–397.<https://doi.org/10.1093/jac/dkm204>
19. Omoshaba E. O, Olufemi F. O, Ojo O. E, Sonibare A. O. and Agbaje M. Multidrug-resistant *Salmonellae* isolated in Japanese quails reared in Abeokuta, Nigeria. *Trop. Anim. Hlth. Prod*. 2017; 49: 1455-1460. doi:10.1007/s11250-017-1347-z.
20. Andrews H. P, Asha K, Vergis P. J, & Rajasekhar R. Phenotypic and Genotypic Characterisation of ESBL Producing and Carbapenem-resistant *Escherichia coli* and *Salmonella* spp. in Japanese Quail Farms of Wayanad District. *International Journal of Current Microbiology and Applied Sciences*.2022; 11(7): 108–116. <https://doi.org/10.20546/ijcmas.2022.1107.012>
21. Singaravelan T, Antony P. X, Vasu J, Srinivas M. V, Shanmugam V. P, Shanmugam S et al. Isolation and Molecular Characterization of Extended Spectrum Beta-lactamase (ESBL) Producing *Escherichia Coli* from Dogs. *the Indian Journal of Veterinary Sciences and Biotechnology*.2022; 18(4):86–91
22. Bhardwaj K, Shenoy M S, Baliga S, B U, Baliga BS, Shetty VK. Research Note: Characterization of antibiotic-resistant phenotypes and linked genes of *Escherichia coli* and *Klebsiella pneumoniae* from healthy broiler chickens, Karnataka, India. *Poult Sci*. 2021 Jun;100(6):101094. doi: 10.1016/j.psj.2021.101094. Epub 2021 Mar 22. PMID: 33989952; PMCID: PMC8135037.
23. Rawat N, Anjali Jamwal R. S, Devi P. G, Yadav K, Kumar N, Rajagopal R. Detection of unprecedented level of antibiotic resistance and identification of antibiotic resistance factors, including QRDR mutations in *Escherichia coli* isolated from commercial chickens from North India. *Journal of Applied Microbiology*. 2022; 132: 268–278. doi:10.1111/jam.15209