

## Effectsofhexanal on shelf life of strawberry (*Fragaria x ananassa*Duch.)

### ABSTRACT

We evaluated the role of 'Enhanced Freshness Formulation' (EFF) having hexanal as the main component on shelf life enhancement of strawberry (*Fragaria x ananassa*) cv. Sweet Charlie. After crop harvest, a laboratory experiment was done at ambient temperature (about 22-30°C) with 5-treatments (dipping of freshly harvested berries in hexanal of varying strengths) viz., 1 % hexanal for 2.5 min (V<sub>1</sub>), 1 % hexanal for 5 min (V<sub>2</sub>), 2 % hexanal for 2.5 min (V<sub>3</sub>) and 2 % hexanal for 5 min (V<sub>4</sub>). Undipped berries were considered as control treatment (V<sub>5</sub>). The study revealed that dipping the berries in 2 % hexanal for 2.5 min proved most effective to prolong the shelf life of strawberry. Maximum shelf life (5.90 days) and fruit firmness (5.88 N mm<sup>-1</sup>) were obtained in the most efficacious treatment, thereby extending the marketing period.

**Keywords:** Enhanced Freshness Formulation, Hexanal, Shelflife, Storage, Strawberry

### INTRODUCTION

Allelochemicals are released from plant tissue in a variety of ways including emission of volatile substances from living plant parts, exudation from roots, or leaching from above ground parts by rain, dew, fog, etc. (30). A product called Enhanced Freshness Formulation (a hexanal formulation) has the potential to preserve quality, reduce postharvest losses and assure longevity of fresh produce without compromising on fruit quality, environmental and human health (19). Hexanal is produced in plant tissues through a lipoxygenase pathway as a naturally occurring volatile with generally-recognized-as-safe status (27,30,35). When the plants are wounded or cut, there is a grassy odour which is chemically referred as "hexanal" (10). Hexanal slows down the ripening process and treated fruits maintains the marketable quality for longer periods due to the retention by the membrane (16). Hexanal is antifungal against *Alternaria alternata*, *Botrytis cinerea* and *Penicillium expansum*. Apart from the antimicrobial activity of hexanal, it enhances the sensory quality of ripe fruit (1,11,36). Pre and postharvest applications of hexanal in several fruits, vegetables, fresh cut produce and flowers enhances their shelf-life (23,32). Synthetic fungicides used to delay spoilage, ripening and senescence of fruits are environmental concerns regarding food safety for the consumers (17). The use of new technologies (hexanal treatment) retains fruit quality, thereby reducing the postharvest losses and extending their shelf life.

Strawberries are perishable and therefore needs very careful postharvest handling primarily due to their susceptibility to mechanical damage, physiological degradation, water loss and microbial deterioration. Economic loss of fruit quality due to short shelf life begins with the loss of membrane integrity that largely plays the leading role in the senescence process (24,29). The susceptibility of strawberries to microbial decay is due to their soft skin, which is easily ruptured facilitating entry of most organisms to attack and multiply (42). In recent years advances has been made in post-harvest storage technology with the onset of Phospholipase D (PLD) inhibiting technology, the important enzyme that initiates membrane phospholipid degradation in the cell membrane during ripening (20). Phospholipase D is a key enzyme involved in membrane deterioration that occurs during

**Comment [Ma1]:** Please briefly describe methodology, statistical designed and parameters studied.

fruit ripening and senescence. The quality of highly perishable horticultural produce is reduced by excessive degradation of membranes. Hexanal, an inhibitor of phospholipase D has been successfully showing significance in enhancing storage life and fruit quality attributes by preserving membrane integrity. Hexanal, a C6 aldehyde, is a naturally occurring compound (12,44) and is responsible for downregulating the expression of PLD enzyme which degrades the phospholipid bilayer (9,21,22).

Enhancing shelf life of fruits and vegetables is a major challenge. In modern agriculture, farmers use synthetic chemicals, whose indiscriminate usage has created numerous problems such as environment and health issues. Increased awareness about these harmful effects has led to the development of alternate ecofriendly shelf life enhancement strategies. Enhanced Freshness Formulation based on hexanal enhances shelf life of many fruit crops. However, studies regarding activity of hexanal for enhancing shelf life in strawberry are rare. Therefore, this study was done to assess the effects of hexanal based Enhanced Freshness Formulation on shelf life of strawberry.



Figure 1. Strawberry population Figure 2. Strawberry plant at fruiting stage

## MATERIALS AND METHODS

Fruits were collected from strawberry plant at fruiting stage during February, 2020 and 2021 from farmer's field at Dhankhuloi, Jorhat district India. Dhankhuloi is located at a latitude  $26.77^{\circ}48'7.3476''N$  and  $94.096^{\circ}3'23.7996''E$  longitude having an altitude of 84 m above mean sea level with annual rainfall of 1879.62 mm. Maximum and minimum temperatures during February are  $25.56^{\circ}C$  and  $11.19^{\circ}C$ , respectively.

The fruits were harvested, when more than 75 % fruit attained colour from strawberry plants grown in open field under plastic mulched condition. The shelf life assessment experiment was done in completely randomized design with 4-replications (Fig 5). Strawberry fruits were subjected to hexanal based Enhanced Freshness Formulation (EFF) for shelf life treatments on the same day of harvest and stored at ambient temperature ( $22-30^{\circ}C$ ). Five treatments consisted of dipping of harvested fruits ( $V_1$ -dipped in 1 % hexanal for 2.5 min,  $V_2$ -1 % hexanal for 5 min,  $V_3$ - 2 % hexanal for 2.5 min,  $V_4$ - 2 % hexanal for 5 min) and Control ( $V_5$ - fruits without dipping). Fruit firmness ( $N\ mm^{-1}$ ) was determined by using a digital Fruit Penetrometer (Model: DFP-001, Parisa Technology).

**Comment [Ma2]:** List objectives of the study at the end of the introduction

**Comment [Ma3]:** Please give reference for rainfall and temperature data

**Comment [Ma4]:** Mention in brief about the formulation of hexanal (EFF) and its preparation clearly telling about the solvent etc. for better understanding of the reader.

The shelf life(days) was estimated by observing the number of days the fruits remained in sound condition from harvesting without spoiling to optimum eating stage at room temperature(28). Physiological loss in weight (%) of fruit was calculated as loss of weight in grams to the initial weight and expressed in percentage (13). The loss due to pathogenic infection, shrivelling and other deformity till fruits became unmarketable was considered as decay loss (%) and was determined visually during the course of the experiment (4,25). Total anthocyanins (mg 100g<sup>-1</sup>) were determined according to the pH differential method on 2<sup>nd</sup> day of harvest (45).

#### Statistical analysis

All the data were subjected to the statistical analysis of variance using completely ~~randomised~~ randomized design ~~as described by~~ (26) at 5 % level of significance with the help of a windows-based computer package OPSTAT (34) and SPSS software.

## RESULTS AND DISCUSSION

#### Fruit firmness (N mm<sup>-1</sup>)

Pooled data of both the years ~~recorded-revealed that~~ highest fruit firmness (5.88 N mm<sup>-1</sup>) ~~was recorded in V<sub>3</sub> strawberry fruits treated with 2% hexanal for 2.5 minutes and~~ minimum fruit firmness (0.44 N mm<sup>-1</sup>) was observed in ~~V<sub>5</sub> in untreated (control) fruits~~ (Table 1 and Fig 3). Fruit firmness is a major quality attribute that often dictates the shelf life of fruits. Genetic background, growing conditions and fruit constitution at the time of testing (degree of ripeness, size, post-harvest handling, internal temperature, method for fruit firmness testing etc.) affect fruit firmness. Further, as the days of storage progressed, the pulp firmness decreased, which might be due to continued ripening of fruits during storage, wherein, insoluble pectic substances of the cell wall are degraded into soluble pectin, resulting in pulp softening (8,35). Higher firmness was observed in hexanal treated samples compared to control which might be due to the delayed ripening and reduced activity of enzyme phospholipase D. Hexanal application on fruits influenced the firmness positively for a longer period of time, primarily because hexanal is a potent inhibitor of enzymes that disturb the membrane integrity. Hexanal application at optimum concentration as dip maintained the best firmness, primarily because of damaging effect at higher concentration. Furthermore, hexanal down regulates the genes responsible for enzymes involved in ethylene biosynthesis, preserves the cell membrane integrity after harvest and resulting in better pulp firmness. The activity of phospholipase D is also ~~stimulated~~ by calcium and low pH. Thus, if the action of phospholipase D is inhibited, then the rest of enzymes are unable to act on the intermediates. This would prevent the accumulation of neutral lipids and the destabilization of the membranes. Similar views were expressed by (2,3,6,9,38,41) in their studies.

#### Shelf life (days)

In the pooled analysis of data of two years showed that ~~V<sub>3</sub>~~ recorded highest shelf life (5.90 days) and shortest shelf life (2.37 days) in ~~V<sub>5</sub>~~ (Table 1 and Fig 4). Maximum shelf life was observed in fruits treated with hexanal of optimum concentration and duration (2% for 2.5 min). The fruits treated with hexanal showed a natural delayed ripening process. Delayed softening of fruits extends shelf life of fresh produces. High concentrations and short duration of exposure resulted in more change than lower concentration treatment for

**Comment [Ma5]:** It should be self-explanatory. Instead of mentioning V1, V2 etc. clearly mention the treatment for better understanding of the reader.

**Comment [Ma6]:** Described as suggested

**Comment [Ma7]:** Check and confirm whether it is stimulated or inhibited

**Comment [Ma8]:** Describe it as suggested in previous parameter.

long time exposure time. This might be due to fact that lower concentration were not enough to inhibit action of PLD enzyme. A simple inhibition of membrane deterioration by hexanal may enhance the shelf life by preserving membrane structure. Hexanal enhances the shelf life of fruits by maintaining fruit firmness of the treated fruits. Similar results were reported for apple, cherry, peach, plum, guava and vegetables by (2,5,33,39,41).

Table 1. Effects of hexanal treatment on fruit firmness (N mm<sup>-1</sup>), shelf life (days) and anthocyanin content (mg 100g<sup>-1</sup>)

Treatment	Fruit firmness (N mm <sup>-1</sup> )			Shelf life (days)			Anthocyanin content (mg 100g <sup>-1</sup> ) on 2 <sup>nd</sup> DAH		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
V <sub>1</sub> (2.5 min 1% hexanal)	5.05 <sup>c</sup>	5.09 <sup>c</sup>	5.07 <sup>c</sup>	4.90 <sup>b</sup>	4.95 <sup>b</sup>	4.92 <sup>b</sup>	51.11 <sup>c</sup>	50.88 <sup>c</sup>	51.00 <sup>c</sup>
V <sub>2</sub> (5.0 min 1% hexanal)	5.30 <sup>b</sup>	5.34 <sup>b</sup>	5.32 <sup>b</sup>	5.00 <sup>b</sup>	5.05 <sup>b</sup>	5.02 <sup>b</sup>	57.42 <sup>b</sup>	56.77 <sup>b</sup>	57.10 <sup>b</sup>
V <sub>3</sub> (2.5 min 2% hexanal)	5.85 <sup>a</sup>	5.91 <sup>a</sup>	<b>5.88<sup>a</sup></b>	5.80 <sup>a</sup>	6.00 <sup>a</sup>	<b>5.90<sup>a</sup></b>	63.13 <sup>a</sup>	62.85 <sup>a</sup>	<b>62.99<sup>a</sup></b>
V <sub>4</sub> (2.5 min 2% hexanal)	5.36 <sup>b</sup>	5.42 <sup>b</sup>	5.39 <sup>b</sup>	5.10 <sup>b</sup>	5.15 <sup>b</sup>	5.12 <sup>b</sup>	47.40 <sup>d</sup>	45.34 <sup>d</sup>	46.37 <sup>d</sup>
V <sub>5</sub> (Control, Fruits not dipped)	0.42 <sup>d</sup>	0.46 <sup>d</sup>	0.44 <sup>d</sup>	2.25 <sup>e</sup>	2.50 <sup>e</sup>	2.37 <sup>e</sup>	45.75 <sup>d</sup>	44.15 <sup>d</sup>	44.95 <sup>e</sup>
SEd(±)	0.10	0.11	0.07	0.21	0.19	0.14	0.82	0.62	0.52
CD(P=0.05)	0.21	0.23	0.15	0.44	0.40	0.29	1.74	1.32	1.05

#### Physiological loss in weight (%)

The physiological loss in weight increased with the increasing period of storage after harvest irrespective of the treatments. Data regarding the physiological loss in weight (%) as influenced by the treatments are presented in Table 2. In the combined analysis of data of both years, on 6th day of harvest, the fruits under V<sub>3</sub> exhibited minimum physiological loss in weight (27.86 %) whereas maximum physiological loss in weight (44.89 %) was recorded under V<sub>5</sub>. Hexanal treated strawberry fruits showed minimum physiological loss in weight than control. Physiological loss in weight of stored produce occurs due to direct water losses vis-à-vis higher respiration rate. Strawberries are subjected to rapid water loss causing them to beshrivelled and deteriorated (13). The reduced weight losses due to hexanal application might be ascribed to phospholipase D inhibition and reduction in phospholipids degradation as well as downstream oxidative processes and preserved membrane integrity. Post harvest application of hexanal might have effectively maintained the cell structures which lead to reduction of respiration rate and turgor loss that contributed to reduced weight loss in treated fruits. Hexanal treatments could have reduced the catabolic processes and quality losses by maintaining the membrane integrity and cell structure. Similar effects of hexanal application on physiological weight loss were observed by (6,7,9,15,23,41).

Table2. Effects of hexanal treatment on physiological loss in weight (%)

Treatments	Days after harvest								
	2 <sup>nd</sup> day			4 <sup>th</sup> day			6 <sup>th</sup> day		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
V <sub>1</sub> (2.5 min 1% hexanal)	5.76 <sup>b</sup>	6.42 <sup>b</sup>	6.09 <sup>bc</sup>	16.70 <sup>b</sup>	16.19 <sup>b</sup>	16.45 <sup>b</sup>	36.25 <sup>b</sup>	37.11 <sup>b</sup>	36.68 <sup>b</sup>
V <sub>2</sub> (5.0 min 1% hexanal)	5.87 <sup>b</sup>	7.30 <sup>b</sup>	6.58 <sup>b</sup>	16.30 <sup>b</sup>	15.75 <sup>b</sup>	16.03 <sup>b</sup>	36.36 <sup>b</sup>	37.45 <sup>b</sup>	36.91 <sup>b</sup>
V <sub>3</sub> (2.5 min 2% hexanal)	4.15 <sup>b</sup>	5.29 <sup>b</sup>	4.72 <sup>c</sup>	13.45 <sup>c</sup>	14.98 <sup>b</sup>	14.22 <sup>c</sup>	26.96 <sup>c</sup>	28.76 <sup>c</sup>	<b>27.86<sup>c</sup></b>
V <sub>4</sub> (2.5 min 2% hexanal)	5.32 <sup>b</sup>	6.08 <sup>b</sup>	5.70 <sup>bc</sup>	14.67 <sup>bc</sup>	16.57 <sup>b</sup>	15.62 <sup>bc</sup>	34.15 <sup>b</sup>	34.97 <sup>b</sup>	34.56 <sup>b</sup>
V <sub>5</sub> (Control, Fruits not dipped)	9.25 <sup>a</sup>	9.81 <sup>a</sup>	9.53 <sup>a</sup>	20.88 <sup>a</sup>	20.12 <sup>a</sup>	20.50 <sup>a</sup>	43.84 <sup>a</sup>	45.94 <sup>a</sup>	44.89 <sup>a</sup>
SEd(±)	0.79	1.06	0.66	1.19	0.75	0.70	2.12	1.56	1.31
CD(P=0.05)	1.68	2.26	1.31	2.53	1.60	1.43	4.52	3.33	2.69

#### Decay loss (%)

The decay loss increased in all the treatments with increasing day of storage (days) after harvest (Table 3). The combined analysis of the data of both the years, on 6<sup>th</sup> day of harvest showed that, the fruits under V<sub>3</sub> exhibited minimum decay loss (59.03 %) whereas maximum decay loss (74.73 %) was recorded under V<sub>5</sub>. Hexanal is highly volatile and possess antifungal properties against *Alternaria alternata*, *Botrytis cinerea* and *Penicillium expansum*. Infection by pathogens may occur during the growing season, at harvest time or during handling and transport. Inoculum load carried by the fruits from growing field and sanitation during storage handling plays an important role to minimize and maximize contamination by decay pathogens. Moreover, fruit composition and fruit maturity status could also affect the decay severity. Higher decay loss could be due to higher presence of disease causing inoculums due to favourable/ favorable environmental conditions. The ability to control previously established infections in the postharvest environment is of crucial importance. Hexanal has the ability to penetrate tissues to control latent or established infections. Hexanal along with other constituents of EFF (Enhanced Freshness Formulation) such as antioxidants may help maintain the membrane structure, as well as provide protection from the oxidative damage. In our study, strawberry fruits treated with hexanal resulted in a significant decrease in fungal decay upto some extent confirming the previous reports about its antifungal effect. Fruits without dip in the hexanal based EFF and stored in the container developed latent fungal growth which resulted in the fruit decay. However dipping in hexanal based EFF for longer duration might have increased the chance of fungal infection due to the increase in surface moisture that favoured fungal infection. The lower concentration and longer exposure times of hexanal treatment might have resulted in less effect on percentage decay. The results of the study have conformity with previous studies of (9,14,37,40,43,46).

**Comment [Ma9]:** Please mentioned about hexanal formulation with EFF in Materials and methods section for better understanding of your treatments.

Table3. Effects of hexanal treatment on decay loss (%)

Treatment	Days after harvest								
	2 <sup>nd</sup> day			4 <sup>th</sup> day			6 <sup>th</sup> day		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
V <sub>1</sub> (2.5 min 1% hexanal)	15.47 <sup>bc</sup>	18.49 <sup>b</sup>	16.98 <sup>bc</sup>	32.14 <sup>b</sup>	35.41 <sup>b</sup>	33.78 <sup>b</sup>	65.62 <sup>b</sup>	64.58 <sup>b</sup>	65.10 <sup>b</sup>
V <sub>2</sub> (5.0 min 1% hexanal)	14.87 <sup>bc</sup>	17.79 <sup>b</sup>	16.33 <sup>bc</sup>	30.62 <sup>b</sup>	34.37 <sup>b</sup>	32.50 <sup>b</sup>	63.33 <sup>b</sup>	62.91 <sup>b</sup>	63.12 <sup>b</sup>
V <sub>3</sub> (2.5 min 2% hexanal)	13.98 <sup>c</sup>	16.78 <sup>b</sup>	15.38 <sup>c</sup>	28.24 <sup>b</sup>	29.11 <sup>c</sup>	28.68 <sup>c</sup>	59.16 <sup>b</sup>	58.89 <sup>c</sup>	<b>59.03<sup>c</sup></b>
V <sub>4</sub> (2.5 min 2% hexanal)	17.49 <sup>b</sup>	18.80 <sup>b</sup>	18.15 <sup>b</sup>	30.95 <sup>b</sup>	35.41 <sup>b</sup>	33.18 <sup>b</sup>	63.54 <sup>b</sup>	62.29 <sup>bc</sup>	62.91 <sup>b</sup>
V <sub>5</sub> (Control, Fruits not dipped)	23.75 <sup>a</sup>	25.17 <sup>a</sup>	24.46 <sup>a</sup>	46.61 <sup>a</sup>	51.39 <sup>a</sup>	50.00 <sup>a</sup>	75.35 <sup>a</sup>	74.10 <sup>a</sup>	74.73 <sup>a</sup>
SEd(±)	1.27	2.06	1.21	1.77	1.78	1.25	2.88	1.77	1.69
CD(P=0.05)	2.71	4.39	2.47	3.78	3.81	2.57	6.14	3.78	3.45

Superscript by same letter means they are at par

#### Anthocyanin content (mg/100g)

The pooled analysis of data of two years (Table 1) showed that highest amount of anthocyanin (62.99 mg 100g<sup>-1</sup>) was recorded in V<sub>3</sub> and minimum (44.95 mg 100g<sup>-1</sup>) in V<sub>5</sub>. Hexanal application significantly reduced the degradation of anthocyanin which might be ascribed to inhibited activities of cell wall degrading phospholipase D enzyme responsible for senescence of tissue. Hexanal formulation also contained cinnamic acid and it might have induced the synthesis of anthocyanin. Supplemented anthocyanin accumulation in hexanal treated fruits might also be ascribed to increased biosynthesis of volatile and antioxidant compounds which contributed to enhanced anthocyanin synthesis. However, there was difference in results between the concentrations of hexanal and the dipping duration of dip in the experiment. Effectiveness of hexanal was significantly affected by duration of application at optimum dose level as effective concentration of hexanal varies with commodity and variety. The results are in agreement with findings of (15,18,24) and in contrast to the findings of (33) in cherry where no change in anthocyanin content was noticed in stored fruits treated with hexanal containing formulations.

#### CONCLUSIONS

The application of hexanal dip at appropriate dose established minimum decay loss, physiological loss in weight and maximum level of fruit firmness, shelf life and anthocyanin content. The results reflected maximum inhibition percentage due to hexanal treatment which showed that hexanal had a strong inhibitory effect on microbial growth in strawberry thereby enhanced its shelf life. Shorter duration of exposure of 2.5 min allowed easy and sufficient penetration of treatment solution in strawberry compared to 5 min resulting to its less effectiveness which could be due to its thin skin. During postharvest handling, strawberry fruit have high susceptibility to mechanical damage, and thus they need to be handled with care, to avoid damage that can create an entry point for the decay causing fungi. For this reason, strawberry fruit are harvested and often packaged in the

field. Dipping of freshly harvested fruits in 2 % hexanal for 2.5 min proved a better option in terms of effective concentration and effective treatment duration respectively for maintaining overall strawberry ~~fruit~~ quality during storage.

## REFERENCES

1. Archbold, D.D., Hamilton-Kemp, T.R. and Fallik, E. (2000). Aroma volatiles as modulators of postharvest mold development on fruit: *In vivo* role and fumigation tools. *Acta Horticulturae* **518**: 87-92.
2. Ashitha, G.N., Preetha, P. and Varadharaju, N. (2019). Effects of post harvest hexanal treatment on shelf life of guava. *International Journal of Current Microbiology and Applied Science* **8**(7): 1057-1067.
3. Ashitha, G.N., Sunny, A.C. and Nisha, R. (2020). Effect of pre-harvest and post-harvest hexanal treatments on fruits and vegetables: A Review. *Agricultural Reviews* **41**(2): 124-131.
4. Ayala-Zavala, J.F., Wang, S.Y., Wang, C.Y. and González-Aguilar, G.A. (2004). Effects of storage temperatures on antioxidant capacity and aroma compounds in strawberry fruit. *Lebensm.-Wiss. u.-Technologie* **37**: 687-695.
5. Baltazari, A., Mtui, H., Chove, L., Msogoya, T., Kudra, A., Tryphone, G., Samwel, J., Paliyath, G., Sullivan, A., Jayasankar, S. and Mwatawala, M. (2020). Evaluation of post-harvest losses and shelf life of fresh mango (*Mangifera indica* L.) in Eastern Zone of Tanzania. *International Journal of Fruit Science* **20**(4): 855-870.
6. Cheema, A., Padmanabhan, P., Subramanian, J., Blom, T. and Paliyath, G. (2014). Improving quality of greenhouse tomato (*Solanum lycopersicum* L.) by pre- and postharvest applications of hexanal - containing formulations. *Postharvest Biology and Technology* **95**: 13-19.
7. Cheema, A., Padmanabhan, P., Amerc, A., Parry, M.J., Limb, L.T., Subramanian, J. and Paliyath, G. (2018). Postharvest hexanal vapour treatment delays ripening and enhances shelf life of greenhouse grown sweet bell pepper (*Capsicum annuum* L.). *Postharvest Biology and Technology* **136**: 80-89.
8. Doving, A., Mage, F. and Vestheim, S. (2005). Methods for testing strawberry fruit firmness. *Small Fruits Review* **4**(2): 11-34.
9. Gill, K.S., Dhaliwal, H.S., Mahajan, B.V.C., Paliyath, G. and Boora, R.S. (2015). Enhancing postharvest shelf life and quality of guava (*Psidium guajava* L.) cv. Allahabad Safeda by pre harvest application of hexanal containing aqueous formulation. *Postharvest Biology and Technology* **112**: 224-232.
10. Gunasekaran, K., Karthika, S., Nandakumar, N.B., Subramanian, K.S., Paliyath, G. and Subramanian, J. (2015). *Biosafety of Hexanal*. Department of Nano Science & Technology Directorate of Natural Resource Management Tamil Nadu Agricultural University, Coimbatore, India & University of Guelph, Canada pp.1-92.
11. Hamilton-Kemp, T.R., McCracken, C.T., Loughrin, J.H., Andersen, R.A. and Hildebrand, D.F. (1992). Effects of some natural volatile compounds in the pathogenic fungi *Alternaria alternata* and *Botrytis cinerea*. *Journal of Chemical Ecology* **18**: 1083-1091.
12. Hildebrand, D.F. (1989). Lipoxygenases. *Physiologia Plantarum* **76**: 249-253.
13. Kannaujia, P.K. (2012). *Influence of Sequential Harvesting on Post Harvest Quality of Strawberry (Fragaria x ananassa Duch.) fruits*. M.Sc (Post Harvest Technology) Thesis, IARI, New Delhi.
14. Kannaujia, P.K. and Asrey, R. (2021). Effects of harvesting season and cultivars on storage behaviour, nutritional quality and consumer acceptability of strawberry (*Fragaria x ananassa Duch.*) fruits. *Acta Physiologia Plantarum* **43**(6): 88.
15. Kaur, S. (2018). *Studies on Quality Improvement and Storage Life in Flame Seedless Grapes (Vitis vinifera L.) Through Use of Pre and Post-Harvest Application of Chemicals*. Ph.D Thesis submitted to Punjab Agricultural University, Ludhiana.
16. Kumar, S.K. (2018). *Enhancement of Post-Harvest Shelf Life of Nectarines (Prunus persica (L.) Batsch var. nectarina) using Hexanal*. M.Sc. Thesis University of Guelph.
17. Mari, M., Martini, C., Spadoni, A., Rouissi, W. and Bertolini, P. (2012). Bio control of apple post-harvest decay by *Aureobasidium pullulans*. *Postharvest Biology and Technology* **73**: 56-62.
18. Misran, A., Padmanabhan, P., Sullivan, J.A., Khanizadeh, S. and Paliyath, G. (2015). Composition of phenolics and volatiles in strawberry cultivars and influence of preharvest hexanal treatment on their profiles. *Canadian Journal of Plant Science* **95**(1): 115-126.
19. Mthembu, S. (2020). *Evaluating the Efficacy of Formulations Containing Hexanal, Moringa Leaf Extracts, and Carboxy Methylcellulose as Postharvest Treatments for Fresh Tomatoes (Solanum lycopersicum L.)*. MSc, University of KwaZulu-Natal, Pietermaritzburg.

20. Paliyath, G. and Droillard, M.J. (1992). The mechanisms of membrane deterioration and disassembly during senescence. *Plant Physiology and Biochemistry* **30**: 789-812.
21. Paliyath, G., Yada, R.Y., Murr, D.P. and Pinhero, R.G. (2003). *Inhibition of Phospholipase D*. US patent no 6514914.
22. Paliyath, G. and Murr, D.P. (2007). *Compositions for the Preservation of Fruit and Vegetables*. US patent 7, 198, 811.
23. Paliyath, G. and Subramanian, J. (2008). Phospholipase D inhibition technology for enhancing shelf life and quality. In: *Postharvest Biology and Technology of Fruits, Vegetables, and Flowers* (Eds., G. Paliyath, D.P.Murr, A.K. Handa and S.Lurie), pp. 240-245. 1st edn. Wiley-Blackwell, Hoboken, N.J.
24. Paliyath, G., Tiwari, K., Yuan, H. and Whitaker, B.D. (2008). Structural deterioration in produce: phospholipase D, membrane deterioration, and senescence. In: *Postharvest Biology and Technology of Fruits, Vegetables, and Flowers* (Eds., G. Paliyath, D.P.Murr, A.K. Handa and S.Lurie), pp. 195-239. 1st edn. Wiley-Blackwell, Hoboken, N.J.
25. Panda, A.K. (2015). *Studies on Modified Atmospheric Packaging of strawberry (Fragaria x ananassa Duch.)*. M.Sc. (Hort.) Thesis submitted to CCS Haryana Agricultural University, Hisar.
26. Panse, V.G. and Sukhatme, P.V. (1985). *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research, New Delhi.
27. Preethi, P., Soorianathasundaram, K., Sadasakthi, A., Subramanian, K.S., Vijay Rakesh Reddy, S., Paliyath, G. and Subramanian, J. (2021). Pre-harvest application of hexanal as a surface treatment improved storage life and quality of mango fruits. *Coatings* **11**, 1267: 1-18.
28. Rahman, M.M., Moniruzzaman, M., Ahmad, M.R., Sarker, B.C. and Alam, M.K. (2014). Maturity stages affect the postharvest quality and shelf-life of strawberry genotypes growing in subtropical regions. *Journal of the Saudi Society of Agricultural Sciences* **124**(3): 1-10.
29. Rashid, M.H.A., Suravi, J.H. and Ashrafun, N. (2020). Effects of vertical staking and different types of manures on growth, yield and quality of strawberry (*Fragaria x ananassa Duch.*). *Journal of Bangladesh Agricultural University* **18**(2): 307-315.
30. Rice, E.L. (1984). *Allelopathy*. 2nd edition. NY: Academic Press.
31. Sáenz-garza N.E., Delaquis P., Durance, T. (2013). Microencapsulation of hexanal by radiant energy vacuum microwave-molecular inclusion for controlled release and inhibition of *Penicillium expansum* in a model system and on apple tissue. *Food Research International* **52**: 496-502.
32. Sekar, C., Subramanian, K.S., Janavi, G.J. and Subramanian, J. (2018). Economic Impact of Hexanal based Nanotechnology on Mango Value Chain in Tamil Nadu State, India. In: *Postharvest Biology and Nanotechnology* (Eds., G. Paliyath, J. Subramanian, L. Lim, K.S. Subramanian, A.K. Handa, and A.K. Mattoo), Pp. 327-364. 1<sup>st</sup> Edition. John Wiley & Sons, Inc.
33. Sharma, M., Jacob, J.K., Subramanian, J. and Paliyath, G. (2010). Hexanal and 1-MCP treatments for enhancing the shelf life and quality of sweet cherry (*Prunus avium* L.). *Scientia Horticulturae* **125**: 239-247.
34. Sheoran, O.P., Tonk, D.S., Kaushik, L.S., Hasija, R.C. and Pannu, R.S. (1998). Statistical software package for agricultural research workers. In: *Recent Advances in information theory, Statistics & Computer Applications*, (Eds., D.S. Hooda and Hasija, R.C.). Department of Mathematics Statistics, CCS HAU, Hisar, pp. 139-143.
35. Solomos, T. and Laties, G. G. (1973). Cellular organization and fruit ripening. *Nature* **245**: 390-392.
36. Song, J., Leepipattanawit, R., Deng, W. and Beaudry, R.M. (1998). Hexanal acts as residue less antifungal agent that enhances aroma biosynthesis in apple fruit. *Acta Horticulturae* **464**: 219-224.
37. Song, J., Hilderbrand, P.D., Fan, L., Fomey, C.F., Renderos, W.E., Campbell-Palmer, L. and Doucette, C. (2007). Effect of hexanal vapor on the growth of postharvest pathogens and fruit decay. *Journal of Food Science* **72**(4): 108-111.
38. Sulaimankhil, Z., Sethi, S., Sharma, R.R., Verma, M.K. and Bhowmik, A. (2021). Influence of hexanal concentration and exposure time on quality of cold stored apples (*Malus domestica*). *Indian Journal of Agricultural Sciences* **91**(5): 57-61.
39. Thakur, J.P., Gothwal, P.P. and Singh, I. (2017). Post-harvest treatments for extension of mango fruit var. Dashehari (*Mangifera indica* L.). *International Journal of Food Science and Nutrition* **2**(4): 156-162.
40. Thavong, P., Archbold, D.D., Pankasemsuk, T. and Koslanund, R. (2010). Effects of hexanal vapour on Longan fruit decay, quality and phenolic metabolism during cold storage. *International Journal of Food Science & Technology* **45**(11): 2313-2320.
41. Tiwari, K. and Paliyath, G. (2011). Microarray analysis of ripening regulated gene expression and its modulation by 1-MCP and hexanal. *Plant Physiology and Biochemistry* **49**(3): 329-340.

42. Tournas, V.H. and Katsoudas, E. (2005). Mould and yeast flora in fresh berries, grapes and citrus fruits. *International Journal of Food Microbiology***23**: 684-688.
43. Utto, W., Mawson, A.J. and Bronlund, J.E. (2008). Hexanal reduces infection of tomatoes by *Botrytis cinerea* whilst maintaining quality. *Post Harvest Biology and Technology***47**: 434-437.
44. Waddell, W.J., Cohen, S.M., Feron, V.J., Goodman, J.I. and Marnett, L.J. (2010). GRAS Flavoring Substances. *Food Technology***14**: 488-495.
45. Yang, L., Rong-Rong, C., Ji-Li, F. and Ke, Y. (2019). Total anthocyanins and Cyanidin-3-O-Glucoside contents and antioxidant activities of purified extracts from eighteen different pigmented plants. *Pharmacognosy Magazine***15**(60): 124-129.
46. Yuan, H.Y., Hu, X.L., Li, X., Li, Y.H. and Gopinadhan, P. (2009). Effect of hexanal treatment on postharvest quality of 'Darselect' strawberry (*Fragaria x ananassa* Duch.) fruit. *Acta Horticulturae***839**: 315-321.

UNDER PEER REVIEW




Figure 3. Effects of Hexanal treatment on Fruit firmness (N mm<sup>-1</sup>)




Figure 4. EffectsofHexanaltreatment onShelf life (days)



Figure 5. Post harvest dip of Strawberry in Hexanal