

## Original Research Article

### MINOR MODIFICATION IN ASBESTOS ROOF OF COW SHED FOR REDUCTION OF HEAT STRESS IN DAIRY COWS DURING SUMMER SEASON

#### Abstract

Thermo neutral zone (TNZ) is the range of ambient temperature & humidity in which animals feel comfortable and their production level is optimum. When temperature humidity index (THI) goes above 72, the dairy cows begin to show symptoms of mild heat stress, which affects the milk production in dairy animals during summer season in India. Asbestos and GI sheet are commonly used as roofing material in many cattle farms throughout the world. This experiment was carried out on apparently healthy cross bred jersey cows receiving balanced ration during the month of May in the livestock farm of OUAT, Bhubaneswar for one week. The existing cattle shed was partitioned into three compartments by raising brick walls, each having the dimension of 30 x 25 ft size. The asbestos roof of first one was covered with 90 % grade green shed-net. Over the asbestos roof of second compartment, heat reflective white colour was painted and asbestos roof of third one was left as such. From 8.00 AM to 5.00PM, the THI of the normal asbestos roof shed varied from mild to moderate heat stress range. But THI of both shed-net covered shed and heat reflective colour painted shed was found to be within the mild heat stress range i.e. 72-79 during noon indicating a comfortable ambient condition for the animals in summer season when the outside ambient temperature went up to 40 °C during noon. No significant ( $P>0.05$ ) change in the milk production was observed among the groups during the week. However, a decrease ( $P<0.05$ ) of 0.43, 0.25 and 0.38 L in milk yield of the cows kept under normal asbestos shed, heat reflective colour painted shed and shed-net covered shed, respectively. From the study, this can be concluded that the impact of heat stress on milk production can be reduced by painting the rood of the shed with heat reflective colour paint.

**Key words:** Heat stress, Heat reflective paint, Shed-net, Housing, THI, Milk yield.

#### 1. Introduction:

Thermo neutral zone is the range of ambient temperature & humidity in which animals feel comfortable. In this zone the health and performance of animals are found to be better. When temperature humidity index (THI) goes above 72, the dairy cows begin to show symptoms of mild heat stress [1,2]. Armstrong [3] considered THI <72 as a thermal comfort zone for dairy cows (assuming the THI does not drop below the thermo-neutral zone) and 72 to 79 as mild heat stress, 80 to 90 as moderate heat stress and >90 as severe heat stress. Heat stress is one of the leading factor

affecting milk production in dairy animals during summer season in India [4,5,6]. Gantner *et al.* [7] observed a highly significant ( $P<0.01$ ) decrease of milk yield, fat and protein content due to heat stress condition. Asbestos and GI sheet are commonly used as roof material in many cattle farms throughout the world [8,9]. But during hot summer, these roofs being good conductor of heat, contribute significantly to heat stress in animals. Roof material plays a key role in determination of thermal heat exchange in animals [10]. Reports also suggests that, Agro-net shading of roof is an efficient way to reduce heat stress [11,12,19,20]. This experiment was conducted to study the effect of modification of asbestos roof of cow shed by painting with heat reflective colour and covering with shed-net to mitigate heat stress in summer season.

## 2. Materials and Methods:

This experiment was carried out on apparently healthy cross bred milking jersey cows receiving balanced ration during the month of May in the livestock farm of OUAT, Bhubaneswar for one week. The cows were fed with a concentrate mixture (Table 1) and hybrid napier grass based ration. The nutrient content of the concentrate mixture is depicted in Table 2.

Table 1. Composition of the concentrate mixture fed to the cattle during experiment

SI No	Ingredient	Quantity (Kg)
1	Maize	36.00
2	Soybean Meal (SBM)	20.00
3	Wheat Bran (WB)	23.00
4	Rice Bran (RB)	18.00
5	Mineral Mix	2.00
6	Salt*	1.00
	Total	100.00
*Commercial mineral mixture available in the market.		

Table 2. Nutrient content of the concentrate mixture fed to the cattle during experiment

SI No	Ingredient	Quantity (%)
1	Dry matter (DM)	86.76
2	Organic matter (OM)	90.77
3	Crude Protein (CP)	17.06
4	Ether Extract (EE)	2.45
5	Total ash (TA)	8.44
6	NDF	35.06
7	ADF	16.16
8	ADL	5.28

The existing cow shed was having asbestos roof, brick wall, concrete floor and other recommended specifications like floor space, air space, ventilation etc. It was partitioned into three compartments by raising brick walls, each having the dimension of 30 x 25 ft size. The asbestos roof of first one was covered with 90 % grade green shed-net. Over the asbestos roof of second compartment, heat reflective white colour was painted and asbestos roof of third one was left as such (Fig.1). The ambient temperature and relative humidity of all three sheds and outside the shed were recorded on hourly basis from 8.00 AM to 5.00 PM by help of digital thermometer and hygrometers and the temperature humidity index (THI) was calculated as suggested by Habeeb *et al.* [1].



**Fig. 1. Outside view of the experimental shed**

Twelve nos. of medium yielder milking cross bred cows were randomly divided to three groups and kept in these three sheds. They were maintained as per the recommended feeding and management practices. The milk yield of each cow was recorded on daily basis during the experiment. The data obtained in the study was analyzed with one way ANOVA [13].

### **3. Results and Discussion:**

During the observation from 8.00 AM to 5.00 PM, the outside temperature ranged from 31.5 to 40.1 °C. The inside temperature of the normal asbestos shed, during the same period ranged from 30.5 – 38.4 °C (Table 3&4). The temperature humidity index (THI) was found to be within mild heat stress range (72 – 79) from 8.00 AM to 12.00 PM and moderate heat stress range (80 – 83) from 12.00 PM to 3.00 PM (Fig. 2). Vanitha and Baskaran [14] reported similar finding of higher temperature range in asbestos roof shed in summer season. As per Roy and Chatterjee [15], macro and micro environment of the shed has a lot of effect on physiological stress or comfort of the animals. The inside temperature of the shed-net covered shed from 8.00 AM to 5.00 PM ranged from 30.0 – 36.3 °C with THI varying from 72 - 79 indicating mild heat stress throughout the day. Whereas, the inside temperature of the heat reflective colour painted shed during the same period ranged from 28.5 – 33.5 °C with THI value of 70 – 76 indicating mild heat stress was prevailed from 10.00 AM to 4.00 PM. It

was observed that the inside temperature of the heat reflective colour painted shed was 3 - 5 °C less than the temperature of the shed-net covered shed and normal asbestos shed during noon in summer season indicating a better comfortable shed environment.

Table 3. Mean ambient conditions of normal asbestos roof shed, shed-net covered shed and heat reflective colour painted shed in summer

Time	Normal Asbestos shed			Shed net covered shed			heat reflective colour painted shed		
	Amb. Temp (°C)	RH (%)	THI	Amb. Temp (°C)	RH (%)	THI	Amb. Temp (°C)	RH (%)	THI
8.00 AM	30.5	85	72	30.0	85	72	28.5	85	70
9.00 AM	32.0	78	74	31.0	80	73	29.0	81	71
10.00 AM	33.0	74	76	32.0	78	74	29.5	79	72
11.00 AM	34.5	66	78	33.0	75	75	30.5	77	73
12.00 PM	36.0	60	80	34.2	71	77	31.5	75	74
1.00 PM	37.5	52	82	35.4	70	78	32.7	72	75
2.00 PM	38.4	46	83	36.3	66	79	33.5	69	76
3.00 PM	37.2	50	82	35.0	62	78	32.0	65	75
4.00 PM	35.0	60	79	33.5	68	76	30.5	69	73
5.00 PM	32.6	72	75	31.0	74	73	29.0	73	71

Table 4. Daily maximum outside and inside temp (°C) of normal asbestos-roof shed, shed-net covered shed and heat reflective colour painted shed in summer

Shed types	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Mean
Outside shed	39.9	40	39.9	40.1	40	39.9	39.9	40.0
Normal asbestos roof shed	38.4	38.5	38.4	38.5	38.4	38.4	38.5	38.4
Shed-net covered shed	36.2	36.3	36.2	36.3	36.3	36.3	36.2	36.3
Heat reflective colour painted shed	33.6	33.5	33.5	33.5	33.5	33.5	33.6	33.5

The daily average milk yield of cows at the beginning of the experiment kept under normal asbestos shed was 5.73 L, which was decreased by 0.45 L over one week of observation during the experiment. West [16] observed a reduction of 0.2 kg in milk yield per unit increase in THI beyond 72 in high yielding dairy cows. Similarly, the daily average milk yield was decreased by 0.38 and 0.25 L in case of cows kept under shed-net covered shed and heat reflective painted shed, respectively during the same period (Table. 6, Fig. 3).

UNDER PEER REVIEW


Table 5. Daily milk production (L) by the cows under different type of modified roofing during peak summer

Shed types	Milk Yield (L)							
	Day-1	Day-2	Day-3	Day-4	Day-5	Day-6	Day-7	Change in milk yield
Normal Asbestos shed	5.70±0.51	5.625±0.53	5.55±0.52	5.45±0.49	5.40±0.51	5.37±0.48	5.27±0.48	-0.43±0.04
Shed net covered shed	5.75±0.56	5.67±0.57	5.55±0.56	5.52±0.55	5.42±0.57	5.40±0.60	5.37±0.55	-0.38±0.04
Heat reflective colour painted shed	5.50±0.32	5.42±0.36	5.40±0.35	5.37±0.33	5.30±0.35	5.27±0.32	5.22±0.29	-0.25±0.05
Values are average of 4 reading.								
Values represents Mean± SD								
Negative values of change in milk yield represents drop in milk production on 7 <sup>th</sup> day as compared to the 1 <sup>st</sup> day								


Table 6. Effect of different roofing materials on Milk yield (L) of dairy cows during peak summer

Shed types	Milk yield (start of Study) in L	Milk yield (End of Study) in L	Change in milk yield* in L
Normal Asbestos shed	5.70±0.51	5.275±0.48	-0.43 ± 0.04
Shed net covered shed	5.75±0.56	5.375±0.55	-0.38 ± 0.04
Heat reflective colour painted shed	5.50±0.32	5.225±0.29	-0.25± 0.05
P value	0.759	0.056	0.002
Values are average of 4 reading.			
Values represents Mean± SD			
*Represents the values differs significantly across the column			

UNDER PEER REVIEW



**Fig. 2. Mean THI of normal asbestos roof shed, shed-net covered shed and heat reflective colour painted shed in summer**



**Fig. 3. Final decrease in milk yield of the cows kept under normal asbestos roof shed, shed-net covered shed and heat reflective colour painted shed in summer**

As per Joksimovic *et al.* [17] there was more decreased milk production in summer than in spring season. The decrease in daily average milk yield of the cows kept under heat reflective colour painted shed was observed to be 50% less than in case of cows kept normal asbestos shed indicating that there was less heat stress on cows in heat reflective colour painted shed during the summer season. Considering the milk yield at the start and end of the study, there was no statistical difference ( $P>0.05$ ) among the group of cows, but the drop in milk yield was significantly higher in asbestos roof animals indicating a higher level of heat stress resulting in drop in milk production. In a similar study, Mohapatra *et al.* [18] reported a drop of 0.2 L in average daily milk yield over one week by sprinkling water over the shed net covered roof during mid may, indicating a better microenvironment for cows in hot summer and advocated for sprinkling over shed net covered roof top as a measure to reduce the heat stress in milch cows. In the present study, only shed net was covered without sprinkling water, thus the reduction in milk was higher as compared to the reports by Mohapatra *et al.* [18].

#### **4. Conclusion:**

From 8.00 AM to 5.00PM, the THI of the normal asbestos roof shed varied from mild to moderate heat stress range. But THI of both shed-net covered shed and heat reflective colour painted shed was found to be within the mild heat stress range ie. 72-79 during noon indicating a comfortable ambient condition for the animals in summer season when the outside ambient temperature went up to 40 °C during noon. There was a decrease of 0.25 and 0.38 L in milk yield of the cows kept under heat reflective colour painted shed and shed-net covered shed, respectively. But a decrease of 0.45 L in milk yield of the cows kept under normal asbestos shed was recorded indicating more environmental stress on animals. Hence, the heat reflective colour painting over the dairy animal shed during peak summer may reduce the heat stress in dairy animals and tend to minimize the drop in milk production during hot environmental condition.

**Acknowledgement:** Authors are thankful to PC, AICRP on UAE, CIAE, Bhopal and ICAR, New Delhi, India for providing necessary support to carry out the experiment.

**DISCLAIMER (ARTIFICIAL INTELLIGENCE):** Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### **Reference:**

- [1] Habeeb AA, Gad AE, Atta MA. Temperature-humidity indices as indicators to heat stress of climatic conditions with relation to production and reproduction of farm animals. *International Journal of Biotechnology: Recent Advances*. 2018;1(1): 35-50.

- [2] Gaughan JB, Mader TL, Holt SM, Lisle A. A new heat load index for feedlot cattle. *Journal of Animal Science*. 2008;86:226-234.
- [3] Armstrong DV. Heat stress interaction with shade and cooling. *Journal of Dairy Science*. 1994;77:2044-2050.
- [4] Bohmanova J, Misztal I, Cole JB. Temperature-humidity indices as indicators of milk production losses due to heat stress. *Journal of Dairy Science*. 2007;90:1947-1956.
- [5] Das R, Sailo L, Verma N, Bharti P, Saikia J, Imtiwati, Kumar R. Impact of heat stress on health and performance of dairy animals: A review. *Veterinary World*. 2016;9:260-268.
- [6] Pragna P, Archana PR, Aleena J, Sejian V, Krishnan G, Bagath M, Manimaran A, Beena V, Kurien EK, Varma G, Bhatta R. Heat stress and dairy cow: Impact on both milk yield and composition. *International Journal of Dairy Science*. 2017;12:1-11.
- [7] Gantner V, Mijic P, Kuterovac K, Solic D, Gantner R. Temperature-humidity index values and their significance on the daily production of dairy cattle, *Mljekarstvo*. 2011;61(1):56-63.
- [8] Samer M. Adjusting dairy housing in hot climates to meet animal welfare requirements. *Journal of Experimental Sciences*. 2010;1(3):14-18.
- [9] Narwaria US, Singh M, Verma KK, Bharti PK. Amelioration of thermal stress using modified roof in dairy animals under tropics: A Review. *Journal of Animal Research*. 2017;7(5):801-812.
- [10] Liberati P. Analysis of the effects of the roofing design on heat stress in dairy cow housing. *Journal of Agricultural Engineering - Riv. di Ing. Agr.* 2008;4:1-7.
- [11] Khongdee S, Sripoon S, Chousawai S, Hinch G, Chaiyabutr N, Markvichitr K, Vajrabukka C. The effect of modified roofing on the milk yield and reproductive performance of heat-stressed dairy cows under hot-humid conditions. *Animal Science Journal*. 2010;81(5): 606-611.
- [12] Kamal R, Dutt T, Patel BHM, Ram RP, Biswas P, Bharti PK, Kaswan S. Effect of roofing materials on micro-climate in loose house for animals during rainy season. *Vet World*. 2013; 6(8):482-485.
- [13] Snedecor GW, Cochran WG. *Statistical Methods* 8th Edn. 1989. Iowa State University Press/Ames, Iowa-50010.

- [14] Vanitha V, Baskaran N. Effect of Ambient temperature in different seasons and roofing materials on ear flapping by Asian Elephants in captivity: A study from Tamil Nadu, India. *Journal of Indian Veterinary Association*, 2009;7(1):22-27.
- [15] Roy PK, Chatterjee A. Effect of different types of dairy cattle shelters on micro-climatic variables in rural West Bengal. *Indian Journal of Animal Science*. 2010;80(8):781-784.
- [16] West JW. Effects of heat-stress on production in dairy cattle. *Journal of Dairy Science*. 2003;86:2131-2144.
- [17] Joksimovic M, Davidovic V, Hristov S, Stankovic B. Effect of heat stress on milk production in dairy cows. *Biotechnology in Animal Husbandry*. 2011;27(3):1017-1023.
- [18] Mohapatra AK, Swain SK, Dash AK, Mohanty GP, Mishra SK and Behera D. Effect of Water Sprinkling Over Shed-net Covered Cattle Shed on Shed Environment and Milk Yield of Cows During Summer Season. *Ecology, Environment and Conservation*, 2023;29(3):1104-1108.
19. Mandal DK, Mandal A, Bhakat C, Dutta TK. Effect of heat stress amelioration through open-ridge ventilated thatched roof housing on production and reproduction performance of crossbred Jersey cows. *Tropical Animal Health and Production*. 2021 Mar;53(1):144.
20. Sahu D, Mandal DK, Podder M. Effect of loose house dairy cattle barn modification on udder health and production performance of Jersey crossbred cows in tropical lower Gangetic plains. *Journal of Dairy Research*. 2022 Nov;89(4):392-6.