

# Metabolic Shifting in Individual Parasitized with Helminths That May Contribute to Stunting

Comment [1]: Individuals

## ABSTRACT

**Aims:** to revisited the condition of chronic metabolic shifting caused by immunomodulatory sequence facilitated by helminth infection together with other deteriorating condition such as persistent exposure to infection which contributes to stunting formation.

Comment [2]: Revisit

**Discussion:** During dynamic interaction between active helminth infection and the host, the Helminth and or its excretory/secretory products induce and arouse the type 2 immune response which drives host tolerance and plays an important role in promoting tissue repair. Helminths also activates M2 Macrophages and induce a metabolic shift, even metabolic reprogramming towards reliance on oxidative phosphorylation, lipid oxidation and amino acid metabolism. Helminth-induced activation and metabolic reprogramming of macrophages precede the improvement in overall whole-body metabolism, denoted by improved insulin sensitivity, body mass in response to high-fat diet and atherogenic index in mammals. Contributions of altered nutrient uptake, adipose tissue function and/or the intestinal microbiota with the ability of helminths to alter metabolic status play a pivotal role in increased metabolism rate and may lead to wasting and even stunting formation.

Comment [3]: .

Comment [4]: helminth

Comment [5]: Remove 'or'

**Conclusion:** Helminth drives the type 2 immunity and activates its cell line which responsible for metabolic shifting and reprogramming.

Comment [6]: which is

Comment [7]: .

**Keywords:** biochemical, stunting, soil transmitted helminths, tropical disease

## 1. INTRODUCTION

Metabolism is the set of life-sustaining biochemical reactions in living organisms, specifically took place at the level of cells; its main business is to change food into energy [1]. This energy, specifically ATP, is the currency of energy among cells [2] to accomplish every aspect in living organism, from simple motoric activity [3] to complex cognitive-intelligentsia [4] even up to sophisticated bodily growth [5]. Specific key signaling proteins in the body control the chemical reactions of metabolism by responds not just to the existence of nutrients or metabolites but also to other growth signals, by that means choreographing the control of cellular metabolic processes [6].

Comment [8]: taking

Unfortunately, this wonderful metabolism blessing could be affected by several condition, one of them is the darling of tropical disease called chronic intestinal parasitic infection (specifically helminth) [7], especially when it happen in vulnerable subpopulation such as pregnant women [8] and can caused metabolic shifting [9]. Metabolic shifting evolved due to the periodic nature of the environment, occurs on a regular basis due to the demand-availability interplay of energy resources (food), expenditure (growth, infection etc.), and depots (excessive food intake), all of which have periodic nature due to the environment's circadian rhythmicity [9].

Comment [9]: responding

Comment [10]: ;

Comment [11]: Conditions;

Comment [12]: a

Comment [13]: happens

Comment [14]: subpopulations

Comment [15]: cause

Comment [16]: expenditures

Comment [17]: a

Metabolic shifting actually is the hallmark of most common diseases and the quest for the underlying unity of its pathogenesis is crucial [10]. Chronic metabolic shifting together with other deteriorating condition such as persistent exposure to infection, can caused catastrophe vicious circle in children called stunting [11,12] and this become the aim of this mini-review.

Comment [18]: is

Comment [19]: remove 'is'

Comment [20]: conditions

Comment [21]: cause a

Comment [22]: became

## 2. STUNTING AT A GLANCE

A condition of chronic and/or recurring deficiency of correct nutritious foods, both in quantity and quality, for mother [8] and children during the whole pre- and postnatal periods [12-14] subsidizes the formation of stunting [7]. Children who are stunted (length-for-age Z-score<-2) are at greater risk of infectious morbidity and mortality [12].

Comment [23]: Replace with 'and/or'

Comment [24]: mothers

Comment [25]: prenatal

The most well-accepted pathway to the existence of stunting is a complex 'vicious cycle' between long term nutritional deprivation [13] and chronic-persistent infection [14], which is progressing negatively to fence in microbiota dysbiosis of the intestine [15] which causes the disappearance of intestinal immune cells function [16], and triggers the local intestinal inflammation [17,18], and when it happens in the specific portion of intestinal walls defined as leaky gut, [19] a weakening of the intestinal walls which in turn allows bacteria and toxins releases into the bloodstream. It then will trigger chronic, full body inflammation or systemic inflammatory responses [20]. Adjacent to energetic, hormonal, and metabolic sequel, anemia is also on numerous occasion experienced as co-morbid condition of stunting [21]. Furthermore, anemia found in mother [8,22] and/or child [23] may be overlapping causing stunting.

Comment [26]: cell

Comment [27]: happens

Comment [28]: to release

Comment [29]: sequela

Common chronic persistent infection among children is caused by intestinal parasites [7], mostly the soil transmitted helminths, or STH [30], which are parasitic nematode worms that live in the human intestine [31]. The three most common STH infecting human are: round worm *Ascaris lumbricoides*, whip worm *Trichuris trichiura* and hookworm *Ancylostoma duodenale* and *Necator americanus*. They easily disseminate through soil or water contaminated by human stool that contain the parasite larvae [30,31]. STH infections are considered the great neglected tropical diseases or NTDs [32].

Comment [30]: humans

Comment [31]: Contains

## 3. HOW INTESTINAL PARASITIC INFECTION CONTRIBUTES TO STUNTING

In the intestine, parasites occupy the same niche as bacteria, member of the normal microbiota [24]. The coming of new comer intestinal parasites can disrupt the normal balance of the gut microbiota [24,25] and can open the door for other bodily derangement [26]. The two-way relationship between intestinal parasites and the microbiota [17, 27], sometime it is already happening in very early life just as revealed by Hakizimana *et al* [28], and their combined reciprocally strengthen effects [14], could play a key role in stunting [12-14,17,23]. Epigenetic regulation of gene expression in children at risk of stunting [29] may link parasitic infections and poor intestinal health and function [14-20] since very early life and may lead to stunting formation.

Comment [32]: members

Comment [33]: strengthened

The normal gut microbiome accommodates all good bacteria [34], specific fungi [35], protozoa which sometime classified as neglected gut microbiome [36], the less common reported archaea [37], and even an abundance of viruses [37]- these all well-function biomes [38] that reside in and along the gastrointestinal tract, especially in the intestine [39] and may occupy their milieu commensally [40]. The composition of the gut microbiota is probably the result of a process involving ecological memory, and not just simple host-based filtering [41]. Its composition is always in dynamic and fluctuant state [42] and, probably, with a double face, favorable or detrimental to the host [43].

Comment [34]: are sometimes

Comment [35]: microbiomes

Comment [36]: are

Comment [37]: a

The gut microbiome plays important roles in (1) process of metabolism - i.e., the metabolism of nutrients, such as indigestible polysaccharides, lipids, vitamins, and AAs [44], (2) gut epithelial well-being [34,37,39,40], and (3) control the systemic innate [45] and adaptive [46] immune responses, even to the extent of promoting immune tolerance [47]. As far as is known, the human host and microbes have coevolved throughout evolutionary history in the gastrointestinal niche [48,49], and it is postulated that their closed interaction [50] have adapted to exert dynamic influences on each other [51]. Helminth infection increases the host's microbiome diversity, together with its richness [52]. Ongoing helminth infection has been associated with the suppression of bodily allergies reaction [53] and also a converted susceptibility to certain microbial infections due to the shift in the host's immune response [54].

Comment [38]: the process

Comment [39]: controlling

Comment [40]: has

Comment [41]: reactions

Study revealed that active and chronic helminth infections can protect against allergic ailment by an active regulatory process [55] by the aid of its intense modulation of the specific host immune system dictated by expanding repertoire of parasite effector molecules [56] (e.g., Anti-inflammatory protein-1, a family member of tissue inhibitor of metalloprotease (TIMP)-like proteins secreted by *N. americanus*, which act as suppression in TNBS colitis model: promotes expression of colon IL-10, TGF- $\beta$ , and TSLP and the accumulation of Treg cells

in the colon and Anti-inflammatory protein-2 also secreted by *N. americanus*, a family member of TIMP-like proteins which causes suppression in model of allergic airway inflammation via Treg cell induction and suppression of T cell proliferation in cells from house dust mite (HDM)-allergic patients), therefore leading to a systemic T-cell hyporesponsiveness to parasite antigen [57], which is triggered by the induction of a complex regulatory network initiated by the regulatory cytokine IL-10 [58] and TGF- $\beta$  family [57]. Basically, helminth-induced immunoregulation occurs through the induction of regulatory T cells [60] and or Th2-type cells [61], and this a role for Treg in limiting inflammation-induced pathology following helminth infection [60]. An increased proportion and expansion of adaptive Treg cells is also reported by Santos et al in Brazil [62] which reported *Ascaris lumbricoides* coinfection with active tuberculosis actually reduces infected tissue damage by reducing IL-6 levels without altering the clinical progression of pulmonary tuberculosis or Th1/Th2/Th17 cytokine profile.

Comment [42]: this plays

Although, significant complex environmental exposure on individual member of the community causing dynamic variation regarding regular daily diet [63], personal hygiene [64], environmental sanitation condition which contributes to ecologically based natural selection [65], helminth related species and burden which infected [66] and so many other factors that make it strenuous to accomplish consensus regarding the consequences of helminths on the gut microbiome reciprocally. Inconsistent associations between active helminth infection and its related parameters such as species abundance, diversity and taxonomic plethora commonly used to profile the microbiome actually already revealed by previous literature review study. These inconsistencies advocate that factors such as host genetics (e.g., in relation to the host's gender and genotype) [67], nutrition [68], the history of exposure to anthelmintic or other treatment [69] and the condition of co-infections [70] may also exert influence on the reciprocal effects of helminths on the microbiome or the microbiome on helminths. This interesting pile of knowledge bolster up the need for more extensive study, perhaps with meta-analyses, to accommodate all the data with the approach of cross-study and cross-population in order get better understanding regarding the role of dynamic interaction between the host's microbiome, the parasite and the environment. In the next section, we are going to link this interaction with the metabolic shifting experienced by individual which chronically infested with parasite.

Comment [43]: members

Comment [44]: conditions

Comment [45]: Remove 's'

Comment [46]: studies

Comment [47]: bolsters

Comment [48]: to

Comment [49]: individuals

Comment [50]: who are

#### 4. METABOLIC SHIFTING IN INDIVIDUAL INFESTATED WITH CHRONIC PARASITIC DISEASE

Here, we provide a comprehensive review regarding this sub topic subtopic. Even during earliest exhibition to the invasion of parasites, hosts can establish behavioral and or physiological responses [72]. Those responses may occur without the host realize or fully aware about it. The tendency of chronicity regarding parasitic infection and inability to combat innumerable species of parasitic helminths that become co-evolved with their preference hosts over history of mankind proposes that certain mammals have accustomed mechanisms which tolerate this specific infectious disease [73]. Those condition expose the host to certain metabolic costs [71,74]. Ongoing inflammatory and immune responses are associated with dramatic shifts in tissue metabolism. These changes include local and persistent depletion of important nutrients [13], increased oxygen consumption [4], and the generation of large quantities of reactive nitrogen and oxygen intermediates [20]. Such enormous shifts in tissue metabolism derive, at least in part, from profound recruitment of inflammatory cell types, particularly myeloid cells, such as neutrophils (polymorphonuclear leukocytes [PMNs]) and monocytes [44]. In reality, the vast majority of inflammatory cells are directed to, as opposed to resident at, the region infected and suffer inflammatory lesions [18]. By stark contrast, adaptive immune responses are characterized by high rates of local T and B cell proliferation and have significantly different metabolic demands. Detrimental effects of parasitism on host fitness are frequently attributed to parasite-associated demands to host's energy [74].

Comment [51]: realizing

Comment [52]: the

Comment [53]: that

Comment [54]: conditions

Unfortunately, few studies have measured these costs directly. Hence, little is known about metabolic costs arising from parasite exposure. Furthermore, no one has yet measured whether and how previous infection history modulates metabolic responses to parasite exposure.

In case of chronic (intestinal) helminth infection, there is an interplay of type 2 immunity, helminth infection and the microbiota in regulating the host's metabolism [75]. During on going helminth infection, type 2 immunity is required to limit worm burdens and to promote the timely repair the damage caused by tissue migrating larval stages of the parasite. With the involvement of Type 2 immunity, which recently has recently noticed as a crucial player in the host's metabolic status [76], with innumerable studies analyzing the role of type 2 immune cells within adipose tissue [77,78].

Comment [55]: of

Comment [56]: been

Metabolic dysfunction which occur during active helminth infection is repeatedly designated as a sustained low-grade or chronic inflammatory state within infected part of the tissues [79], and type 2 immunity may simplify a return to the condition of metabolic homeostasis, as part of the host protective function [80]. Ideally, the immune response is designed to the specific milieu in which it takes place [12,16]. Immune cells have the special ability to sense and adapt to fluctuates in their surroundings, and it is now getting more noticed that in

Comment [57]: the

Comment [58]: fluctuations

addition to cytokines made by stromal and epithelial cells, metabolic cues provide key adaptation signals. Changes in immune cell activation states are linked to changes in cellular metabolism that support its function [81]. Furthermore, metabolites themselves can signal between as well as within cells [40,79,81]. Metabolic regulation relates to type 2 immunity firstly by considering specifics of metabolism within type 2 immune cells and secondly by stressing how type 2 immune cells are integrated more broadly into the metabolism of the organism as a whole [81].

A complex network of type 2 resident cells including ~~The~~ anti-inflammatory M2 macrophages [82], eosinophils [83] and group 2 innate lymphoid cells (ILC2s) [84] has been properly identified within adipose tissue. In spite of the fact that the exact effector cells in this equilibrium have not been intelligibly recognized, any change of the type 2 microenvironment resulted in a shifted metabolic state [85]. This shifted metabolic state of the host sometime also called metabolic reprogramming [9,10,85-87].

The characterization [metabolic reprogramming toward glycolysis, which favors immune cell activation and its effector, is commonly seen in sepsis [86,87]. Their functional implication in anti-inflammatory [88], immune regulation [89] and later pro-inflammatory [20] responses are actually important built in features to survive from catastrophic infection. In the case of helminth parasites, they are complex metazoans that belong to different taxonomic families but collectively share the ability to downregulate the host immune response directed toward themselves (parasite-specific immunoregulation). During long-standing chronic infection, these helminths appear to have the ability to dictate its host by way of suppress immune responses to bystander pathogens/antigens and atopic, autoimmune, and metabolic disorders.

At least, there are two major features of metabolic reprogramming in inflammatory condition, in innate and as well as in adaptive immune cells, namely as follows:

- 1) energy production and biosynthesis reprogramming, including increased glycolysis and decreased oxidative phosphorylation, in order to secure faster ATP production and biosynthesis for defense response and damage repair, and
- 2) epigenetic reprogramming, including enhanced histone acetylation and suppressed DNA methylation, due to altered accessibility of acetyl/methyl group donor and metabolite-modulated enzymatic activity. Finally, we discuss current strategies of metabolic and epigenetic therapy in cardiovascular disease and recommend cell-specific metabolic and gene-targeted site-specific epigenetic alterations for future therapies.




Fig 1. Schematic steps of general interplay between immune response and metabolic reprogramming. Activation of innate immune cells by pathogen/danger-associated molecular pattern or metabolite-associated danger signal via pattern recognition receptors or metabolic sensor to facilitate the downstream signaling cascade to initiate an immune response. Metabolic reprogramming exists during this type of immune responses, in which various cellular metabolic processes in immune cells are altered to achieve an adequate function. Complex process may be involved and finally lead to a metabolic optimization in which distinct metabolic signature determines specific inflammatory or anti-inflammatory immune cell subset differentiation [86 with modification].

Comment [59]: sometimes

Comment [60]: of

Comment [61]: implications

The metabolic response of the host to helminth infection usually leads to modification in the pattern of amino acids in the body fluids [90], remodeling of the lipid metabolism, *i.e.*, a depletion of the amino acids pool, an alteration of the ketogenic pathways [79], and changes in composition of microbiota-related metabolites [54]. This profound metabolic shift also has an intense influence on the host immune status and eventually affects organs that are not directly involved in the immune response.




Fig 2. Schematic representation of putative mechanisms by which helminth infection could alter the metabolic state of the host. Helminths typically elicit a type 2 immune response which has been associated with modulation of adipose tissue homeostasis, whole-body metabolic changes and changes in intestinal physiology [75, with modification].

Helminths can make adjustments to harsh environmental conditions inside the host [91], such as utilizing anaerobic cycle such as fermentation and malate dismutation to obtain energy from carbohydrate they steal from their host [92]. Carbohydrate is an essential energy source in all mature helminths and its metabolism process is often primarily through anaerobic cycle, even in the milieu where oxygen is abundant. Helminths also use amino acid, polyunsaturated fatty acid (PUFA), and cholesterol metabolism, a possible strategy favoring the production of immunomodulatory compounds that may influence survival in the host and evade host immunity [93].

It is clear that helminth infections can cause severe problems such as stunting in undernourished children by way of attenuation of systemic inflammation and metabolic reprogramming. Further study need to be conducted in order to explore metabolically mediated immunosuppressive status of the vulnerable host which infected with helminth; what species, and for how long, and also with the history of previous treatment for a chronic helminth infection.

#### 4. CONCLUSION

Chronic metabolic shifting and reprogramming during helminth infection usually occurring together with other deteriorating condition such as persistent exposure to infection and prolonged inflammation that all together can cause stunting. Type 2 immune response induced by helminth infection drives host tolerance and plays an important role in promoting tissue repair. Helminths polarizes in to M2 Macrophages and induce a metabolic shift towards reliance on oxidative phosphorylation, lipid oxidation and amino acid metabolism. Helminth-induced activation and metabolic reprogramming of macrophages underlie improvement in overall whole-body metabolism, denoted by improved insulin sensitivity, body mass in response to high-fat diet and atherogenic index in mammals. Contributions of altered nutrient uptake, adipose tissue function and/or the intestinal microbiota with the ability of helminths to alter metabolic status play a pivotal role in increased metabolism rate and may lead to wasting and even stunting formation.

Comment [62]: acid  
 Comment [63]: the

Comment [64]: cycles  
 Comment [65]: an

Comment [66]: studies  
 Comment [67]: is

Comment [68]: occur

Comment [69]: a

## CONSENT (WHERE EVER APPLICABLE)

Not needed

## ETHICAL APPROVAL (WHERE EVER APPLICABLE)

Not needed

## REFERENCES

1. Sánchez López de Nava A, Raja A. Physiology, Metabolism. [Updated 2022 Sep 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK546690/>
2. Pinna S, Kunz C, Halpern A, Harrison SA, Jordan SF, Ward J, et al. A prebiotic basis for ATP as the universal energy currency. *PLoS Biol.* 2022 Oct 4;20(10):e3001437. <https://doi.org/10.1371/journal.pbio.3001437>.
3. Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. *Nat Metab.* 2020; 2, 817–28. <https://doi.org/10.1038/s42255-020-0251-4>
4. Meijer A, Königs M, Pouwels PJW, Smith J, Visscher C, Bosker RJ, Hartman E, Oosterlaan J. Effects of aerobic versus cognitively demanding exercise interventions on brain structure and function in healthy children-Results from a cluster randomized controlled trial. *Psychophysiology.* 2022 Aug;59(8):e14034. <https://doi.org/10.1111/psyp.14034>.
5. Mahdi S, Dickerson A, Infield Solar G, Caton SJ. Timing of energy intake and BMI in children: differential impacts by age and sex. *British Journal of Nutrition.* 2023;130(1):71-82. <https://doi.org/10.1017/S0007114522003014>
6. Szwed A, Kim E, Jacinto E. Regulation and metabolic functions of mTORC1 and mTORC2. *Physiol Rev.* 2021 Jul 1;101(3):1371-1426. <https://doi.org/10.1152/physrev.00026.2020>.
7. Siagian FE. Intestinal Parasitic Infection Responsible for Undernourishment and Stunted Growth in Children of School Going Ageing. *Asian Journal of Research in Infectious Diseases.* 2023; 14(1): 18-25. <https://doi.org/10.9734/AJRID/2023/v14i1278>
8. Nainggolan S, Siagian FE. The prevalence of anemia in pregnant women in the 10 priority villages for stunting control in Sumedang district, West Java: a community-based survey. *International Journal of Community Medicine And Public Health.* 2019; 6 (9): 3760-7. <https://doi.org/10.18203/2394-6040.ijcmph20193966>
9. Zilberter T, Paoli A. Editorial: Metabolic Shifting: Nutrition, Exercise, and Timing. *Front Nutr.* 2020 Dec 10;7:592863. <https://doi.org/10.3389/fnut.2020.592863>.
10. Schwartz L, Henry M, Alfarouk KO, Reshkin SJ, Radman M. Metabolic Shifts as the Hallmark of Most Common Diseases: The Quest for the Underlying Unity. *Int J Mol Sci.* 2021 Apr 12;22(8):3972. <https://doi.org/10.3390/ijms22083972>.
11. de Moraes RCS, Sawaya AL, Vieira ACA, et al. Food addiction symptoms and metabolic changes in children and adolescents with the double burden of malnutrition. *British Journal of Nutrition.* 2021;126(12):1911-1918. <https://doi.org/10.1017/S0007114521000313>
12. Mutasa K, Tome J, Rukobo S, Govha M, Mushayanembwa P, Matimba FS, Chiorera CK, Majo FD, Tavengwa NV, Mutasa B, Chasekwa B, Humphrey JH, Ntozini R, Prendergast AJ, Bourke CD. Stunting Status and Exposure to Infection and Inflammation in Early Life Shape Antibacterial Immune Cell Function Among Zimbabwean Children. *Front Immunol.* 2022 Jun 13;13:899296. <https://doi.org/10.3389/fimmu.2022.899296>.
13. Millward DJ. Nutrition, infection and stunting: the roles of deficiencies of individual nutrients and foods, and of inflammation, as determinants of reduced linear growth of children. *Nutrition Research Reviews.* 2017;30(1):50-72. <https://doi.org/10.1017/S0954422416000238>
14. Gabain IL, Ramsteijn AS, Webster JP. Parasites and childhood stunting - a mechanistic interplay with nutrition, anemia, gut health, microbiota, and epigenetics. *Trends Parasitol.* 2023 Mar;39(3):167-180. <https://doi.org/10.1016/j.pt.2022.12.004>.

15. Hrnčir T. Gut Microbiota Dysbiosis: Triggers, Consequences, Diagnostic and Therapeutic Options. *Microorganisms*. 2022 Mar 7;10(3):578. <https://doi.org/10.3390/microorganisms10030578>.
16. Fakharian F, Thirugnanam S, Welsh DA, Kim W-K, Rappaport J, Bittinger K, Rout N. The Role of Gut Dysbiosis in the Loss of Intestinal Immune Cell Functions and Viral Pathogenesis. *Microorganisms*. 2023; 11(7):1849. <https://doi.org/10.3390/microorganisms11071849>
17. Lefebo BK, Kassa DH, Tarekegn BG. Factors associated with stunting: gut inflammation and child and maternal-related contributors among under-five children in Hawassa City, Sidama Region, Ethiopia. *BMC Nutr* 2023;9: 54. <https://doi.org/10.1186/s40795-023-00701-4>
18. Zeng MY, Inohara N, Nuñez G. Mechanisms of inflammation-driven bacterial dysbiosis in the gut. *Mucosal Immunol*. 2017 Jan;10(1):18-26. <https://doi.org/10.1038/mi.2016.75>.
19. Camilleri M. Leaky gut: mechanisms, measurement and clinical implications in humans. *Gut*. 2019 Aug;68(8):1516-1526. <https://doi.org/10.1136/gutjnl-2019-318427>.
20. Patterson GT, Osorio EY, Peniche A, Dann SM, Cordova E, Preidis GA, Suh JH, Ito I, Saldarriaga OA, Loeffelholz M, Ajami NJ, Travi BL, Melby PC. Pathologic Inflammation in Malnutrition Is Driven by Proinflammatory Intestinal Microbiota, Large Intestine Barrier Dysfunction, and Translocation of Bacterial Lipopolysaccharide. *Front Immunol*. 2022 May 26;13:846155. <https://doi.org/10.3389/fimmu.2022.846155>.
21. Sahiledengle B, Mwanri L, Petrucka P, Agho KE. Coexistence of Anaemia and Stunting among Children Aged 6-59 Months in Ethiopia: Findings from the Nationally Representative Cross-Sectional Study. *Int J Environ Res Public Health*. 2023 Jun 29;20(13):6251. <https://doi.org/10.3390/ijerph20136251>.
22. Dessie G, Li J, Nghiem S, Doan T. Prevalence and Determinants of Stunting-Anemia and Wasting-Anemia Comorbidities and Micronutrient Deficiencies in Children Under 5 in the Least-Developed Countries: A Systematic Review and Meta-analysis, *Nutrition Reviews*, 2024;, nuae063, <https://doi.org/10.1093/nutrit/nuae063>
23. Nadhiroh SR, Micheala F, Tung SEH, Kustiawan TC. Association between maternal anemia and stunting in infants and children aged 0-60 months: A systematic literature review. *Nutrition*. 2023 Nov;115:112094. <https://doi.org/10.1016/j.nut.2023.112094>.
24. Glendinning L, Nausch N, Free A, Taylor DW, Mutapi F. The microbiota and helminths: sharing the same niche in the human host. *Parasitology*. 2014;141(10):1255-71. <https://doi.org/10.1017/S0031182014000699>
25. Beyhan YE, Yildiz MR. Microbiota and parasite relationship. *Diagn Microbiol Infect Dis*. 2023 Aug;106(4):115954. <https://doi.org/10.1016/j.diagmicrobio.2023.115954>.
26. Ramírez-Carrillo E, Gaona O, Nieto J, Sánchez-Quinto A, Cerqueda-García D, Falcón LI, Rojas-Ramos OA, González-Santoyo I. Disturbance in human gut microbiota networks by parasites and its implications in the incidence of depression. *Sci Rep*. 2020 Feb 28;10(1):3680. <https://doi.org/10.1038/s41598-020-60562-w>.
27. Siagian FE. Intestinal Microflora Vs Protozoan Parasites: From Interaction to Competition. *South Asian Journal of Research in Microbiology*, 2022;13(1): 36-46, <https://doi.org/10.9734/SAJRM/2022/v13i130290>
28. Hakizimana E, Kim JY, Oh S, Yoon M, Yong TS. Intestinal parasitic infections among children aged 12-59 months in Nyamasheke District, Rwanda. *Parasites Hosts Dis*. 2023 Aug;61(3):304-309. <https://doi.org/10.3347/PHD.23045>.
29. Ramsteijn AS, Ndiaye M, Kalashikam RR. Epigenetic studies in children at risk of stunting and their parents in India, Indonesia and Senegal: a UKRI GCRF Action Against Stunting Hub protocol paper. *BMJ Paediatrics Open*, 2024;8:e001770. <https://doi.org/10.1136/bmjpo-2022-001770>
30. Loukas A, Maizels RM, Hotez PJ. The yin and yang of human soil-transmitted helminth infections. *Int J Parasitol*. 2021 Dec;51(13-14):1243-1253. <https://doi.org/10.1016/j.ijpara.2021.11.001>.
31. Al Amin ASM, Wadhwa R. Helminthiasis. [Updated 2023 Jul 17]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK560525/>
32. Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. *J Clin Invest*. 2008 Apr;118(4):1311-21. <https://doi.org/10.1172/JCI34261>.
33. Sunarti LS. Microbial Normal Flora: Its Existence And Their Contribution To Homeostasis. *Journal of Advances in Microbiology*,2022; 22(9): 1-15, <https://doi.org/10.9734/JAMB/2022/v22i930483>

34. Zhang F, Aschenbrenner D, Yoo JY, Zuo T. The gut mycobiome in health, disease, and clinical applications in association with the gut bacterial microbiome assembly. *Lancet Microbe*. 2022 Dec;3(12):e969-e983. [https://doi.org/10.1016/S2666-5247\(22\)00203-8](https://doi.org/10.1016/S2666-5247(22)00203-8).
35. Guzzo GL, Andrews JM, Weyrich LS. The Neglected Gut Microbiome: Fungi, Protozoa, and Bacteriophages in Inflammatory Bowel Disease. *Inflamm Bowel Dis*. 2022 Jul 1;28(7):1112-1122. <https://doi.org/10.1093/ibd/izab343>.
36. Mafra D, Ribeiro M, Fonseca L, Regis B, Cardozo LFMF, Fragoso Dos Santos H, Emiliano de Jesus H, Schultz J, Shiels PG, Stenvinkel P, Rosado A. Archaea from the gut microbiota of humans: Could be linked to chronic diseases? *Anaerobe*. 2022 Oct;77:102629. <https://doi.org/10.1016/j.anaerobe.2022.102629>.
37. Cao Z, Sugimura N, Burgermeister E, Ebert MP, Zuo T, Lan P. The gut virome: A new microbiome component in health and disease. *EBioMedicine*. 2022 Jul;81:104113. <https://doi.org/10.1016/j.ebiom.2022.104113>.
38. Vemuri R, Shankar EM, Chieppa M, Eri R, Kavanagh K. Beyond Just Bacteria: Functional Biomes in the Gut Ecosystem Including Virome, Mycobiome, Archaeome and Helminths. *Microorganisms*. 2020; 8(4):483. <https://doi.org/10.3390/microorganisms8040483>
39. Dupont HL, Jiang ZD, Dupont AW, Utay NS. The intestinal microbiome in human health and disease. *Trans Am Clin Climatol Assoc*. 2020;131:178-197.
40. Krishnamurthy HK, Pereira M, Bosco J, George J, Jayaraman V, Krishna K, Wang T, Bei K, Rajasekaran JJ. Gut commensals and their metabolites in health and disease. *Front Microbiol*. 2023 Nov 8;14:1244293. <https://doi.org/10.3389/fmicb.2023.1244293>.
41. Letourneau J, Holmes ZC, Dallow EP, Durand HK, Jiang S, Carrion VM, Gupta SK, Mincey AC, Muehlbauer MJ, Bain JR, David LA. Ecological memory of prior nutrient exposure in the human gut microbiome. *ISME J*. 2022 Nov;16(11):2479-2490. <https://doi.org/10.1038/s41396-022-01292-x>.
42. Minagar A, Jabbour R, Jabbour H (eds). *The Human Gut Microbiota: A Dynamic Biologic Factory*. In: *Advances in Biochemical Engineering/Biotechnology*. Springer, Berlin, Heidelberg. 2023. [https://doi.org/10.1007/10\\_2023\\_243](https://doi.org/10.1007/10_2023_243)
43. Cani PD. Human gut microbiome: hopes, threats and promises *Gut* 2018;67:1716-1725.
44. Fujisaka S, Watanabe Y, Tobe K. The gut microbiome: a core regulator of metabolism. *Journal of Endocrinology* 2023;256 (3): e220111, <https://doi.org/10.1530/JOE-22-0111>
45. Jordan CKI, Clarke TB. How does the microbiota control systemic innate immunity? *Trends Immunol*. 2024 Feb;45(2):94-102. <https://doi.org/10.1016/j.it.2023.12.002>.
46. Li Y, Ye Z, Zhu J, Fang S, Meng L, Zhou C. Effects of Gut Microbiota on Host Adaptive Immunity Under Immune Homeostasis and Tumor Pathology State. *Front Immunol*. 2022 Mar 10;13:844335. <https://doi.org/10.3389/fimmu.2022.844335>.
47. Ning X, Lei Z, Rui B, Li Y, Li M. Gut Microbiota Promotes Immune Tolerance by Regulating RORγt+ Treg Cells in Food Allergy, *Advanced Gut & Microbiome Research*, 2022, 8529578, 9 pages, 2022. <https://doi.org/10.1155/2022/8529578>
48. Shahab M, Shahab N. Coevolution of the Human Host and Gut Microbiome: Metagenomics of Microbiota. *Cureus*, 2022; 14(6): e26310. <https://doi.org/10.7759/cureus.26310>
49. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, et al. Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nat Commun*. 2019 May 16;10(1):2200. <https://doi.org/10.1038/s41467-019-10191-3>.
50. Llinás-Caballero K, Caraballo L. Helminths and Bacterial Microbiota: The Interactions of Two of Humans' "Old Friends". *International Journal of Molecular Sciences*. 2022; 23(21):13358. <https://doi.org/10.3390/ijms232113358>
51. Loke P, Harris NL. Networking between helminths, microbes, and mammals. *Cell Host Microbe*. 2023 Apr 12;31(4):464-471. <https://doi.org/10.1016/j.chom.2023.02.008>.
52. Walusimbi B, Lawson MAE, Jacent N, Kateete DP, David P, Webb EL, et al. The effects of helminth infections on the human gut microbiome: a systematic review and meta-analysis. *Frontiers in Microbiomes*, 2023;2. <https://doi.org/10.3389/frmbi.2023.1174034>
53. Maizels RM. Regulation of immunity and allergy by helminth parasites. *Allergy*. 2020 Mar;75(3):524-534. <https://doi.org/10.1111/all.13944>.
54. Brosschot TP, Reynolds LA. The impact of a helminth-modified microbiome on host immunity. *Mucosal Immunol*. 2018 Jul;11(4):1039-1046. <https://doi.org/10.1038/s41385-018-0008-5>.

55. Smits HH, Everts B, Hartgers FC, Yazdanbakhsh M. Chronic helminth infections protect against allergic diseases by active regulatory processes. *Curr Allergy Asthma Rep.* 2010 Jan;10(1):3-12. <https://doi.org/10.1007/s11882-009-0085-3>.
56. Maizels RM, Smits HH, McSorley HJ. Modulation of Host Immunity by Helminths: The Expanding Repertoire of Parasite Effector Molecules. *Immunity.* 2018 Nov 20;49(5):801-818. <https://doi.org/10.1016/j.immuni.2018.10.016>.
57. White MPJ, McManus CM, Maizels RM. Regulatory T-cells in helminth infection: induction, function and therapeutic potential. *Immunology.* 2020 Jul;160(3):248-260. <https://doi.org/10.1111/imm.13190>.
58. Figueiredo CA, Barreto ML, Rodrigues LC, Cooper PJ, Silva NB, Amorim LD, et al. Chronic intestinal helminth infections are associated with immune hyporesponsiveness and induction of a regulatory network. *Infect Immun.* 2010 Jul;78(7):3160-7. <https://doi.org/10.1128/IAI.01228-09>.
59. Ince MN, Elliott DE, Setiawan T, Metwali A, Blum A, Chen HL, Urban JF, Flavell RA, Weinstock JV. Role of T cell TGF-beta signaling in intestinal cytokine responses and helminthic immune modulation. *Eur J Immunol.* 2009 Jul;39(7):1870-8. <https://doi.org/10.1002/eji.200838956>.
60. Smith KA, Filbey KJ, Reynolds LA, Hewitson JP, Harcus Y, Boon L, Sparwasser T, Hämmerling G, Maizels RM. Low-level regulatory T-cell activity is essential for functional type-2 effector immunity to expel gastrointestinal helminths. *Mucosal Immunol.* 2016 Mar;9(2):428-43. <https://doi.org/10.1038/mi.2015.73>.
61. Nutman TB. Looking beyond the induction of Th2 responses to explain immunomodulation by helminths. *Parasite Immunol.* 2015 Jun;37(6):304-13. <https://doi.org/10.1111/pim.12194>.
62. Santos JHA, Bühner-Sékula S, Melo GC, Cordeiro-Santos M, Pimentel JPD, Gomes-Silva A, et al. *Ascaris lumbricoides* coinfection reduces tissue damage by decreasing IL-6 levels without altering clinical evolution of pulmonary tuberculosis or Th1/Th2/Th17 cytokine profile. *Rev. Soc. Bras. Med. Trop* 2019; 52: e20190315. <https://doi.org/10.1590/0037-8682-0315-2019>
63. Myhill LJ, Williams AR. Diet-microbiota crosstalk and immunity to helminth infection. *Parasite Immunology.* 2023 Apr;45(4):e12965. <https://doi.org/10.1111/pim.12965>.
64. Loke P, Lim YA. Helminths and the microbiota: parts of the hygiene hypothesis. *Parasite Immunol.* 2015 Jun;37(6):314-23. <https://doi.org/10.1111/pim.12193>.
65. Salloum PM, Jorge F, Dheilly NM, Poulin R. Eco-evolutionary implications of helminth microbiomes. *Journal of Helminthology.* 2023;97:e22. <https://doi.org/10.1017/S0022149X23000056>
66. Rosa BA, Supali T, Gankpala L, Djuardi Y, Sartono E, Zhou Y, et al. Differential human gut microbiome assemblages during soil-transmitted helminth infections in Indonesia and Liberia. *Microbiome,* 2018;6. <https://doi.org/10.1186/s40168-018-0416-5>
67. Ling F, Steinel N, Weber J, Ma L, Smith C, Correa D, et al. The gut microbiota response to helminth infection depends on host sex and genotype. *ISME J.* 2020 May;14(5):1141-1153. <https://doi.org/10.1038/s41396-020-0589-3>.
68. Cattadori IM, Sebastian A, Hao H, Katani R, Albert I, Eilertson KE, et al. Impact of Helminth Infections and Nutritional Constraints on the Small Intestine Microbiota. *PLoS One.* 2016 Jul 20;11(7):e0159770. <https://doi.org/10.1371/journal.pone.0159770>.
69. Easton AV, Quiñones M, Vujkovic-Cvijin I, Oliveira RG, Kepha S, Odiere MR, et al. The Impact of Anthelmintic Treatment on Human Gut Microbiota Based on Cross-Sectional and Pre- and Postdeworming Comparisons in Western Kenya. *mBio.* 2019 Apr 23;10(2):e00519-19. <https://doi.org/10.1128/mBio.00519-19>.
70. Schmid, DW, Fackelmann G., Wasimuddin. A framework for testing the impact of co-infections on host gut microbiomes. *anim microbiome,* 2022; 4, 48. <https://doi.org/10.1186/s42523-022-00198-5>
71. Nadler LE, Bengston E, Eliason EJ, Hassibi C, Helland-Riise SH, Johansen IB, et al. A brain-infecting parasite impacts host metabolism both during exposure and after infection is established. *Functional Ecology,* 2020; (00) : 1 -12. <https://doi.org/10.1111/1365-2435.13695>
72. Thompson SN, Kavaliers M. Physiological bases for parasite-induced alterations of host behaviour. *Parasitology.* 1994;109(S1):S119-S138. <https://doi.org/10.1017/S0031182000085139>
73. King IL, Li Y. Host-Parasite Interactions Promote Disease Tolerance to Intestinal Helminth Infection. *Front Immunol.* 2018 Sep 20;9:2128. <https://doi.org/10.3389/fimmu.2018.02128>.
74. Robar N, Murray DL, Burness G. Effects of parasites on host energy expenditure: the resting metabolic rate stalemate. *Canadian Journal of Zoology.* 2011;89(11): 1146-1155. <https://doi.org/10.1139/z11-084>

75. Moyat M, Coakley G, Harris NL. The interplay of type 2 immunity, helminth infection and the microbiota in regulating metabolism. *Clin Transl Immunology*. 2019 Nov 7;8(11):e01089. doi: <https://doi.org/10.1002/cti2.1089>.
76. Sikder S, Pierce D, Sarkar ER, McHugh C, Quinlan KGR, Giacomini P, Loukas A. Regulation of host metabolic health by parasitic helminths. *Trends Parasitol*. 2024 May;40(5):386-400. doi: <https://doi.org/10.1016/j.pt.2024.03.006>.
77. Man K, Kallies A, Vasanthakumar A. Resident and migratory adipose immune cells control systemic metabolism and thermogenesis. *Cell Mol Immunol*, 2022; 19: 421–431. <https://doi.org/10.1038/s41423-021-00804-7>
78. Wu D, Yifu Qiu Y. Type 2 immune regulation of adipose tissue homeostasis. *Current Opinion in Physiology*, 2019; 12:20-5. <https://doi.org/10.1016/j.cophys.2019.04.018>.
79. Kokova D, Verhoeven A, Perina EA, Ivanov VV, Heijink M, Yazdanbakhsh M, Mayboroda OA. Metabolic Homeostasis in Chronic Helminth Infection Is Sustained by Organ-Specific Metabolic Rewiring. *ACS Infect Dis*. 2021 Apr 9;7(4):906-916. <https://doi.org/10.1021/acsinfectdis.1c00026>.
80. Luo X, Villablanca EJ. Type 2 immunity in intestinal homeostasis and inflammatory bowel disease. *Biochem Soc Trans* 1 November 2021; 49 (5): 2371–2380. doi: <https://doi.org/10.1042/BST20210535>
81. Kabat AM, Pearce EL, Pearce EJ. Metabolism in type 2 immune responses. *Immunity*, 2023; 56(4):723-41 <https://doi.org/10.1016/j.immuni.2023.03.007>
82. Fujisaka S, Usui I, Nawaz A, Takikawa A, Kado T, Igarashi Y, Tobe K. M2 macrophages in metabolism. *Diabetol Int*. 2016 Nov 1;7(4):342-351. <https://doi.org/10.1007/s13340-016-0290-y>.
83. Hosseini A, Germic N, Markov N, Stojkov D, Oberson K, Yousefi S, Simon HU. The regulatory role of eosinophils in adipose tissue depends on autophagy. *Front Immunol*. 2024 Jan 3;14:1331151. <https://doi.org/10.3389/fimmu.2023.1331151>.
84. Misawa T, Wagner M, Koyasu S. ILC2s and Adipose Tissue Homeostasis: Progress to Date and the Road Ahead. *Front Immunol*. 2022 Jun 16;13:876029. <https://doi.org/10.3389/fimmu.2022.876029>.
85. Kominsky DJ, Campbell EL, Colgan SP. Metabolic Shifts in Immunity and Inflammation. *J Immunol*. 2010; 184 (8): 4062–4068. <https://doi.org/10.4049/jimmunol.0903002>
86. Sun (孙李哲) L, Yang (杨晓峰) X, Yuan (袁祖贻) Z, Wang (王虹) H. Metabolic Reprogramming in Immune Response and Tissue Inflammation. *Arterioscler Thromb Vasc Biol*. 2020 Sep;40(9):1990-2001. <https://doi.org/10.1161/ATVBAHA.120.314037>.
87. Assis PA, Allen RM, Schaller MA, Kunkel SL, Bermick JR. Metabolic reprogramming and dysregulated IL-17 production impairs CD4 T cell function post sepsis. *iScience*, 2024; 27(7):110114. <https://doi.org/10.1016/j.isci.2024.110114>
88. Pålsson-McDermott EM, O'Neill LAJ. Targeting immunometabolism as an anti-inflammatory strategy. *Cell Res*, 2020; 30: 300–314. <https://doi.org/10.1038/s41422-020-0291-z>
89. Gazzinelli-Guimaraes PH and Nutman TB. Helminth parasites and immune regulation. *F1000Research* 2018, 7(F1000 Faculty Rev):1685 <https://doi.org/10.12688/f1000research.15596.1>
90. Kokova D, Mayboroda OA. Twenty Years on: Metabolomics in Helminth Research. *Trends Parasitol*. 2019 Apr;35(4):282-288. <https://doi.org/10.1016/j.pt.2019.01.012>.
91. Wakelin D. Helminths: Pathogenesis and Defenses. In: Baron S, editor. *Medical Microbiology*. 4th edition. Galveston (TX): University of Texas Medical Branch at Galveston; 1996. Chapter 87. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK8191/>
92. Komuniecki R, Tielens A. Carbohydrate and Energy Metabolism in Helminths. *Molecular Medical Parasitology*.2002. <https://doi.org/10.1016/B978-012473346-6/50017-X>.
93. Wangchuk, P, Yeshi K, Loukas A. Metabolomics and lipidomics studies of parasitic helminths: molecular diversity and identification levels achieved by using different characterisation tools. *Metabolomics*, 2023; 19: 63. <https://doi.org/10.1007/s11306-023-02019-5>