

## Assessment of Eco-friendly management of *Alternaria* leaf spot of brinjal (*Solanum melongena* L.)

### ABSTRACT

Among vegetables, brinjal or eggplant is an economically important vegetable crop and second major vegetable crop next to potato in India. This crop suffers from various diseases, among different fungal diseases of brinjal, *Alternaria* leaf spot caused by *Alternaria alternata* has currently become one of the most important diseases for all brinjal varieties. Loss due to the disease has been estimated up to 25-30 per cent. Therefore, present investigation was undertaken at Central Research Field of Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj during the *Kharif* of 2023-2024, for the management of leaf spot using botanicals, bio-agent, cow urine and a fungicide (treated check) for their efficacy *in vivo* against *Alternaria* leaf spot of brinjal incited by *Alternaria alternata*.

Effects of different botanicals, bio-agent and cow urine were evaluated on different growth, disease management and yield parameters of brinjal crop under field condition by foliar spray of botanicals, cow urine and a bio-agent, along with seedling treatment with *Trichoderma viride* at selected concentration. The effects of treatments on growth parameter were recorded i.e. maximum plant height at 30 DAT (22.78 cm), 60 DAT (37.16 cm), 90 DAT (53.87 cm) and maximum number of leaves at 30 DAT (10.83), 60 DAT (31.80), 90 DAT (47.33) and maximum number of branches at 30 DAT (4.23), 60 DAT (11.40), 90 DAT (13.83). Among the treatments, neem leaf extract + *Trichoderma viride* was recorded significantly reducing disease intensity at 45 DAT (10.32%), 60 DAT (16.11%), 75 DAT (27.93%) of brinjal followed by clove extract + *Trichoderma viride* and *Trichoderma viride* + *T. viride* as compared to mancozeb (treated check) and control. The maximum yield was found to be in neem leaf extract + *Trichoderma viride* i.e. (16.63 t/ha) followed by clove extract + *Trichoderma viride*, and *Trichoderma viride* + *T. viride* as compared to mancozeb (treated check) and control and the most economical treatment was Neem leaf extract + *Trichoderma viride* i.e. (1:2.62).

**Keywords:** *Alternaria alternata*, botanicals, brinjal, clove extract, cow urine, fungicide, neem leaf extract, *Trichoderma viride*.

## 1. Introduction

“Brinjal (*Solanum melongena* L.) belongs to the family Solanaceae (USDA, 2008) is an important and indigenous vegetable crop of India. It is a very good source of protein, dietary

fiber and highly productive and usually finds a place as Poor man's crop" (**Rajan and Markose, 2002**).

Brinjal or Eggplant is an important vegetable crop grown in various parts of the world. It is native to northern India and is grown extensively in China and South-East Asia. It is known as Eggplant in English and Baingan in Hindi and synonymously as Aubergine (French name).

"The main brinjal producing countries are China, Egypt, Indonesia, Japan, Turkey, France and Italy. In India, brinjal is grown in practically all parts of the country except the higher altitudes. However, Maharashtra, Gujarat, Punjab, Madhya Pradesh, Kerala, Bihar and Uttar Pradesh are the major producing states. Total production of Brinjal is about 58.6 million of metric tonnes in the world wherein, India is world's second largest producer after China (37.4 million metric tonnes) (**FAO, 2022**). India's share is a cultivated area of 752.80 thousand hectares with an annual production of 13023.20 thousand MT and productivity of 17.30 MT per hectare wherein, Uttar Pradesh covered the cultivated area of 8.82 thousand hectares with an annual production of 312.98 thousand MT and productivity of 35.47 MT per hectare" (**National Horticulture Board, 2022/ Indiatat, 2022**). The productivity of brinjal is quite low (17.30 MT/ha) in India compared to those countries, where the productivity ranges from 17.8 - 37.4 MT/ha. Insect pests and diseases are responsible for the low productivity of brinjal.

"Eggplant due to longer growth period more exposed to number of diseases causing pathogens like nematode, bacteria and fungi. Bacterial wilt, Fusarium wilt, Alternaria leaf spot, damping-off, Phytophthora blight, fruit rot and little leaf of brinjal are most common diseases of eggplants" (**Taher et al., 2017**).

"In global agriculture, about 18% yield losses are caused by animals, 34% by weeds and 16% by microbial diseases. Among microbial diseases, 70-80% of losses are caused by fungal pathogens" (**Moore et al., 2011**). "Fungal species are found all over the world and cause 20-80% yield loss in agriculture by attacking the fruit flower and stem" (**Bashir et al., 2014**).

"Among all the fungal diseases, Alternaria leaf spot, Alternaria leaf blight and fruit rot diseases are of regular occurrence in moderate to severe proportion in India and causes extensive damage to the quality of fruits" (**Pandey and Vishwakarma, 1999**). Alternaria leaf spot caused by *Alternaria alternata* (first reported by (**Fries**) **Keisslerin 1912**), that severely damages the plants of brinjal. *Alternaria* spp. causes up to 25% yield loss in brinjal by

affecting fruits, leaves and flowers in Jaipur district (**Raina et al., 2018**).“In Uttar Pradesh, this disease causes up to 29% yield loss in brinjal by affecting fruit, leaf and flower”(National Horticulture Board, 2022).

“*Alternaria alternata* a fungal pathogen responsible for the development of small, circular, concentric, brownish dark necrotic spots that spread and cause leaf senescence”(El-Gali, 2015).“The necrotic spots enlarge and sporulate under favourable conditions, damaging the leaves and then spreading through the fruit, causing fruit decay” (Batta, 2003).

“Biological control of plant pathogens has been considered as a potential disease management strategy in recent years and appears to be the most promising in disease management. Biocontrol agents colonise the rhizosphere and provide protection against various soil-borne plant pathogens”(Klopper et al., 1989). Fungi of the genus *Trichoderma* are of increasing interest as biocontrol agents.

“Plant extracts have significant and economic applications in the management of plant diseases”(Obongoya et al., 2010). “Chemical fungicides used against plant fungal diseases are toxic to the environment and harmful to human health, hence the need to identify antifungal resources. Plant extracts have natural antifungal bioactive compounds such as saponins, alkaloids, flavonoids and tannins. Plant extracts can be used directly or as a precursor for the development of better molecules”(Arif et al., 2009).“Natural plant products are important sources of new agrochemicals for managing plant diseases”(Kagale et al., 2004).

“The use of cow urine is a cost effective and ecofriendly approach to control phytopathogenic fungi. The antifungal property of cow urine can be explained by the presence of non-volatile active constituents like phenolic acids which are found abundant in the chloroform fraction of cow urine when subjected to high performance liquid chromatographic (HPLC) analysis, the antioxidant property and the ability to reduce germination of spores”(Kekuda et al., 2008; Kekuda et al., 2010; Vats and Miglani, 2011).“Cow urine can be incorporated to minimize fungal infections. Future research is needed to clarify the role of actual lead compounds present in cow urine against fungal pathogens”. (Emehet et al., 2020).“The aqueous cow urine could be exploited for the management of *Alternaria* leaf spot diseases by foliar application”. (Sharma et al., 2023).

Bio-agent and botanicals belonging to various groups are recommended for the management of *Alternaria* leaf spot of brinjal. Generally, farmers are using only the chemicals for managing the disease, but it has negative impact on the environment and develop resistance in the pathogen.

Considering the effects of this disease, the present paper discusses the efficacy of *Trichoderma* sp., cow urine and botanicals for the management of *Alternaria* leaf spot of brinjal.

## **2. Materials and Methods**

The study was conducted field condition at department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, during the *Kharif* season of 2023-24. Field experiment was laid-out in Randomized block design with three replications.

### **2.1 Field preparation**

The selected field area was well prepared and plot marked as per the layout plan. The selected field was ploughed, cleaned and the soil was well pulverized after which the total area was divided into sub-plots.

### **2.2 Experimental site**

The present study was carried out in the Central Research Field under the Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during *Kharif*2023-24. The field experiment was laid-out in Randomized Block Design (RBD) with three replications of eight treatments, each in 2×1 m sized plots and 60×45cm plant spacing was used.

### **2.3 Seedling treatment**

Before transplanting, the seedling was treated by *Trichoderma viride* and mancozeb. For application to seedling treatment of *T. viride*, thick slurry of powder formulation was made at 5 g/litre and seedlings were dipped for 30 minutes in the slurry and for mancozeb (50WP / 0.2 %) applied as seedling treatment as same and then allowed to air dry in shade for 15 minutes, followed by transplanting in designated treatments plots.

### **2.4 Preparation of plant extracts**

The plants were used for the experiment are neem (*Azadirachta indica*), clove (*Syzygium aromaticum*), turmeric (*Curcuma longa*), aloe vera (*Aloe barbadensis*). Plant part used for treatment were neem leaf at 5%, clove at 5%, turmeric rhizome at 5%, aloe vera leaves at 5% to prepare extract. All these plant extracts prepared with the help of mortar and pestle with equal amount of distilled water (1:1 wt/vol). The extracts then filter through muslin cloth to get the stock solution of 100% concentration help of mortar and pestle with the equal amount of distilled water (1:1 wt/vol). The extracts then filtered through muslin cloth to get the stock solution of 100% concentration and further it was diluted according to the 5% concentration in sterile water for *in-vivo* evaluation (Mahapatra and Das, 2013).

## 2.5 Collection of cow urine

For the evaluation of antifungal activity of cow urine, the fresh cow urine of morning period is collected in sterilized container i.e. 100% concentration for stock solution of cow urine further it was diluted according to the 15% concentration in sterile water for *in-vivo* evaluation.

## 2.6 Symptoms

Symptoms first appear as small, isolated, scattered, necrotic brown spots on the leaves. In these necrotic spots, concentric rings appear on the older leaves and darkened areas on the stem. There is usually a narrow chlorotic zone around the spots, which fades into normal green and enlarges in the size of the leaf. This is due to the toxin-alternaric acid, produced by the fungus and translocated through the veins.



**Plate 1: Symptoms on leaf of brinjal**


## **2.7 Isolation of fungal organism**

Initially the diseased leaves were collected from infected plants and thoroughly washed under running tap water. The symptomatic leaf tissue or diseased portion of the leaves were cut under aseptic conditions into small bits and surface sterilized with 0.1% mercuric chloride solution for 1 minute and was washed three times with sterile distilled water and placed on media. To avoid bacterial contamination streptomycin at 100 ppm, was added in the medium at lukewarm stage before pouring PDA into Petri plates. Then Petri plates were wrapped and incubated at  $27\pm 2^{\circ}\text{C}$  in BOD. To get the pure culture of the fungus, hyphal tip method was used for sub-culturing the fungus in media slants/petri-plates. The culture was periodically transferred to fresh media.

## **2.8 Identification of pathogen**

Examination of the fungal colony characteristics was done through microscopic examination. Using a sterile needle, a small portion of the culture was taken and placed on a sterile glass slide. It was stained using lactophenol and cotton blue. Then, the microscope was used for the examination of morphology and culture characteristics of fungal structures.

After 6 days of incubation at  $25^{\circ}\text{C}$  with a 12 h photoperiod, fungal colonies had round margins, and the cottony mycelia were dark olivaceous with a mean diameter of 7.5 cm. Conidiophores were septate and light to olive golden brown with a conidial scar, from which conidia were produced. Conidia were borne singly or in short chains and were obpyriform to obclavate with zero to three longitudinal and two to six transversal septa (**Fr.) Keissl. (1912).**



**Plate 2: Microscopic view of conidia of *Alternaria alternata* on 40x**

### **2.9 Purification, maintenance and preservation of culture**

The culture of *Alternaria alternata* was purified by hyphal tip method and maintained by periodic sub-culturing on PDA petri plates and slants. The pathogen was sub cultured regular intervals of 1 month to maintain the live culture. These were incubated at  $25\pm 2^{\circ}\text{C}$  temperature.



**Plate 3: Culture plate of *Alternaria alternata* Plate 4: Pure Culture tubes of *Alternaria alternata***

## 2.10 Application of treatments

Four plant extracts were taken for treatments along with cow urine, bio-agent, fungicide (treated check) and control. The treatments namely (Clove extract + *Trichoderma viride*), (Turmeric rhizome extract + *T. viride*), (Aloe vera leaf extract + *T. viride*), (Neem leaf extract + *T. viride*), (Cow urine + *T. viride*), (*Trichoderma viride* + *T. viride*), (Mancozeb) and a control as T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>, T<sub>7</sub>, T<sub>0</sub> respectively. All the four plant extracts applied as foliar spray at 5 per cent concentration and bio-agent i.e. *Trichoderma viride* applied as seedling treatment at 5g/litre. In T<sub>5</sub>, cow urine applied as foliar spray at 15 per cent concentration along with seedling treatment of *T. viride* at 5g/kg of seedling. In the treatment T<sub>6</sub> (*Trichoderma viride*) applied as foliar spray at 5 per cent concentration along with seedling treatment of at 5g/litre. Mancozeb (50WP / 0.2 %) applied as foliar spray and seedling treatment as same. For application to seedling treatment of *T. viride*, thick slurry of powder formulation was made at 5 g/litre and seedlings were dipped for 30 minutes in the slurry and then allowed to air dry in shade for 15 minutes, followed by transplanting in designated treatment plots.

## 2.11 Observations recorded

Plant height (cm): Plant height was recorded at 30, 60 and 90 DAT.

No. of leaves: No. of leaves was recorded at 30, 60 and 90 DAT.

No. of branches: No. of branches was recorded at 30, 60 and 90 DAT.

Disease intensity: Disease intensity was recorded as grades in five randomly selected plants in each plot at 45 DAT, 60 DAT and 75 DAT at different times i.e. 15 days after the first spray, 15 days after the second spray and 15 days after the third spray as per the scale of **Mayee and Datar (1986)**.

**Disease intensity (%) was calculated by using the following formula (Wheeler, 1969)**

$$\text{PDI} = \frac{\text{Total sum of rating}}{\text{No. of leaves examined} \times \text{Maximum rating}} \times 100$$

Yield (tonnes/ha): Yield is calculated after the harvesting of the crop.

## 2.12 Cost benefit ratio

Gross return was calculated by multiplying total yield with the market price of the produce. Cost of cultivation and cost of treatment imposition will be deducted from the gross returns, to find out net returns and cost benefit ratio by following formula (Reddy and Reddi, 2004).

$$\text{C:B ratio} = \frac{\text{Gross Return}}{\text{Total cost of cultivation}}$$

Where,

C: B = Cost Benefit ratio

### 3. Results and Discussion

#### 3.1 Effect of treatments on plant growth parameters

##### 3.1.1 Effect of treatments on plant height at 30 DAT, 60 DAT and 90 DAT

Data presented in Table 2 and depicted in figure 1 revealed that the plant height of brinjal significantly increased at 30 DAT, 60 DAT and 90 DAT in the treatment T<sub>4</sub> neem leaf extract + *T. viride* i.e. (22.78 cm, 37.16 cm, 53.87 cm respectively) which was superior over all treatments, followed by T<sub>1</sub> clove extract + *T. viride* i.e. (21.25 cm, 36.19 cm, 52.84 cm respectively), T<sub>6</sub> *Trichoderma viride* + *T. viride* i.e. (18.37 cm, 35.18 cm, 51.59 cm respectively), T<sub>2</sub> turmeric rhizome extract + *T. viride* i.e. (18.30 cm, 33.21 cm, 50.11 cm respectively), T<sub>5</sub> cow urine + *T. viride* i.e. (18.22 cm, 34.26 cm, 50.60 cm respectively), T<sub>3</sub> aloe vera leaf extract + *T. viride* i.e. (18.10 cm, 31.92 cm, 48.63 cm respectively) in comparison of T<sub>0</sub> control (16.16 cm, 27.96 cm, 45.77 cm respectively). However, all the treatments were significant over control.

##### 3.1.2 Effect of treatments on plant leaves at 30 DAT, 60 DAT and 90 DAT

Data presented in Table 3 and depicted in figure 2 revealed that the number of leaves of brinjal significantly increased at 30 DAT, 60 DAT and 90 DAT in the treatment T<sub>4</sub> neem leaf extract + *T. viride* i.e. (10.83, 31.80, 47.33 respectively) which was superior over all the treatments, followed by T<sub>1</sub> clove extract + *T. viride* i.e. (10.63, 31.36, 46.20 respectively), *Trichoderma viride* + *T. viride* i.e. (9.26, 30.50, 45.10 respectively), T<sub>5</sub> cow urine + *T. viride* i.e. (8.26, 29.73, 44.40 respectively), T<sub>2</sub> turmeric rhizome extract + *T. viride* i.e. (8.23, 28.20, 43.23 respectively), T<sub>3</sub> aloe vera leaf extract + *T. viride* i.e. (8.06, 28.20, 43.23 respectively)

in comparison of T<sub>0</sub> control (6.73, 24.93, 40.20 respectively). All the treatments were significant over control.

### 3.1.3 Effect of treatments on plant branches at 30 DAT, 60 DAT and 90 DAT

Data presented in Table 4 and depicted in figure 3 revealed that the number of branches of brinjal significantly increased at 30 DAT, 60 DAT and 90 DAT in the treatment T<sub>4</sub> neem leaf extract + *T. viride* i.e. (4.23, 11.40, 13.83 respectively) which was superior over all the treatments, followed by T<sub>1</sub> clove extract + *T. viride* i.e. (3.80, 10.36, 13.46 respectively), T<sub>6</sub> *Trichoderma viride* + *T. viride* i.e. (3.76, 9.46, 12.23 respectively), T<sub>5</sub> cow urine + *T. viride* i.e. (3.63, 8.36, 11.16 respectively), T<sub>2</sub> turmeric rhizome extract + *T. viride* i.e. (3.56, 7.36, 10.14 respectively), T<sub>3</sub> aloe vera leaf extract + *T. viride* i.e. (3.43, 7.06, 10.13 respectively) in comparison of T<sub>0</sub> control (2.03, 5.26, 8.16 respectively). All the treatments were significant over control.

In the present studies, plant height, number of leaves, number of branches at 30, 60 and 90 DAT was recorded in neem leaf extract + *Trichoderma viride* followed by clove extract + *T. viride* was found to be effective over other treatments. This is because all growth parameters increased in comparison to control this may be due to reduction in insect pests or the percentage infestation by the pests or both. The growth stimulating effect is not exclusively by its adverse effect on pathogen or by an increase in nutrient uptake. However, substances with hormone-like properties can stimulate of effect biomass allocation in plants. In addition to hormone, medicinal plant extracts contain saponins and polyphenols which could be the active compounds causing the effect on growth and yield of the plants. By seedling treatment with *Trichoderma viride* it gives effect by enhancing physiological responses to stress, boost nitrogen-use efficiency in certain crop, and enhance photosynthetic efficiency. So, combination of neem leaf extract (foliar spray) and seedling treatment with *Trichoderma viride* gives effective results on the growth parameters of brinjal plant. These results were similar to the findings of Netam *et al.* (2014) in that the highest dry shoot weight was found in *T. viride* and was effective over other treatments and similar finding of Kakraliya *et al.* (2017) were *in-vitro* and field conditions, the effect of bio-agents and botanicals showed maximum plant height i.e. in *T. viride* followed by *T. harzianum* as compared to control.

Nahak and Sahu (2014) also tested “the aqueous neem extract for the growth, yield and disease control of a common vegetable plant brinjal and found that the neem extracts increased shoot height, number of leaves, number of buds, number of flowers and number of

fruits of brinjal plant over control. Disease incidence was calculated in percentage. The percentage of reduction was calculated after the spray of neem extract. Neem extract was found as most effective agent in controlling leaf spot (82.33%) in comparison to control under field conditions. The results showed that neem leaf extract act as potential source for the management of severe diseases in addition to its positive effect on growth parameters”.

These results were also similar to the findings of Reddy *et al.* (2021) in that number of leaves significantly increased in the treatment combination of neem oil + *T. viride*. Netam *et al.* (2014) and Gabal *et al.* (1999) also concluded that the highest dry shoot weight was found in treatment *T. viride* and was effective over other treatment.

These results were also similar findings to Reddy *et al.* (2021) in which the number of branches significantly increased in the treatment combination of Neem and *Trichoderma viride* and in tomato by Mehta *et al.* (1989) and in brinjal by Netam *et al.* (2014).

**Table No: 1 Details of treatments on field management of Alternaria leaf spot of brinjal**

S.no	No. of Treatments	Name of Treatments	Concentration
1	T <sub>0</sub>	Control (untreated check)	--
2	T <sub>1</sub>	<i>Trichoderma viride</i> + Clove extract	0.5% (S.T) + 5% (F.S)
3	T <sub>2</sub>	<i>Trichoderma viride</i> + Turmeric rhizome extract	0.5% (S.T) + 5% (F.S)
4	T <sub>3</sub>	<i>Trichoderma viride</i> + Aloe vera leaves extract	0.5% (S.T) + 5% (F.S)
5	T <sub>4</sub>	<i>Trichoderma viride</i> + Neem leaf extract	0.5% (S.T) + 5% (F.S)
6	T <sub>5</sub>	<i>Trichoderma viride</i> + Cow urine	0.5% (S.T) + 15% (F.S)
7	T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	0.5% (S.T) + 5% (F.S)
8	T <sub>7</sub>	Mancozeb	75% WP (0.2%) (S.T + F.S)

\*S.T.- Seedling Treatment;


\*F.S.- Foliar Spray

**Table No: 2 Effect of treatments on plant height (cm) of brinjal at different days interval**

Tr. No.	Treatment	Plant height (cm)*		
		30 DAT	60 DAT	90 DAT
T <sub>0</sub>	Control	16.16	27.96	45.77
T <sub>1</sub>	Clove extract + <i>T. viride</i>	21.25	36.19	52.84
T <sub>2</sub>	Turmeric rhizome extract + <i>T. viride</i>	18.30 <sup>a</sup>	33.21	50.11 <sup>a</sup>
T <sub>3</sub>	Aloe vera leaf extract + <i>T. viride</i>	18.10 <sup>a</sup>	31.92	48.63
T <sub>4</sub>	Neem leaf extract + <i>T. viride</i>	22.78	37.16	53.87
T <sub>5</sub>	Cow urine + <i>T. viride</i>	18.22 <sup>a</sup>	34.26 <sup>a</sup>	50.60 <sup>a</sup>
T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	18.37 <sup>a</sup>	35.18 <sup>a</sup>	51.59
T <sub>7</sub>	Mancozeb	24.36	38.30	54.84
CD (p=0.05)		1.03	0.95	0.89
SEd (±)		0.48	0.44	0.41

\* Average of five plants

• Data followed by same alphabet in a column are non-significant to each other at 5% level.



**Figure- 1: Effect of treatments on plant height (cm) of brinjal at different days interval**


**Table- 3: Effect of treatments on number of leaves in brinjal at different day intervals**

Tr. No.	Treatment	No. of. Leaves*		
		30 DAT	60 DAT	90 DAT

T <sub>0</sub>	Control	6.73	24.93	40.20
T <sub>1</sub>	Clove extract + <i>T. viride</i>	10.63 <sup>a</sup>	31.36 <sup>a</sup>	46.20
T <sub>2</sub>	Turmeric rhizome extract + <i>T. viride</i>	8.23 <sup>b</sup>	28.20	43.23
T <sub>3</sub>	Aloe vera leaf extract + <i>T. viride</i>	8.06 <sup>b</sup>	27.23	42.26
T <sub>4</sub>	Neem leaf extract + <i>T. viride</i>	10.83 <sup>a</sup>	31.80 <sup>a</sup>	47.33
T <sub>5</sub>	Cow urine + <i>T. viride</i>	8.26 <sup>b</sup>	29.73	44.40 <sup>a</sup>
T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	9.26	30.50	45.10 <sup>a</sup>
T <sub>7</sub>	Mancozeb	12.40	32.50	48.26
CD (p=0.05)		0.98	0.46	0.85
SEd (±)		0.46	0.22	0.40

\* Average of five plants

- Data followed by same alphabet in a column are non-significant to each other at 5% level.



**Figure- 2: Number of leaves in brinjal at different days interval**


**Table- 4: Effect of treatments on number of branches in brinjal at different day interval**

Tr. No.	Treatment	No. of. Branches*		
		30 DAT	60 DAT	90 DAT

T <sub>0</sub>	Control	2.03	5.26	8.16
T <sub>1</sub>	Clove extract + <i>T. viride</i>	3.80 <sup>a</sup>	10.36	13.46 <sup>a</sup>
T <sub>2</sub>	Turmeric rhizome extract + <i>T. viride</i>	3.56 <sup>a</sup>	7.36 <sup>a</sup>	10.14 <sup>b</sup>
T <sub>3</sub>	Aloe vera leaf extract + <i>T. viride</i>	3.43 <sup>a</sup>	7.06 <sup>a</sup>	10.13 <sup>b</sup>
T <sub>4</sub>	Neem leaf extract + <i>T. viride</i>	4.23	11.40	13.83 <sup>a</sup>
T <sub>5</sub>	Cow urine + <i>T. viride</i>	3.63 <sup>a</sup>	8.36	11.16
T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	3.76 <sup>a</sup>	9.46	12.23
T <sub>7</sub>	Mancozeb	4.86	12.80	14.90
CD (p=0.05)		0.42	0.69	1.00
SEd (±)		0.20	0.32	0.47

\* Average of five plants

• Data followed by same alphabet in a column are non-significant to each other at 5% level.



**Figure- 3: Effect of treatments on number of branches in brinjal at different days interval**

### 3.2 Effect of treatment on disease intensity (%) at different DAT of Alternaria leaf spot of brinjal

The efficacy studies for the management of Alternaria leaf spot of brinjal. The data recorded in table 5 and depicted in figure 3 revealed that all the treatments were significantly superior

to untreated (control) T<sub>0</sub> in reducing disease severity and percent disease incidence at 45 DAT, 60 DAT and 75 DAT. The minimum disease intensity percent at 45 DAT, 60 DAT and 75 DAT was recorded with T<sub>4</sub> neem leaf extract + *T. viride* (10.32%, 16.11%, 27.93% respectively) followed by T<sub>1</sub> clove extract + *T. viride* (10.44%, 16.92%, 28.27% respectively), T<sub>6</sub> *Trichoderma viride* + *T. viride* (11.51%, 18.05%, 29.84% respectively), T<sub>5</sub> cow urine + *T. viride* (12.66%, 19.45%, 30.98% respectively), T<sub>2</sub> turmeric rhizome extract + *T. viride* (13.05%, 19.93%, 31.35% respectively) and T<sub>3</sub> aloe vera leaf extract + *T. viride* (14.23 %, 20.97%, 32.88% respectively), maximum percent disease incidence was recorded in T<sub>0</sub> control (16.48%, 23.98%, 34.99% respectively).

In the present studies minimum disease incidence at 45, 60 and 75 DAT was recorded with neem leaf extract + *Trichoderma viride* followed by clove extract + *T. viride* and *Trichoderma viride* + *T. viride* at selected concentration was found to be effective over other treatments. The probable reasons may be that Neem leaf extract is highly efficient in controlling leaf spot in *Solanum melongena* L. due to its active compounds such as azadirachtin, nimbin, nimbidin, nimbinene and azadirone which are antifungal, antibacterial and anti-insecticidal in nature. Quantitative estimation of phytochemical compounds of neem leaves showed presence of phenols, flavonoids, tannins, alkaloids and saponins. This class of compounds independently or in combination may be responsible for the broad range of medicinal properties of neem and effective against controlling leaf spot diseases of brinjal plant. So, *Azadirachtaindica* is highly efficient in controlling leaf spot disease in brinjal plant by using aqueous leaf extract to control pathogen and substantially reduced number of infected leaves and capsules from infection and reduction of leaf spot is up to 50%. It proves that the plant extracts of neem can serve as alternatives to the synthetic pesticides for controlling leaf spot of brinjal without any adverse effect on crop and yield parameters.

So, in combination of seedling treatment with *Trichoderma viride* it shows effective results probably due to antagonistic and antifungal activities of these agents. *Trichoderma viride* reduces the severity of plant diseases by inhibiting soil pathogens through its potent antagonistic and mycoparasitic activities. The antagonistic activity of *T. viride* against plant pathogens has been produced secondary metabolites such as harzianic acid, alamethicins, peptaibols, antibiotics, 6- pethyl-  $\alpha$  -pyrone, massoilactone, viridin, glioviridin, gliovirin, glisoprenins, heptelidic acid, pentyl pyrone etc which have antifungal properties. So, by combination of neem leaf extract by using (foliar spray) and *Trichoderma viride* as seedling treatment it gives very effective results without harming environments.

Similar findings were given by Ravishanker *et al.* (2018) that “bio-agents and combinations in *in-vivo* and *in-vitro*, *Trichoderma viride*, and neem oil and their combination the results show the combination of bio-agents had a great impact in controlling the blight disease of *Alternaria* blight followed by fungicide”. Vijayalakshmi *et al.* (2018) reported “maximum inhibition of radial growth of *Alternariahelianthi* was observed in *T. viride* (85.33%) which was followed by *T. viride* strain 16 (79.33%), and *T. harzianum*. Bagwan (2010) results given that, neem oil (5%), neem leaves extract (10%), wild sorghum leaves extract (10%), neem cake, castor cake and mustard cake extract (10%) enhanced the growth of *Trichoderma*”.

Similar findings, in regards of treatment clove extract + *Trichoderma viride*.e preceded by neem leaf extract + *T. viride* is of Walter *et al.* (1997) that “they discovered a special formulation of rectified clove oil and surfactant composed of ionic and anionic compounds was highly effective in inhibiting fungal diseases. The formulation, which forms a stable emulsion when diluted with water, is applied as an aqueous solution. Upon dilution, the clove water formulation is highly effective in application as soil drench, or when incorporated into the soil prior to planting for soil-borne diseases or as foliar spray for foliar diseases”.

Chrapacieneet *al.* (2022) also gave similar findings by conducting assay, in which they revealed that the development of fungal mycelium depended on the concentration of the extract in the medium. His results revealed that clove extract was effective *in vitro* and concluded, *Syzygiumaromaticum* extract has promising antifungal properties, causing *Alternaria* spp. growth suppression.


**Table- 5: Effect of treatments on disease intensity (%) in brinjal at different day interval**

Tr. No.	Treatment	Disease intensity (%)		
		45 DAT	60 DAT	75 DAT

T <sub>0</sub>	Control	16.48	23.98	34.99
T <sub>1</sub>	Clove extract + <i>T. viride</i>	10.44 <sup>b</sup>	16.92	28.27 <sup>b</sup>
T <sub>2</sub>	Turmeric rhizome extract + <i>T. viride</i>	13.05 <sup>a</sup>	19.93 <sup>a</sup>	31.35 <sup>a</sup>
T <sub>3</sub>	Aloe vera leaf extract + <i>T. viride</i>	14.23	20.97	32.88
T <sub>4</sub>	Neem leaf extract + <i>T. viride</i>	10.32 <sup>b</sup>	16.11	27.93 <sup>b</sup>
T <sub>5</sub>	Cow urine + <i>T. viride</i>	12.66 <sup>a</sup>	19.45 <sup>a</sup>	30.98 <sup>a</sup>
T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	11.51	18.05	29.84
T <sub>7</sub>	Mancozeb	9.34	14.95	26.87
CD (p=0.05)		0.91	0.59	0.85
SEd (±)		0.43	0.28	0.40

\* Average of five plants

- Data followed by same alphabet in a column are non-significant to each other at 5% level.



**Figure- 4: Effect of treatments on disease intensity (%) in brinjal at different day interval**

### 3.3 Effect of treatments on yield and C:B ratio

Data presented in Table 5 and figure 4 showed, among all the treatments, maximum fruit yield of brinjal (t/ha) was recorded in treatment T<sub>4</sub> neem leaf extract + *T. viride* i.e. (16.63)


which was superior over all treatments, followed by T<sub>1</sub> clove extract + *T.viride*.e. (15.60) as comparison of T<sub>0</sub> (9.23).

The treatment wise economics of brinjal productions under the field demonstrations were estimated and the results have been in Table 6. The economics analysis of the data over the session that T<sub>4</sub> neem leaf extract + *T. viride* recorded higher gross returns i.e. Rs.2,16,190.00, net returns Rs. 1,33,720.00 with C:B ratio 1:2.62 followed by T<sub>6</sub> *Trichodermaviride*+ *T. viride* gross return Rs. 1,88,500.00, net returns Rs. 1,05,055.00 with C:B ratio 1:2.25 and T<sub>5</sub>cow urine + *T. viride* gross return Rs. 1,78,971.00, net returns Rs. 96,156.00 with C:B ratio 1:2.16.

**Table- 6: Effect of treatments on yield and C:B ratio**

Tr. No	Treatment	Fruit yield (t/ha)	Total cost of cultivation (Rs/ha)	Gross return (Rs/ha)	C:B ratio
T <sub>0</sub>	Control	9.23	79,045	1,19,990	1:1.51
T <sub>1</sub>	Clove extract + <i>T. viride</i>	15.60	110,820	2,02,800	1:1.82
T <sub>2</sub>	Turmeric rhizome extract + <i>T. viride</i>	13.46 <sup>a</sup>	89,070	1,75,071	1:1.96
T <sub>3</sub>	Aloe vera leaf extract + <i>T. viride</i>	12.40	83,820	1,61,200	1:1.92
T <sub>4</sub>	Neem leaf extract + <i>T. viride</i>	16.63	82,470	2,16,190	1:2.62
T <sub>5</sub>	Cow urine + <i>T. viride</i>	13.76 <sup>a</sup>	82,815	1,78,971	1:2.16
T <sub>6</sub>	<i>Trichoderma viride</i> + <i>T. viride</i>	14.50	83,445	1,88,500	1:2.25
T <sub>7</sub>	Mancozeb	17.60	82,185	2,28,800	1:2.76

- Data followed by same alphabet in a column are non-significant to each other at 5% level.



**Figure 5: Effect of treatments on fruit yield (t/ha) of brinjal**

#### 4. Conclusions

Based on the results obtained from present investigations it was found that foliar spray of neem leaf extract + *T. viride* (seedling treatment) was most effective against Alternaria leaf spot of brinjal which causes leaf spot disease in brinjal followed by foliar spray of clove extract + *T. viride*(seedling treatment)and foliar spray of *Trichoderma viride* + *T. viride*(seedling treatment), therefore it may be recommended for the better management of Alternaria leaf spot of brinjal, in this the input cost of clove extract + *T. viride* is little bit high but very effective treatment preceded by neem leaf extract + *T. viride* for Alternaria leaf spot of brinjal however, the all treatment is used in this research work is effective except aloe vera leaf extract + *T. viride*.i.e less effective as compared to above mentioned treatments. It is important for us to adapt these alternative management approaches as increased use of agrochemicals have created serious problems for existing and upcoming generations.

Foliar spray of neem leaf extract + *T.viride*(seedling treatment) under field condition resulted in minimum disease intensity (%) at 75 DAT i.e. 27.93, on the basis of growth parameter before final harvesting the plant height is (53.87 cm), number of leaves (47.33), number of branches (13.83) and the yield i.e. 16.63 t/ha.

Results of the present study were found to be significantly effective under Prayag raj agroclimatic conditions. It may vary with region and climatic conditions, therefore for validation of the results mores such trials should be carried out in future.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## References

- Arif, T., Bhosale, J. D., Kumar, N., Mandal, T. K., Bendre, R. S.,Lavekar, G. S. and Dabur, R. (2009).** Natural products– antifungal agents derived from plants. *Journal of Asian Natural Products Research*. 11 (7): 621–638.
- Bashir, U., Mushtaq, S. and Akhtar, N. (2014).** First report of *Alternaria metachromatica* from Pakistan causing leaf spot of tomato. *Pakistan Journal of Agricultural Sciences*.51: 305-308.

- Chrapaciene, S., Rasiukeviciute, N., Rudinskaite, A. and Valiuskaite, A. (2022).** Clove extract for carrot *Alternaria* spp. management. *LAMMC Institute of Horticulture*. 592-593.
- El-Gali, Z. I. (2015).** *Trichoderma harzianum* as a root treatment to control *Alternaria alternata* ceratoni leaf spot on *Ceratonia siliqua* L. *Sky Journal of Agricultural Research*. 4:161-166.
- Emeh, A. A., Anyanwu, G. O., Amadi, R. E., Onyewenjo, S. C. and Chukwudifu, A. J. (2020).** Investigation on anti-fungal properties of cow urine on pathogenic fungi. *IOSR Journal of Pharmacy (IOSRPHR)*. 10(6): 20-24.
- Food and Agriculture organization. (2022).** <https://www.fao.org/>.
- Gabal, Oben G. M. and Garell, R. (1999).** Effect of G.A on morphophysiological characters and yield of kidney beans (*Phaseolus vulgaris*). *Journal of Agronomy and Crop Science*. 160(2): 91-101.
- India-stat. Indiastat.com. (2022).** Area, production and productivity of Brinjal in India (1987-1988, 1991-1992 and 1993-1994 to 2022-2023(2nd advance estimate) (indiastat.com).
- Kagale, S., Marimuthu, T., Thayumanavan, B., Nandakumar, R. and Samiyappan, R. (2004).** Antimicrobial activity and induction of systemic resistance in rice by leaf extract of *Datura metel* against *Rhizoctonia solani* and *Xanthomonas oryzae* pv. *oryzae*. *Physiological and Molecular Plant Pathology*. 65:91-100.
- Kakraliya., Dechan, C., Devanshi, P. and Sonali, A. (2017).** Effect of bio-agents, neem leaf extract and fungicides against *Alternaria* leaf blight of wheat (*Triticum aestivum* L.) *Natural Products Chemistry and Research*. ISSN. 2329-6836.
- Keissler, K. (1912).** Zur Kenntnis der Pilzflora Krains. *Beihefte zum Botanischen Centralblatt*. 29:395-440.
- Kekuda, P. T. R., Kavya, R., Shrungashree, R. M. and Suchitra, S. V. (2008).** *In vitro* studies on antifungal activity of cow urine distillate against fungi causing opportunistic infections. *BioTechnology: An Indian Journal*. 2(1): 47-51.
- Kekuda, P. T. R., Nishanth, B. C., Praveen, S. K., Kamal, D., Sandeep, M. and Megharaj, H. K. (2010).** Cow urine concentrate: A potent agent with antimicrobial and anthelmintic activity. *Journal of Pharmacy Research*. 3(5): 1025-1027.
- Kloepper, J.W., Lifshitz, R and Zablutowicz, R.M. (1989).** Free-living bacterial inocula for enhancing crop productivity. *Trends biotechnology*. 7:39-43.

- Mahapatra, S. and Das, S. (2013).** Bio-efficacy of botanicals against *Alternaria* leaf blight of mustard under field condition. *The Bioscan*. 8(2): 675-679.
- Mayee, C. D. and Datar, V. V. (1986).** Technical Bulletin-1 (Special Bulletin-3). *Phytopathometry*. Marathwada Agricultural University, Parbhani, Maharashtra, India. 95.
- Mehta, A. K., Singh, R. P. and Lal, G. (1989).** Effect of concentration and methods of application of 2,4-D phenoxy acetic acid on yield, fruit quality and seed quality of tomato (*Lycopersicon esculentum* Mill). *Vegetable Science*. 16 (1): 1-8.
- Moore, D., Robson, G. D. and Trinci, A.P. (2011).** 21st century guidebook to fungi with CD. Cambridge University Press. 368-370.
- Nahak, G. and Sahu, R. K. (2014).** Bio-efficacy of leaf extract of neem (*Azadirachta indica* A. Juss) on growth parameters, wilt and leafspot diseases of brinjal. *Research Journal of Medicinal Plant*. 8: 269-276.
- National Horticulture Board. (2022).** Ministry of Agriculture and Farmers Welfare. Government of India.
- Netam, J. L. and Sharma, R. (2014).** Efficacy of plant growth regulators on growth characters and yield attributes in brinjal (*Solanum melongena* L.) cv. Brinjal 3112. *IOSR Journal of Agriculture and Veterinary Science*. 7:27-30.
- Obongoya, B. O., Wagai, S. O. and Odhiambo, G. (2010).** Phytotoxic effect of selected crude plant extracts on soil-borne fungi of common bean. *African Crop Science Journal*. 18:15–22.
- Pandey, K. K. and Vishwakarma, S. N. (1999).** Morphological and symptomatological variations in *Alternaria alternata* causing leaf blight in brinjal. *Journal of Mycology and Plant Pathology*. 29: 350-353.
- Raina, Z. K., Rather, R. A., Bhat, S. A., Wani, T. A., Bhat, N. A. and Dorjey, S. (2018).** Severity of *Alternaria* leaf spot of brinjal caused by *Alternaria alternata* in Kashmir, India. *International Journal of Current Microbiology and Applied Sciences*. 7: 322-328.
- Rajan, S. and Markose, B. L. (2002).** Propagation of horticultural crops. *Horticulture Science Series*. 6: 94.
- Ravishankar and Shashi, T. (2018).** Biological management of *Alternaria* leaf blight in coriander (*Coriandrum Sativum*). *Journal of Pharmacognosy and Phytochemistry*. 7(6): 1867-1869.

- Reddy, A. K., Tiwari, S. and Rohini, T. (2021).** Eco-friendly management of leaf spot of brinjal (*Solanum melongena* L.) caused by *Alternaria* spp. *International Journal of Current Microbiology and Applied Sciences*. 10(05): 132-142.
- Reddy, T. Y. and Reddi, G. H. S. (2004).** Principles of Agronomy 3rd edition, Kalyanipublisher, *Plant Pathology*. 527.
- Sharma, S., Sharma, A. and Sharma, M. (2023).** Antifungal effect of aqueous and cow urine extracts of native plants against *Alternariabrassicola* causing leaf spot of cauliflowers. *Plant Diseases Research*. 38(2): 132-139.
- Taher, D., Solberg, S. O., Prohens, J., Chou, Y. Y., Rakha, M. and Wu, T. H. (2017).** World vegetable center eggplant collection: origin, composition, seed dissemination and utilization in breeding. *Frontiers in Plant Science*. 8:1-12.
- United States Department of Agriculture. (2008).** National Agricultural Statistics Service. <https://www.nass.usda.gov>.
- Vats, S. and Miglani, K. (2011).** Synergistic antimicrobial effect of cow urine and *Azadirachtaindica* on infectious microbes. *International Journal of Pharmaceutical Sciences and Research*. 2(7): 1781–1785.
- Vijayalakshmi, G., Karuna, K. and Mahadevaswamy, G. (2018).** Evaluation of microbial biocontrol agents and fungicides against *Alternaria helianthi* causing leaf blight of sunflower. *International Journal of Current Microbiology and Applied Sciences*. 7(01): 2726-2730.
- Walter, J. F., Locke, J. C. and Normoyle, M. C. (1997).** Clove oil as Plant Fungicide. United States Patent. Patent Number: 5,679,351.
- Wheeler, B. E. J. (1969).** *An Introduction to Plant Disease*. John Wiley and Sons Limited, London. 301.