

Original Research Article

Insecticidal and Anti-feedant activity of *Clerodendrum* extracts on growth and development of Castor Spiny Caterpillar, *Ariadne merione*

Abstract

The castor spiny caterpillar *Ariadne merione* (F.) is a major lepidopteran pest affecting castor (*Ricinus communis*) crops mainly by causing defoliation which in turn reduces the photosynthesis and yield. The conventional pest control methods have been based on chemical control which although has been effective, has its drawbacks of being hazardous to the environment and human health besides leading to resistance by the pests. The present work aims at investigating the possibility of using *Clerodendrum inerme*, a plant belonging to the Lamiaceae family, as an eco-friendly solution for pest management. Insecticidal and anti-feedant efficacy of *C. inerme* leaf extracts, which were prepared using solvents hexane, acetone, and petroleum ether, was tested against *A. merione* larvae using leaf dip bioassay under no-choice situation. The findings showed that the hexane extract had the highest insecticidal and anti-feedant action with a mortality rate of 73 percent. The petroleum ether extract also showed moderate insecticidal and anti-feedant activity but not as potent as the acetone extract. The results imply that *C. inerme* extracts, especially the hexane extract, could be used as botanical insecticides in IPM programs as a natural replacement for chemical insecticides.

Keywords: *Clerodendrum inerme*, *Ariadne merione*, Anti-feedant, Mortality, Integrated Pest Management.

Introduction

Ariadne merione (F.), a lepidopteran pest, is responsible for causing severe harm to the yield and quality of castor (*Ricinus communis*). Its larvae, being an economically significant pest, feed extensively on castor leaves, resulting in defoliation and drop in photosynthetic ability, which ultimately causes a major yield loss. When tilling crops in tropical and subtropical areas, the pest is challenging to manage because of its wide host range and ability to thrive in a variety of environmental circumstances.

Synthetic pesticides are the mainstay of traditional pest management strategies. Although the effectiveness of these compounds in controlling *A. merione* has been demonstrated, their usage is linked to a number of drawbacks, including environmental

contamination, toxicity to organisms that are not the intended targets, and the possibility of encouraging the emergence of resistance in pest species (Navatha *et al.*, 2011). These issues have forced the farming community to search for safer alternatives for pest management (Ahmed *et al.*, 2021).

Another set of research idea can be associated with the administration of plant origin substances known as botanicals, as opposed to synthetic compounds. It is important to note that the insecticidal qualities of the plant species of the current generation that belong to the genus *Clerodendrum inerme* have attracted a lot of attention. *C. inerme*, which are members of the Lamiaceae family and are found throughout the world's tropical and subtropical regions are distinguished by an abundance of bioactive substances. These substances have been used in numerous medicinal procedures in the past, and recent studies have shown that they may also be effective natural pesticides.

Various alkaloids, flavonoids, terpenoids and phenolic acids isolated from the extracts of *C. inerme* genus demonstrated the general biological activities such as insecticidal, anti-feedant and repellent activities (Kar *et al.*, 2014) These properties make *C. inerme* extracts a viable additive to Integrated Pest Management (IPM) regimes designed to mitigate dependence on synthetic insecticides.

In the current study, we aim to evaluate the insecticidal and anti-feedant properties of *C. inerme* extracts against the castor spiny caterpillar (*Ariadne merione*). Our goal is to determine the effectiveness of these extracts as a sustainable pest control method along with their influence on its feeding behaviour. This research seeks to identify potent botanicals that can be integrated into Integrated Pest Management (IPM) strategies, offering farmers an eco-friendly alternative to chemical insecticides. The findings from this study could pave the way for developing new bio-insecticides derived from *C. inerme* species, potentially enhancing agricultural practices while minimizing the environmental impact of pest control methods.

Materials and Methods

Plant material and Preparation of extracts

The fresh leaves of *C. inerme* were harvested from the Botanical garden of PGP College of Agricultural Sciences, Namakkal, Tamil Nadu, India situated in the geographical co-ordinates of 11.21⁰ N Latitude and 78.16⁰ E longitude. The leaves were washed with distilled water several times and then dried under the shade and powdered mechanically. The powdered plant material was sealed in an air-tight container before extraction and kept at ambient temperature.

Preparation of crude extracts

The extraction was carried out using solvents of increasing polarity such as acetone, petroleum ether and hexane. For extraction, 100g of the powdered leaf material was mixed with 500ml of the respective solvent and shaken in a shaker at room temperature for 48 hours. These were followed by filtration through Whatman no 1 filter paper and removal of solvents by evaporation under reduced pressure in a rotary evaporator at 40°C. The dried extracts were then weighed and kept in a closed glass containers in refrigerator at 4°C until analysis.

Leaf dip bioassay

To evaluate the toxicity and anti-feedant activity of *C. inermis* extracts to *A. merione* larvae, leaf dip method under no choice condition was employed. The prepared crude acetone, petroleum ether and hexane extracts were taken and six different concentrations *viz.*, 0.7, 1.0, 3.0, 5.0, 7.0 and 10.0 per cent were prepared using the respective solvents. Each concentration was prepared and added with 0.1 per cent Triton X-100 as a sticking agent so that the extracts stick to the leaves and did not drain out from the leaves. Triton X-100 with respective solvents were kept as control. The castor leaf discs with known size (4 cm diameter) were dipped in different concentrations of acetone, petroleum ether and hexane crude extracts for about 30 seconds. The leaf discs were taken out using forceps and placed in filter paper and shade dried. After drying, the leaves were kept in the Petri dishes and 4 h pre-starved larvae were released into each Petri dish and allowed to feed under no choice conditions. For each concentration, three replicates with ten larvae per replication were used. The experiment was done under Completely Randomized Design. The mortality due to direct toxicity/growth regulatory activity to the larvae were calculated at different time intervals starting with 24 h after treatment up to the treated larva stops feeding (Duraipandiyani *et al.*, 2011). The mortality percentage and the anti feedant index (Jannet *et al.*, 2000) was calculated by using the formula,

$$\text{Mortality (\%)} = \frac{\text{Number of insects dead}}{\text{Total number of insects released}} \times 100$$

$$\text{Antifeedant Index (\%)} = \frac{\text{Leaf area consumed in control} - \text{Leaf area consumed in treated}}{\text{Leaf area consumed in control} + \text{Leaf area consumed in treated}} \times 100$$

All collected data were analysed using analysis of variance (ANOVA) on a Statistical Package for Social Science (SPSS) software. Differences of greater magnitude between the treatments were established using Duncan's Multiple Range Test at 5 % level of significance.

Results and Discussion

1. Extraction yield of *C. inerme* extracts

The highest yield was obtained with hexane ($9.6 \pm 0.22\%$) while petroleum ether yielded $6.2 \pm 0.14\%$ and acetone yielded $2.9 \pm 0.03\%$ (Table 1). The variation in extraction yield of different solvents can be attributed to the polarity of the solvents because the solubility of bioactive compounds in the plant leaves varies with polarity of the solvent used. The yield of hexane is higher than that of the other solvents and this indicates that non polar compounds form a large proportion of the bio active compounds in *C. inerme* leaves.

2. Anti-Feedant activity of *C. inerme* extracts

The antifeedant activity of *C. inerme* extracts was tested with petroleum ether, hexane and acetone as solvents. Out of all the extracts, hexane gave the highest results at 10% concentration with the highest AFI of 87% at all the concentrations followed by Acetone (74%) and Petroleum ether (48%). The similar trend was observed at the lowest concentration (1%) with Hexane (61%), Acetone (64%) and Petroleum ether (48%). The anti-feedant action increases with increasing concentration (Fig. 1). On the other hand, the control group exhibited low AFI suggesting that *C. inerme* extracts are efficient in reducing feeding by *A. merione* larvae with hexane as the best solvent.

3. Toxicity of *C. inerme* extracts against *A. merione*

The results obtained from the experiment on the larval mortality of *A. merione* when treated with *C. inerme* extracts of various concentrations presented in Fig. 2 affirm the insecticidal property of the extracts. Out of the three extracts used, hexane extract was the most toxic followed by acetone extract and petroleum ether. The mortality gradually increased from day 2 and extended to day 5. At 10% concentration of hexane extract showed 73.33% mortality by day 5 as compared to Acetone (56%) and Petroleum ether (50%). This can be attributed to the fact that hexane extract contains higher concentration of non-polar bioactive compounds that are soluble in non-polar solvents and possess good insecticidal properties.

The LC_{10} , LC_{20} and LC_{50} values also support the effectiveness of the hexane extract of the plant. It had the least LC_{50} value (4.963%), this implying that it was the most toxic to *A. merione* larvae at lower concentration than acetone ($LC_{50} = 9.747\%$) and petroleum ether ($LC_{50} = 14.70\%$). The confidence intervals for these LC_{50} values especially the narrow one for the hexane extract supports the claim that it could be a potent insecticide. The mortality rates obtained for *A. merione* larvae decreased with time and with higher concentrations of *C.*

inerme extracts, which indicates that *C. inerme* extracts have both acute and chronic toxic effects on *A. merione* larvae as was found in other plant-based pest control systems.

The findings of this study are evident to show that the extracts of *C. inerme*, especially the hexane and acetone extracts, have potent insecticidal and anti-feedant effects against the *A. merione*. The high mortality rates and strong feeding deterrence indicate that these extracts could be useful components of IPM. The reason for the better performance of hexane extract is due to the fact that it is capable of solubilizing non-polar compounds which possess high biological activity. These findings are in agreement with earlier studies that the presence of several bioactive compounds including terpenoids and flavonoids in *C. inerme* species exhibiting insecticidal and anti-feedant effects.

This is in concordance with our study, especially the work done by Talukdar *et al.* (2014a) who postulated that root and stem extracts of *C. viscosum* were toxic to *T. castaneum* and resulted upto 88% mortality. Azmi (2004) also determined LC₅₀ value of 1460 µg/cm² for *C. inerme* against *Rhizopertha dominica*. In addition, Adesina *et al.* (2019) showed that *C. capitatum* extracts were efficacious against stored grain insects including *Sitophilus oryzae*, *R. dominica*, and *Tribolium castaneum*. They discovered that the ethyl extract had good repellent properties while the hexane extract was toxic through fumigation and killed 80%–90% of pests within 24 hours. Lu *et al.* (2021) and colleagues also revealed that the essential oil of *C. bungei* was very effective as a repellent against stored product insects like *Tribolium castaneum* and the major active compounds were myristicin and linalool.

Nambiar and Ranjini, (2018) on *C. infortunatum* and *Chromolaena odorata* revealed that these botanicals significantly reduced carbohydrate levels in *Orthaga exvinacea* larvae and the latter was more effective. Jadhav *et al.* (2016) reported that *C. inerme*, *C. viscosum* and *C. philippinum* were found to be effective against *Spodoptera litura* and *Helicoverpa armigera* and were found to possess potent anti-feedant and toxic activity. Our findings on the efficacy of *C. inerme* against *A. merione* are in conformity with the result of Jadhav *et al.* (2016) on *C. inerme*, *C. viscosum*, and *C. philippinum*. Talukdar *et al.* (2014b) also found that *C. viscosum* root extract had 44-84% mortality and strong repellent activity against rice weevils.

This indicates the possibility of these botanicals to be used as natural bio-pesticides instead of the synthetic insecticides. The findings of the study are far reaching in pest control since it provides a natural method of controlling pests instead of using synthetic pesticides which are environmentally unfriendly and cause pest resistance. Nevertheless, more studies

are required to determine the exact phytochemicals that are responsible for the observed effects and to assess the effectiveness of these compounds under real-world conditions.

Conclusion

In conclusion, the hexane and acetone extracts of *C. inerme* show promise as eco-friendly insecticides for controlling *A. merione* in castor cultivation. These findings contribute to the growing body of knowledge on botanical pesticides and their role in sustainable agriculture.

Table 1: Extraction yield of *C. inerme* extracts

| Solvent | Extraction yield (%) |
|-----------------|-------------------------|
| Acetone | 2.9 ± 0.03 ^c |
| Petroleum ether | 6.2 ± 0.14 ^b |
| Hexane | 9.6 ± 0.22 ^a |

All values are mean ± SD for triplicate experiments (n=3). Different letters within the same column indicate significant difference between the values (p<0.05). Extraction yield (%) calculated as g of dried extract/500 g DW.

Table 2: Lethal Concentration of Petroleum ether, Hexane and Acetone extracts of *C. inerme* on *A. merione*

| Extracts | Lethal Concentration | Estimated value | Confidence interval | Slope ± SE | Chi square value |
|-----------------|----------------------|-----------------|---------------------|---------------|------------------|
| Petroleum ether | LC10 | 0.80 | 0.140 - 1.548 | 1.016 ± 0.267 | 1.993 |
| | LC20 | 2.18 | 0.932 - 3.486 | | |
| | LC50 | 14.70 | 8.032 - 72.258 | | |
| Hexane | LC10 | 0.525 | 0.156 - 0.952 | 1.314 ± 0.252 | 5.012 |
| | LC20 | 1.136 | 0.512 - 1.744 | | |
| | LC50 | 4.963 | 3.523 - 7.866 | | |
| Acetone | LC10 | 0.382 | 0.029 - 0.905 | 0.911 ± 0.246 | 2.848 |
| | LC20 | 1.161 | 0.283 - 2.044 | | |
| | LC50 | 9.747 | 5.597 - 37.925 | | |

SE – Standard Error




Fig.1 Effect of different extracts of *C. inerme* on feeding action of *A. merione* larvae




Fig.2. Effect of different extracts of *C. inerme* on mortality of *A. merione* larvae

References

- Adesina, J. M., Raghavendra, A., Rajashekar, Y., & Ofuya, T. I. (2019). Potential use of *Clerodendrum capitatum* extracts and its formulation for control of three major stored product beetles. *Food Quality and Safety*, 3(3), 179-185.
- Ahmed, N., Alam, M., Saeed, M., Ullah, H., Iqbal, T., Al-Mutairi, K. A., Shahjeer, K., Ullah, R., Ahmed, S., NAAH Ahmed, Fathy Khater, H., & Salman, M. (2021). Botanical insecticides are a non-toxic alternative to conventional pesticides in the control of insects and pests. *Global decline of insects*, 11(1), 1-19.
- Azmi, M. A. (2004). Toxicity of *Clerodendrum inerme* extract and cyhalothrin against *Rhizopertha dominica* PARC strain and their effect on acid phosphatase and cholinesterase activity. *Biological Sciences-PJSIR*, 47(5), 394-397.
- Duraipandiyan, V., Ignacimuthu, S., & Paulraj, M. G. (2011). Antifeedant and larvicidal activities of Rhein isolated from the flowers of *Cassia fistula* L. *Saudi Journal of Biological Sciences*, 18(2), 129-133. <https://doi.org/10.1016/j.sjbs.2010.11.001>
- Jadhav, G. S., Devarshi, A. A., & Yankanchi, S. R. (2016). Efficacy of certain *Clerodendrum* leaf crude extracts against cutworm, *Spodoptera litura* Fab and cotton bollworm, *Helicoverpa armigera* Hub. *Journal of Entomology and Zoology Studies*, 4(4), 466-472.
- Jannet, H. B., Skhiri, F. H., Mighri, Z., Simmonds, M. S. J., & Blaney, W. M. (2000). Responses of *Spodoptera littoralis* larvae to Tunisian plant extracts and to neoclerodane diterpenoids isolated from *Ajuga pseudoiva* leaves. *Fitoterapia*, 71(2), 105-112.
- Kar, P., Goyal, A. K., Das, A. P., & Sen, A. (2014). Antioxidant and pharmaceutical potential of *Clerodendrum* L.: An overview. *International Journal of Green Pharmacy*, 8(4).
- Lu, X.-X., Hu, N.-N., Du, Y.-S., Borjigidai, A., Zhang, X., & Du, S.-S. (2021). Chemical compositions and repellent activity of *Clerodendrum bungei* essential oil against three stored product insects. *DARU Journal of Pharmaceutical Sciences*, 29(2), 469-475.
- Nambiar, J. G. N., & Ranjini, K. R. (2018). Efficacy of *Clerodendrum infortunatum* and *Chromolaena odorata* on carbohydrate content in haemolymph of the sixth instar larvae of *Orthaga exvinacea* Hampson. *Management Research*, 5(4), 62-68.
- Navatha, L., & Sreedevi, K. (2024). Neem seed attracts adults of castor butterfly, *Ariadne merione* Cramer. *Current Biotica*, 5(2), 241-243.

Talukdar, M. W., Yeasmin, A. M., Islam, M. W., & Hassan, P. (2014a). Insecticidal and repellent activity of *Clerodendrum viscosum* Vent. (Verbenaceae) against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionoidea). *Journal of Entomology*, 7(2), 63-69.

Talukdar, M. W., Akter, M. Y., Islam, M. W., & Hassan, P. (2014b). Mortality and repellent activity of *Clerodendrum viscosum* Vent. (Verbenaceae) against *Sitophilus oryzae* (Coleoptera: Curculionidae). *International Journal of Pharmacognosy*, 1(4), 250-257.

UNDER PEER REVIEW