

**Effect of insecticides on generative potential of Entomopathogenic Nematode,  
*Steinernema abbasi* PN-1 in *Spodoptera litura* Fabricius**

**Abstract**

This study aimed to evaluate the effects of different insecticides at varying concentrations on the progeny production of the entomopathogenic nematode *Steinernema abbasi* PN-1 in *Spodoptera litura* larvae, with a focus on determining the insecticides' compatibility with biological control agents. *S. litura* larvae were treated with a mixture of *S. abbasi* PN-1 and insecticides (fipronil, emamectin benzoate, cyantraniliprole, indoxacarb, and chlorantraniliprole) at three different concentrations (lower, medium, and higher). Progeny production of *S. abbasi* PN-1 was assessed and compared across treatments, with control groups consisting of larvae treated with *S. abbasi* PN-1 alone. The study found that progeny production was significantly influenced by the type and concentration of insecticides. Fipronil exhibited the least negative impact on progeny production across all concentrations, followed by emamectin benzoate, cyantraniliprole, indoxacarb, and chlorantraniliprole, with the latter showing the most significant inhibitory effects. Higher concentrations of insecticides resulted in a greater reduction in progeny production, indicating a dose-dependent relationship. The results suggest that certain insecticides, particularly at higher concentrations, can negatively affect the progeny production of *S. abbasi* PN-1, potentially compromising the effectiveness of EPNs in integrated pest management (IPM) programs. The study highlights the importance of selecting compatible insecticides and their appropriate concentrations to enhance the success of biological control strategies involving EPNs.

**Key words:** Biological control, *Steinernema abbasi* PN-1, Insecticides and *Spodoptera litura*

## **1. Introduction**

The Entomopathogenic nematodes (EPNs) are soil inhabiting, delicate or soft bodied, non-segmented roundworms that are obligate or facultative parasites of insect pests. Entomopathogenic nematodes (EPNs) are members of the Steinernematidae and Heterorhabditidae families. The genera *Steinernema* and *Neosteinerema* belong to the Steinernematidae family, while *Heterorhabditis* is represented by the genus *Heterorhabditis*

[1]. Nematodes of the genera *Heterorhabditis* and *Steinernema* are found worldwide, infecting about 250 distinct insect species belonging to ten orders [2]. The EPNs have simple life cycle, viz., egg, four juvenile stages (separated by moults) and adult stages. The infective stage is J3 (third juvenile stage), and it is known as infective juveniles (IJs). The IJs are non-feeding, free-living active stage capable of withstanding adverse environmental circumstances and the non-availability of host for an extended time period. Nowadays EPNs are mostly utilized in environments where chemical substances fail, such as galleries of boring insects, soil, or situations where pesticide resistance has developed [3]. Farmers in India use a variety of insecticides, including those from both novel classes (such as insect growth regulators, diamides, spinosyns, and avermectins) and conventional classes (such as organophosphates, carbamates, and pyrethroids), to manage insect pests [4]. However, many pesticides have become ineffective due to the development of resistance in pests and their negative impacts on the environment and human health [5]. Therefore, it is essential to reduce the frequency of spraying or the amount of active ingredient in pesticide applications, integrating them with other management strategies like biological control. Combining different control agents can enhance the effectiveness of integrated pest management (IPM) strategies, providing a quicker, more cost-effective, and efficient method for controlling insect pests. When two control agents act independently on a target host, their combined effects can be antagonistic, potentiating, or additive, depending on whether one agent's toxicity is influenced by the other agent [6-7]. Numerous studies have examined the EPNs compatibility with a variety of insecticides to combat insect pests [8-9]. This study has been conducted to determine the effect of selected insecticides (targeting common lepidopteran pests) on generative potential of *Steinernema abbasi* PN-1 under laboratory conditions.

## **2. Material and methods**

### **2.1 Insect and Nematode culture:**

The EPN *Steinernema abbasi* PN-1 was obtained from the Department of Entomology, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, Uttarakhand. The EPN was cultured on late-instar larvae of the greater wax moth, *Galleria mellonella* Linnaeus. A fresh 2-day-old infective juvenile (IJ) solution was used for experiments. Egg masses of *Spodoptera litura* were collected from castor and guava trees at the CRC, Pantnagar, Uttarakhand. Upon hatching, the young larvae were placed in rearing boxes and provided with fresh, sterilized soft castor leaves. The rearing boxes were cleaned and sterilized daily, with

fresh leaves provided regularly. A running culture of *S. litura* was maintained and used for laboratory experiments. Larvae of similar size and weight were selected for the study.

## 2.2 Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticide.

The commonly used insecticides for lepidopteran pests were selected. The 7 days old larvae of *S. litura* were placed in a 9cm petri dish lined with filter paper. The larvae were provided with one castor leaf (5cm× 5cm) treated with selected doses of insecticides (based on a preliminary trial) for one minute and left for 3 minutes to air-dry. The Leaves were applied with 1 ml suspension containing 500IJs/ml of EPN. The Petri dishes were sealed tight immediately and incubated at 27±2 °C and 75±10 per cent RH under laboratory conditions. Control Petri dishes provided with EPN alone. Ten larvae were maintained in each petri dish, with three replicates maintained. The larvae which were dead were washed three times and placed on a white trap according to treatments [10]. The emerged new IJs are harvested regularly and counted under a stereomicroscope.

## 3. Results and discussion

### 3.1 Progeny production of *S. abbasi* PN-1 with lower concentrations of insecticide.

Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticide as shown in Table 1. The progeny production of *S. abbasi* PN-1 in *S. litura* treated with different insecticides was significantly different. The maximum progeny production (9517.33 IJs/larva) was recorded in fipronil which was followed by emamectin benzoate, cyantraniliprole, indoxacarb and chlorantraniliprole where the progeny production was 9412.3, 9386.3, 9375 and 9216.3 IJs/larva respectively. The treatment of *S. abbasi* PN-1 alone yielded 9564.33 IJs/larva. Among the tested insecticides fipronil showed less negative effect followed by emamectin benzoate, cyantraniliprole, indoxacarb and chlorantraniliprole showed a more negative effect on progeny production of *S. abbasi* PN-1.

**Table 1: Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticides.**

| Treatments  | Progeny production/larvae |
|---|---------------------------|
| T1: <i>S. abbasi</i> +Fipronil (0.0013%)            | 9517.33 (97.56) *         |
| T2: <i>S. abbasi</i> +Emamectin benzoate (0.00012%) | 9412.33 (96.89) *         |
| T3: <i>S. abbasi</i> +Indoxacarb (0.008%)           | 9375 (97.02) *            |

|   |                   |
|---|-------------------|
| T4: <i>S. abbasi</i> +Chlorantraniliprole (0.0015%) | 9216.33 (96) *    |
| T5: <i>S. abbasi</i> +Cyantraniliprole (0.003%)     | 9386.33 (96.88) * |
| T6: Control ( <i>S. abbasi</i> )                    | 9564.33 (97.8) *  |
| CD  | 0.627             |
| SE(m)   | 0.201             |

*S. abbasi* PN-1 @ 500IJs/ml, \*values in the parenthesis are square root transformed values.

### 3.2 Progeny production of *S. abbasi* PN-1 with medium concentrations of insecticide.

Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticide as shown in Table 2. The progeny production of *S. abbasi* PN-1 in *S. litura* treated with different insecticides was significantly different. The maximum progeny production (9414 IJs/larva) was recorded in fipronil which was followed by emamectin benzoate, cyantraniliprole, indoxacarb and chlorantraniliprole where the progeny production was 9287, 9267, 9033.6 and 8923 IJs/larva respectively. The treatment of *S. abbasi* PN-1 alone yielded 9564.33 IJs/larva. Among the tested insecticides fipronil showed less negative effect followed by emamectin benzoate, cyantraniliprole, indoxacarb and chlorantraniliprole showed a more negative effect on progeny production of *S. abbasi* PN-1.

**Table 2: Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticides.**

| Treatments:   | Progeny production/larvae |
|---|---------------------------|
| T1: <i>S. abbasi</i> +Fipronil (0.0025%)            | 9414 (96.38) *            |
| T2: <i>S. abbasi</i> +Emamectin benzoate (0.00024%) | 9287 (94.48) *            |
| T3: <i>S. abbasi</i> +Indoxacarb (0.0016%)          | 9033.67 (95.46) *         |
| T4: <i>S. abbasi</i> +Chlorantraniliprole (0.003%)  | 8923 (93.38) *            |
| T5: <i>S. abbasi</i> +Cyantraniliprole (0.005%)     | 9267.33 (94.48) *         |
| T6: Control ( <i>S. abbasi</i> )                    | 9564.33 (97.80) *         |
| CD  | 0.461                     |
| SE(m)   | 0.148                     |

*S. abbasi* PN-1 @ 500IJs/ml, \*values in the parenthesis are square root transformed values.

### 3.3 Progeny production of *S. abbasi* PN-1 with higher concentrations of insecticide.

Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticide as shown in Table 3. The progeny production of *S. abbasi* PN-1 in *S. litura* treated with different insecticides was significantly different. The maximum progeny

production (9289.6 IJs/larva) was recorded in fipronil which was followed by emamectin benzoate, indoxacarb, cyantraniliprole and chlorantraniliprole where the progeny production was 9112.3, 8927, 8926.3 and 8719.6 IJs/larva respectively. The treatment of *S. abbasi* PN-1 alone yielded 9564.33 IJs/larva. Among the tested insecticides fipronil showed less negative effect followed by emamectin benzoate, indoxacarb, cyantraniliprole, and chlorantraniliprole showed a more negative effect on progeny production of *S. abbasi* PN-1.

All the treatments with lower concentration insecticides have less effect on progeny production in comparison to the higher concentrations of insecticides with *S. abbasi* PN-1. Among the tested insecticides fipronil showed less negative effect followed by indoxacarb, cyantraniliprole, emamectin benzoate and chlorantraniliprole showed a more negative effect on progeny production of *S. abbasi* PN-1. Some insecticides are less toxic to EPNs may be because of the presence of the butyrylcholinesterase in the synapse of EPNs. Selkirk *et al.* [11] reported the action of butyrylcholinesterase as a first line of defense in nematodes and it may protect against early attack by inhibitors to acetylcholinesterase therefore, in contrast to such inhibitor compounds, it is directly involved in the defense. Bhat *et al.* [12] reported that sometime IJs are sheathed with previous molt cuticle and it may not allow the penetration of some insecticides and other substances into the IJs body. The findings were similar to studies conducted [13], where they tested the effect of insecticide treated larvae of *Hyphantria cunea* (Fall webworm) on the progeny production of *S. carpocapsae* and they observed no negative effect of insecticides on progeny production of *S. carpocapsae* under laboratory conditions. Ozdemir *et al.* [14] studied the progeny production of *S. feltiae* and *H. bacteriophora* exposed to insecticide and observed that imidacloprid recorded no negative effect on progeny production but spinosad recorded a negative effect on progeny production both the EPN species in *G. mellonella*.

**Table 3: Progeny production of *S. abbasi* PN-1 in *S. litura* treated with a mixture of *S. abbasi* PN-1 and insecticides.**

| Treatments  | Progeny production /larvae |
|---|----------------------------|
| T1: <i>S. abbasi</i> +Fipronil (0.005%)             | 9289.67 (97.03) *          |
| T2: <i>S. abbasi</i> +Emamectin benzoate (0.00036%) | 9112.33 (95.05) *          |
| T3: <i>S. abbasi</i> +Indoxacarb (0.0024%)          | 8927 (96.34) *             |
| T4: <i>S. abbasi</i> +Chlorantraniliprole (0.0045%) | 8719.67 (94.46) *          |
| T5: <i>S. abbasi</i> +Cyantraniliprole (0.007%)     | 8926.33 (96.27) *          |
| T6: Control ( <i>S. abbasi</i> )                    | 9564.33 (97.80) *          |

|       |       |
|-------|-------|
| CD    | 0.089 |
| SE(m) | 0.6   |

*S. abbasi* PN-1 @ 500IJs/ml, \*values in the parenthesis are square root transformed values.

#### **4. Conclusion:**

This study evaluated the effects of different concentrations of insecticides on the progeny production of *Steinernema abbasi* PN-1 in *Spodoptera litura*. Our findings indicate that the progeny production of *S. abbasi* PN-1 was significantly affected by the type and concentration of the insecticides used. Among the insecticides tested, fipronil consistently showed the least negative impact on progeny production across all concentrations, followed by emamectin benzoate, cyantraniliprole, indoxacarb, and chlorantraniliprole, with the latter exhibiting the most substantial inhibitory effects. The negative impact on progeny production increased with the concentration of the insecticides, underscoring the sensitivity of *S. abbasi* PN-1 to higher doses of chemical treatments. The presence of butyrylcholinesterase and the sheathing of infective juveniles (IJs) with the previous molt cuticle may provide some level of protection against certain insecticides. These findings contribute to the understanding of the interaction between entomopathogenic nematodes (EPNs) and insecticides, offering insights for integrated pest management (IPM) strategies where the compatibility of biocontrol agents and chemical treatments is crucial. The study suggests that careful selection and concentration of insecticides are essential to minimize the negative effects on beneficial EPNs, thereby enhancing their efficacy in biological control programs.

#### **Data availability statement**

All the data have been provided in the main text of the manuscript.

#### **Ethics approval statement**

Not Applicable

#### **Consent to participate**

Not Applicable

#### **Consent to publish**

Not Applicable

## 5. References:

1. Vireesha P, Pandey R, Shashikala M, Mehra N. Susceptibility of Different Developmental Stages of the Tobacco Cutworm, *Spodoptera litura* (Fabricius) to Entomopathogenic Nematode, *Steinernema abbasi* PN-1. *Uttar Pradesh J Zool.* 2024 Jun 24;45(14):87-93.
2. Poinar GO. Nematodes for biological control of insects. Boca Raton, Florida: CRS Press; 2018. 289 p.
3. Abdel Gawad MM, Ruan W, Hammam M. Entomopathogenic nematodes: Integrated pest management and new vistas. *Egypt J Agronematol.* 2023;22(1):1-18.
4. Natikar PK, Balikai RA. Present status on bio-ecology and management of tobacco caterpillar, *Spodoptera litura* (Fabricius) - an update. *Int J Plant Prot.* 2017;10(1):193-202.
5. Gandhi K, Patil RH, Srujana Y. Field resistance of *Spodoptera litura* (Fab.) to conventional insecticides in India. *Crop Prot.* 2016;88:103-8.
6. Robertson JL, Jones MM, Olguin E, Alberts B. Bioassays with arthropods. Boca Raton, Florida: CRC Press; 2017.
7. Pavan, J.S., Patel, N.B., Raghunandan, B.L., Baldaniya, A.M. and Bhatt, N.A., 2024. Comparative efficacy of nucleopolyhedrovirus (NPV) alone and in conjunction with chemical insecticides against fall armyworm, *Spodoptera frugiperda* (JE Smith) (Noctuidae: Lepidoptera) under laboratory conditions. *Int J Trop Insect Sci.*, pp.1-12.
8. Khan RR, Ali RA, Ali A, Arshad M, Majeed S, Ahmed S, et al. Compatibility of entomopathogenic nematodes (Nematoda: Rhabditida) and the biocide, spinosad for mitigation of the armyworm, *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Egypt J Biol Pest Control.* 2018;28(1):1-6.
9. Khan RR, Arshad M, Aslam A. Additive interactions of some reduced-risk biocides and two entomopathogenic nematodes suggest implications for integrated control of *Spodoptera litura* (Lepidoptera: Noctuidae). *Sci Rep.* 2021;11(1):1-9.
10. Woodring JL, Kaya HK. *Steinernematid* and *heterorhabditid* nematodes: A handbook of biology and techniques. Southern Cooperative Series Bulletin. USA; 1988. 331 p.
11. Selkirk ME, Henson SM, Russell WS, Hussein AS. Acetylcholinesterase secretion by nematodes. In: Kennedy H, editor. Parasitic nematodes—molecular biology, biochemistry and immunology. Wallingford, UK: CAB International; 2001. p. 211-28.
12. Bhat AH, Chaubey AK, Upadhyay SK. Morphotaxometric and molecular validation of entomopathogenic nematode, *Steinernema abbasi* (Rhabditida: Steinernematidae) with

mucronate processes in adults of second generations off sub-humid region, Uttar Pradesh, India. World J Pharm Pharm Sci. 2016;5(1):1558-79.

13. Nalinci E, Karagoz M, Gulcu B, Ulug D, Gulsen SH, Cimen H, et al. The effect of chemical insecticides on the scavenging performance of *Steinernema carpocapsae*: Direct effects and exposure to insects killed by chemical insecticides. J Invertebr Pathol. 2021;184(1):107-41.
14. Özdemir E, İnak E, Evlice E, Laznik Z. Compatibility of entomopathogenic nematodes with pesticides registered in vegetable crops under laboratory conditions. J Plant Dis Prot. 2020;127(4):529-35.

UNDER PEER REVIEW