

Influence of green synthesized silver and zinc nanoparticle on seed quality attributes in Red sanders (*Pterocarpussantalinus* Linn.)

Abstract: Red Sanders (*Pterocarpussantalinus*), or Raktachandana, is an endangered tree native to the Southern Indian Peninsula, valued for its medicinal and dye properties. However, its seed germination is hindered by a hard seed coat, phenolic compounds and hormonal imbalances. The laboratory experiment was conducted at the Green Nano-Technology Laboratory and Department of seed science and technology, University of Agricultural Sciences, Dharwad, during 2023. The experiment used a completely randomized design (CRD) with three replications and evaluated seven presowing treatments: T₁ (AgNPs at 1000 ppm for 24 hours), T₂ (ZnNPs at 2000 ppm for 24 hours), T₃ (Conc. H₂SO₄ for 12 minutes), T₄ (NaDC 500 ppm), T₅ (Dewinging), T₆ (Soaking in hot water at 100°C for 5 minutes) and T₇ (control). The results indicated that seeds treated with ZnNPs at 2000 ppm (T₂) exhibited highest germination rate (56%), germination speed (1.04) along with root length (17.05cm), shoot length (16.33cm) and highest dry weight (6.22 g) and highest seedling vigour Index-I (1869). In comparison, the H₂SO₄ treatment (T₃) resulted in 51.67 per cent germination and a speed of 0.95, with root length (15.28 cm), shoot length (16.22 cm), dry weight (5.24 g) and seedling vigour Index (1628) slightly lower but still significant. The control (T₇) had the lowest performance with 31 per cent germination, a speed of 0.36, a root length (15.28 cm), shoot length (16.22 cm), lower dry weight (1.63 g) and the lowest Seedling Vigour Index (649). Overall, ZnNPs at 2000 ppm (T₂) were the most effective in enhancing germination, seedling growth and overall vigour in red sanders.

Keywords: *Pterocarpussantalinus* Linn, Seed priming, Zn nanoparticle

I. Introduction

Red sanders (*Pterocarpussantalinus* Linn.) is an endangered, endemic tree of the Southern Indian Peninsula, belonging to the Fabaceae family. It thrives in dry, hilly, rocky areas. The wood of red sanders is prized for its rich red colour and fine grain, making it highly sought after for furniture, musical instruments, and carvings. Its high market value drives both legal and illegal trade. However, it has poor and staggered germinability (33%) (Anuradha et al., 2019) due to the presence of phenolic compounds, hormonal imbalance and

Comment [A1]: Its not necessary to mention in abstract

dormancy in pods and seed respectively. Seed germination and early seedling establishment depend on several factors, with seed dormancy being a major constraint in forest species. Dormancy, where viable seeds fail to germinate under suitable conditions, can be due to environmental factors or the seeds own inhibitory mechanisms, which can be external, internal, or both. In the process of dormancy breaking, germination improvement is a complex phenomenon through silver and zinc nanoparticles and comprises of a series of physiological and biochemical changes. Hence the present investigation is aimed to evaluate the effects of green synthesized silver (Ag) and zinc (Zn) nanoparticles and different treatments on the seed quality of Red sanders.

Comment [A2]: List the factors

II. Materials and Methods

The laboratory experiment was conducted to investigate the influence of green synthesized Ag and Zn nanoparticles on seed quality parameters in Red sander (*Pterocarpussantalinus* Linn.) during 2023 at the Department of Seed Science and Technology and green synthesis of nanoparticles carried out in Green Nanotechnology Laboratory, University of Agricultural Sciences, Dharwad. The experiment was laid out using an incompletely randomized design with 3 replications and 17 treatments viz., Red sander seeds were primed with Silver and Zinc nanoparticles (ZnNPs) of concentration 500, 1000, 1500 and 2000 ppm respectively for 12 h and 24 h and seeds primed with 1mM AgNO₃ and 1 mM ZnSO₄, respectively along with control (untreated). The seed-to-solution ratio for the nanopriming treatment was 1:2. Later the seeds were shade dried to bring down to original moisture content under ambient conditions. Observations of the seed quality parameters were recorded.

Comment [A3]: Which design is used weather completely or incomplete RBD

The synthesis, characterization and standardization of nanoparticles silver (Ag) and zinc (Zn) nanoparticles were confirmed through UV-visible spectroscopy, PSA, SEM and XRD. The effective concentrations of Ag and ZnNPs for priming duration were standardized through seed priming. The observations of germination, root length, shoot length, best priming duration and concentration of AgNPs and ZnNPs were standardized and primed the seeds were primed with 17 different treatments. Red sander seeds were primed as per according to best silver and zinc nanoparticle treatment, and T₁ and T₂ were subjected to out with other final treatments. The treated seeds were evaluated for seed quality parameters i.e. seed germination (%), speed of germination, final mean germination, shoot length, root length, seedling dry weight and seedling vigour index. Treatment details are as follows:

T₁: Seed priming with AgNPs at 1000 ppm for 24 h, T₂: Seed priming with ZnNPs at 2000 ppm for 24 h, T₃: Seed treatment with Concentrated H₂SO₄ for 12 min, T₄: Seed

treatment with NaDC 500 ppm for 15 min, T₅: Dewinging by secature, T₆:Soaking seeds in hot water at 100°C for 5 minutes, T₇: Control (Untreated).

III. Observations to be recorded

Germination (%)

The standard germination test was conducted (ISTA, 1976) by sowing the seeds in sand media in four replications. For the final count of the germination test, number of seeds germinated was counted and expressed as germination percentage.

$$\text{Germination (\%)} = \frac{\text{Number of normal seedlings}}{\text{Total number of seeds}} \times 100$$

Speed of germination

Number of normal seedlings was counted everyday until the end of the test period. The speed of emergence was calculated using the formula suggested by Maguire (1962).

$$\text{Speed of emergence} = n_1/d_1 + n_2/d_2 + n_3/d_3 + \dots$$

where, n = the number of emerged seedlings,

d = number of days.

Final mean germination

The final mean germination rate was calculated by dividing the cumulative germination percentage/total number of days.

Seedling dry weight (g)

For the germination test, the same ten normal seedlings used for measuring the root and shoot length were kept in a butter paper packet and dried in hot air oven maintained at 80 ± 2°C for 24 hours. Then the seedlings were subsequently cooled in desiccators for 30 minutes and the weights of the dry seedlings were recorded using an electronic balance and was expressed in milligrams.

~~Ten normal seedlings after observing root and shoot length they are put in butter paper (Hot air oven).~~ After observing the length of the roots and shoots, ten normal seedlings were placed on butter paper (hot air oven).

Root length (cm)

~~From~~For the germination test, the ~~same~~ ten normal seedlings which are used for measuring the shoot length ~~was~~were used for measuring the root length. The root length was measured from ~~the~~tip of ~~the~~root to the hypocotyl point and the mean length was calculated and expressed in centimetres.

Shoot length (cm)

~~From~~For the germination test, ten normal seedlings were randomly selected from each treatment on ~~the~~twelfth day and the shoot length was measured from the tip of ~~the~~shoot to the hypocotyl point and the mean length was calculated and expressed in centimeters.

Seedling Vigour Index

The Seedling Vigour Index was calculated by using the formula suggested by Abdul-Baki and Anderson (1973).

Seedling vigour index = Germination (%) x [root length (cm) + shoot length (cm)]

SVI-I = Germination (%) × Total seedling length (cm)

SVI-II = Germination (%) × Seedling dry weight (mg)

IV. Results and Discussion

Among the treatments, T_2 (ZnNPs 2000 ppm) primed seeds recorded significantly higher seed germination (56.00 %), speed of germination (1.04), final mean germination (1.20) ~~was recorded in T_2 (ZnNPs 2000 ppm) primed seeds when~~ compared to control T_7 seeds (31.00 %), (0.36), (0.45) respectively (Table. 1). Thenanoparticles entering into seed coat pores, causes increased penetration of water molecules and inducing ROS-generating/starch-degrading enzyme activity for physiological enhancement of seed germination (Mahakhamet *et al.*, 2017; nanoprimering technology for enhancing seed germination and starch metabolism of aged rice seeds using photosynthesized silver nanoparticles, Khodakovskaya *et al.*, 2011). Zinc is essential for the preservation and defence of cell membrane structural stability, ~~as~~reported by Cakmak (2000), ~~and~~ Ajouriet *al.* (2004) found that Zn seed priming significantly improved barley seed germination and seedling development; these findings may indicate that high Zn concentrations in seeds play critical physiological roles during seed germination and early seedling growth (Prasad *et al.*, 2012). Zinc oxidenanoparticles ~~additionally boosted also increased~~ peanut germination, growth and yield. Nanoparticles ~~contribute~~ positively ~~contribute to increasing~~

Comment [A4]: Give

enhancing seed quality ~~by implying in process that involves~~ via processes that involve oxidation and reduction during germination. This is achieved by generating superoxide ion radicals, which effectively neutralize free radicals present during seed germination. ~~Enhanced—The increased amount of~~ oxygen produced in this process leads to ~~higher~~ increased respiration ~~and this, which~~ further encourages germination. These results are in ~~compliance~~ accordance with the findings of (Shyla and Natarajan, 2014).

Significantly higher mean shoot (cm) length and root length ~~was~~ recorded in T₂ (ZnNPs 2000 ppm) primed seeds (16.33 and 17.05 cm) respectively as compared to control *i.e.*, T₇ (Untreated) (11.02 and 9.92 cm) respectively in Table.2. This is mainly due to the ~~enhancement—of increase~~ in the growth hormones in seeds ~~caused~~ by the treatment ~~of~~ with different concentrations of zinc oxide nanoparticles, ~~which~~ improved other growth parameters in green gram (Lakshmi *et al.*, 2017). Zinc nanoparticles enhance plant growth and improve nutrient uptake ~~due to~~ because of their small size and high surface area, ~~making zinc more—available~~ increasing the availability of zinc for absorption by plant roots. Micronutrients ~~is~~ are vital for physiological processes such as enzyme activation, protein synthesis and hormone regulation (Raliya *et al.*, 2019).

Significantly higher mean seedling dry weight was recorded in T₂ (ZnNPs 2000 ppm for 24 h) primed seeds (6.22 g) as compared to control *i.e.*, T₇ (Untreated) (1.63 g). Suriyaprabha *et al.* (2012) reported that the content of organic substances in the primed seeds, including protein, chlorophyll, and phenols, was increased by the application of nanoparticles to the seeds. ~~Owing to~~ the increase in seedling length (cm) through nanopriming, primed seeds recorded higher seedling dry weight.

Significantly higher mean Seedling Vigour Index-I and Seedling Vigour Index – II was recorded in T₂ (ZnNPs 2000 ppm for 24 h) primed seeds (1869 and 125) respectively as compared to control *i.e.*, T₇ (untreated) (649 and 51) respectively in Table 3. This is due to the ~~result of~~ increased synthesis of tryptophan and auxin, a hormone that stimulates longitudinal growth, as well as the ~~effects~~ ability of nanoparticles ~~on—boosting to increase~~ cell division and growth, particularly in meristematic points (Kaya and Higgs, 2002). This could lead to an increase in the length and height of internodes. This conclusion is consistent with findings from studies on encouraging and accelerating of longitudinal growth in *Tanacetum parthenium* (Mahajan *et al.*, 2011; Shahhoseini *et al.*, 2020).

The mechanism of NP uptake ~~is~~ generally considered ~~as~~ active transport, ~~which that~~ includes cellular processes such as signalling and regulation of the plasma membrane (Etxeberria *et al.*, 2006). NPs can ~~move~~ pass through tissues ~~via:~~ apoplastic and

symplastic routes. Apoplastic transport occurs outside the plasma membrane and through the extracellular spaces, cell walls of adjacent cells, and xylem vessels (Sattelmacher, 2001); in contrast, symplastic transport involves movement of water and substances between the cytoplasm of adjacent cells through specialized structures called plasmodesmata (Roberts and Oparka, 2003) and sieve plates. The apoplastic pathway is important for radial movement within plant tissues; it allows nanomaterials to reach the root central cylinder and vascular tissue.

Compared ~~to~~with the untreated control, ~~the~~ seed treatment with nanoparticles considerably changed the seed quality parameters and consistently preserved the viability and vigour of the seeds. In terms of seed germination (%), ~~the~~ speed of germination, final mean germination, seedling shoot length (cm), seedling root length (cm), dry weight and Seedling Vigour indices significantly increased in the seeds treated with nanoparticles ~~significantly increased as compared to~~ compared with the control. The seed treatment with ZnNPs outperforms other treatments examined, including the control.

V. Conclusion

Green synthesis of AgNPs and ZnNPs was successfully carried out ~~through leaf extracts of guava (*Psidium guajava*)~~ as reducing and capping agents. As compared to control significantly higher seed quality parameters were recorded in seeds primed with ZnNPs (2000 ppm for 24 h) recorded in Red sander seeds. The outcome of this experiment unequivocally demonstrates that ZnNPs can be used as an effective priming agent for enhancing seed quality.

Comment [A5]: Mention the process how it is processed with guava leaves in material and methods

Table 1: Effects of green synthesized silver and zinc nanoparticles in combination with other treatments on seed germination, speed of germination and final mean germination of Red sander

Treatments	Germination (%)	Speed of germination	Final mean germination
T ₁ - Seed priming with AgNPs at 1000 ppm for 24 h	42.00 (40.38)*	0.77	0.76
T ₂ - Seed priming with ZnNPs at 2000 ppm for 24 h	56.00 (48.43)*	1.04	1.20
T ₃ - Seed treatment with Conc H ₂ SO ₄ for 12 min	51.67 (45.94)*	0.95	1.16
T ₄ - Seed treatment with NaDC 500 ppm for 15 min	35.33 (36.45)*	0.50	0.60
T ₅ - Dewinging by secature	48.33 (44.03)*	0.89	0.93
T ₆ - Soaking seeds in hot water at 100 °C for 5 min	37.67 (37.85)*	0.62	0.64
T ₇ - Control (Untreated)	31.00 (33.82)*	0.36	0.45
S.Em (±)	1.09	0.02	0.02
CD at 1%	3.31	0.05	0.07

Note :* values in the parenthesis are arc sine transformed

AgNPs : Silver nanoparticles,

ZnNps : Zinc nanoparticles, Conc

H₂SO₄ : Concentrated sulphuric acid,

NaDC : Sodium de-oxycholate

Table 2: Effects of green synthesized silver and zinc nanoparticles in combination with other treatments on the root length, root length and seedling dry weight of Red sander

Treatments	Shoot length (cm)	Root length (cm)	Seedling dry weight (g)/10 seedlings
T ₁ - Seed priming with AgNPs at 1000 ppm for 24 h	13.75	12.17	2.97
T ₂ - Seed priming with ZnNPs at 2000 ppm for 24 h	16.33	17.05	6.22
T ₃ - Seed treatment with Conc H ₂ SO ₄ for 12 min	16.22	15.28	5.24
T ₄ - Seed treatment with NaDC 500 ppm for 15 min	11.67	11.79	1.96
T ₅ - Dewinging by secature	14.39	13.05	5.09
T ₆ - Soaking seeds in hot water at 100 °C for 5 min	12.55	12.08	2.48

T ₇ - Control (Untreated)	11.02	9.92	1.63
S.Em (±)	0.33	0.31	0.08
CD at 1%	1.00	0.95	0.24

Table 3: Effects of green synthesized silver and zinc nanoparticles in combination along with other treatments on seedling the vigour indices of Red sander seedlings

Treatments	Seedling Vigour Index-I	Seedling Vigour Index-II
T ₁ - Seed priming with AgNPs at 1000 ppm for 24 h	1089	125
T ₂ - Seed priming with ZnNPs at 2000 ppm for 24 h	1869	348

T ₃ - Seed treatment with Conc H ₂ SO ₄ for 12 min	1628	271
T ₄ - Seed treatment with NaDC 500 ppm for 15 min	829	69
T ₅ - Dewinging by secature	1326	246
T ₆ - Soaking seeds in hot water at 100°C for 5 min	928	93
T ₇ - Control (Untreated)	649	51
S.Em (±)	28.47	3.76
CD at 1%	86.36	11.40

UNDER PEER REVIEW

fig .1 Effect of green synthesized silver and zinc nanoparticles along with other treatments on germination (%), speed of germination (%) and final mean germination of Red sander (*Pterocarpus santalinus* Linn.)





fig . 2 Effect of green synthesized silver and zinc nanoparticles along with other treatments on seedling vigour indices of Red sander (*Pterocarpus santalinus* Linn.)



References:

- Abdul Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1, *crop science*. 1973, 13(6):630-633.
- Ajouri A, Asgedom and Becker M, 2004, Seed priming enhances germination and seedling growth of barley under conditions of P and Zn deficiency. *Journal of Plant Nutrition and Soil Science*, 167(5): 630-636.
- Anonymous. International Rules for Seed Testing, Seed Science and Technology. 2019: 14:1-14.
- Anuradha M, Indu B K and Balasubramanya S, 2019, Propagation of red sanders: an overview. *Red Sanders: Silviculture and Conservation*: 85-100.
- Cakmak I, 2000, Tansley Review No. 111 Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *The New Phytologist*, 146(2): 185-205.
- Etxeberria E, Gonzalez P, Baroja-Fernandez E and Romero J P, 2006, Fluid phase endocytic uptake of artificial nano-spheres and fluorescent quantum dots by sycamore cultured cells: evidence for the distribution of solutes to different intracellular compartments. *Plant Signal Behav* 1: 196-200
- Kaya C and Higgs D, 2002, Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. *Scientia Horticulturae*, 93(1): 53-64.
- Khodakovskaya M V, de Silva K, Nedosekin D A, Dervishi E, Biris A S, Shashkov E V, Galanzha E I and Zharov V P, 2011, Complex genetic, photothermal and photoacoustic analysis of nanoparticle-plant interactions. *Proceedings of the National Academy of Sciences*, 108(3): 1028-1033.
- Lakshmi S J, Bai R R, Sharanagouda H, Ramachandra C T, Nadagouda S and Doddagoudar S R, 2017, Biosynthesis and characterization of ZnO nanoparticles from spinach (*Spinacia oleracea*) leaves and its effect on seed quality parameters of green gram (*Vignaradiata*). *International Journal of Current Microbiology and Applied Sciences*, 6(9): 3376-3384.
- Maguire J D, 1962, Speed of germination-aid in selection and evaluation for seedling emergence and vigour.

- Mahajan P, Dhoke S K and Khanna A S, 2011, Effect of nano ZnO particle suspension on growth of mung (*Vignaradiata*) and gram (*Cicerarietinum*) seedlings using plant agar method. *Journal of Nanotechnology*, 2011(1): 696535.
- Mahakham W, Sarmah A K, Maensiri S and Theerakulpisut P, 2017, Nanopriming technology for enhancing germination and starch metabolism of aged rice seeds using photosynthesized silver nanoparticles. *Scientific Reports*, 7(1): 8263.
- Prasad T N V KV, Sudhakar P, Sreenivasulu Y, Latha P, Munaswamy V Reddy, KR, Sreeprasad TS, Sajanlal PR. and Pradeep T, 2012, Effect of nanoscale zinc oxide particles on the germination, growth and yield of peanut. *Journal of Plant Nutrition*, 35(6): 905-927.
- Raliya R, Saharan V, Choudhary K, Summarwar S, Gulecha K, Gupta V and Sain P M, 2019, Nanomaterials synthesis and characterization. In *Nanoscale Engineering in Agricultural Management*: 1-10.
- Roberts A G, Oparka K J, 2003, Plasmodesmata and the control of symplastic transport. *Plant Cell Environ* 26:103–124.
- Sattelmacher B, 2001, The apoplast and its significance for plant mineral nutrition. *New Phytology*, 149: 167–192
- Shahhoseini R, Azizi M, Asili J, Moshtaghi N and Samiei L, 2020, Effects of zinc oxide nanoelicitors on yield, secondary metabolites, zinc and iron absorption of Feverfew (*Tanacetum parthenium* (L.)). *Acta Physiologiae Plantarum*, 42: 1-18.
- Shyla K K and Natarajan N, 2014, Customizing zinc oxide, silver and titanium dioxide nanoparticles for enhancing groundnut seed quality. *Indian Journal of Science and Technology*, 7(9): 1376-1381.
- Suriyaprabha R, Karunakaran G, Yuvakkumar R, Prabu P, Rajendran V, Kannan N. Growth and physiological responses of maize (*Zea mays* L.) to porous silica nanoparticles in soil. *Journal of Nanoparticle Research*. 2012: 14:1-14.