

Original Research Article

Analysis of Growth and Instability of Rice Production in Madhya Pradesh, India: A District Level Study

ABSTRACT

Aim: Rice is one of the important food crops cultivated in Madhya Pradesh state covering 4.55% rice area and 3.72% rice production of the country. Its contribution is very vital in agricultural growth of the country. But, yield of the state is less than the national average yield. So, for further improvement of its contribution in the agriculture sector of the country, it is essential to study the growth pattern of rice production in the state. Therefore the present study has been undertaken to examine the growth rate and instability in area, production and yield of rice in Madhya Pradesh state.

Place and Duration of Study: The district-wise data on rice of Madhya Pradesh state was collected from the Directorate of Economics and Statistics, Ministry of Agriculture and Farmers Welfare, Government of India and from the online sources of the state departments for the period 1992-93 to 2019-20.

Methodology: The data was divided into four periods for analysis. Compound annual growth rate and Cuddy-Della Valle Instability index have been computed for the four periods. The districts have been classified into five classes based on the classification of instability. These classes are very low instability, low instability, medium instability, high instability and very high instability.

Results: For the state, the growth rate for area ranges from -0.80 to 4.23; for production ranges from 0.90 to 12.82 and for yield ranges from 1.31 to 8.24. The highest growth rate for area and production was observed in Hosangabad and Raisen districts respectively in period III; and for yield Datia district in period I. For the state instability ranges from 3.77 to 9.00 for area, 5.93 to 30.70 for production and 6.29 to 22.57 for yield. None of the districts were classified as very low instability for production and yield.

Conclusion: Most of the districts recorded medium instability in period I, II and III for yield.

Keywords: *Rice, Production, Growth Rate, Instability, Madhya Pradesh*

1. INTRODUCTION

Rice is the staple food for about half of the world's population and more than two-thirds of the Indian population. India ranks first in rice area and second in rice production next to China. In India, rice was grown in more than 46.27 million hectare area with the production of 129.47 million tons of milled rice in 2021-22 contributing approximately 25% of the global rice production. Rice cultivation engages most of the workforce in the economy as the source of livelihood for those people. Rice accounts for 40% of the total food grain production occupying 35.6% of the food grain area of the country.

Madhya Pradesh is the tenth largest rice producing state of the country producing 4.81 million tons of rice in 2.11 million hectares of area with an average yield of 2.80 tons per hectare contributing 3.72% of rice production and 4.56% rice area of the country.

For higher growth of agriculture, a quantitative assessment of the contribution of different factors to agricultural output growth is important for reorienting the programmes and prioritizing agricultural development. Various factors affect the growth of agricultural output. Major of these factors is area and yield [1,2]. These major sources of output growth have significance in finalizing programmes of agricultural development and priorities of investment in it [3,4]. Hence, it may be vital to find the reasons for growth rates different from one another, to remove the bottlenecks to achieve the fast development of the agricultural sector [5]. The study of instability is also required to find out the fluctuation in the trend for area, production and yield of rice which is severely affecting the production, and indirectly employment and income distribution thereby hampering the economic growth of the state.

Samal et al [6] studied the state-wise analysis of production of rice in India. District wise growth rate and instability of rice have been analyzed by the researchers for some states [7-12]. Some researchers studied growth and instability in Madhya Pradesh. Kumar et al [13] studied the growth and instability of paddy and wheat crop in Gwalior district of Madhya Pradesh; Mishra et al [14] analyzed the growth and instability of major *kharif* crops in Madhya Pradesh; Balai et al [15] studied the growth, decomposition and instability of major *rabi* pulse crops in Madhya Pradesh, but the district-wise study on rice in Madhya Pradesh state over the long period is not available.

Therefore, the present study is undertaken to analyze the district wise growth rate and instability of area, production and yield of rice in Madhya Pradesh state.

2. MATERIAL AND METHODS

The present study is done based on the secondary rice data of Madhya Pradesh state collected during the years 1992-93 to 2019-20. The district-wise data on the area, production and yield of rice of Madhya Pradesh state was collected from the Directorate of Economics and Statistics, Department of Agriculture, Cooperation and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Government of India during the last ten years. Last twenty eight years rice data from 1992-93 to 2019-20 have been used in the present study. There are fifty five districts in Madhya Pradesh. Some districts were carved out to become new districts during last few years. The analysis have been done from 1992-93 data, therefore the bifurcated districts were merged for the analysis with sum of twenty five districts. For lucidity, the twenty eight years rice data have been divided into four periods; period I (1992-93 to 1999-2000), period II (2000-01 to 2009-10), period III (2010-11 to 2019-20) and lastly the overall period as a period IV (1992-93 to 2019-20).

The district-wise growth rate of area, production and yield of Madhya Pradesh state for each period is computed to study the growth pattern in area, production and yield of that district.

The compound growth rate was estimated using the following exponential [16].

$$Y = ab^t$$

$$\text{Log } Y = \log a + t \log b$$

$$\text{CGR}(r) = [\text{Antilog}(\log b) - 1] \times 100$$

where,

CGR = Compound growth rate

t = time in the year

Y = area/production/productivity

a and b = Regression parameters

The performance of agricultural output is affected by climatic factors, the growth rate has been calculated based on three years of average data [16-18].

Instability means deviation from the trend. In agriculture, instability is an inherent characteristic due to weather conditions, seasonal variation in area, yields and production of crops from year to year. The instability in area, production and yield of rice is computed to measure the variability using an index of instability called the Cuddy-Della Valle index [19]. This method is used to examine the extent of risk involved in crop production.

The instability in area, production and yield has been computed using the following Cuddy-Della Valle Index.

$$CDVI = CV \times \sqrt{(1 - \text{Adj. } R^2)}$$

where,

CDVI = Cuddy-Della Valle Instability index (per cent)

CV = Coefficient of variation (per cent)

Adj. R^2 = Coefficient of determination from a time trend regression adjusted by the number of degrees of freedom

Classification methodology for instability proposed by Jambhulkar et. al., 2023a, b) [10,11] has been used to classify the instability into five classes as follows:

| Class | Range of instability |
|-----------------------|----------------------|
| Very low instability | 0 to 5 |
| Low instability | 5 to 15 |
| Medium instability | 15 to 30 |
| High instability | 30 to 50 |
| Very high instability | > 50 |

3. RESULTS AND DISCUSSION

The area (million hectares), production (million tons) and yield (t/ha) of rice in Madhya Pradesh state from 1992-93 to 2019-20 is depicted in Figure 1. The rice area is increased by 1.36 times, rice production is increased by 4.41 times and rice yield is increased by 3.22 times during the last twenty eight years. The rice area is almost similar, but the rice production and rice yield increased significantly.




Figure 1: Area (million hectares), production (million tons) and yield (t/ha) of rice in Madhya Pradesh during 1992-93 to 2019-20

3.1 Compound Annual Growth Rate

The district-wise growth rate of area, production and yield of rice in Madhya Pradesh has been presented in Table 1. It revealed that the growth rate ranges from -19.31 to 39.40 across the districts for area, production and yield during the studied four periods.

During the period I(1992-93 to 1999-2000),the highest growth rate for area (12.74) was observed in Gwalior district and for production (20.53) and yield (18.47) was recorded in Datia district. The lowest growth rate for area (-4.62) and production (-5.74) was observed in Sehore district and lowest yield (-1.69) was recorded in recorded in Tikamgarh district.The growth rate for the state was positive for all area, production and yield.

During period II(2000-01 to 2009-10),the highest growth rate for the area (19.57) and production (24.11) was recorded inRaisen district while highest growth rate for yield(5.37) was observed inChhindwaradistrict.The lowest growth rate for all area (-19.31) and production(-19.04) was found in Bhind district and for yield was found in Tikamgarh district. In this period,the growth rate for the state was observed to be negative for area, and positive for production and yield.

During period III (2010-11 to 2019-20), the highest growth rate for area (28.22), production (39.40) and yield (16.86) was observed in Hosangabad, Raisen and Rewa districts respectively. The lowest growth rate for the area (-15.62) and production (-15.43) was recorded in Tikamgarh district while lowest growth rate for yield (-1.02) was observed in Chhindwara district. In this period, the growth rate for production and yield was observed to be positive for all the districts except two districts; and for the state it was positive for all area, production and yield.

During the overall period IV (1992-93 to 2019-20), the highest growth rate for the area (18.64) and production (22.45) was recorded in Datia and highest growth rate for yield (6.34) was observed in Sehore district. The lowest growth rate for the area (-5.94), production (-8.40) and yield (-2.62) was observed in Tikamgarh district. The growth rate yield was observed to be positive for all the districts except two districts; and for the state it was positive for all area, production and yield.

In most of the cases, the highest and lowest growth rate was observed in same districts. A similar trend was observed in Punjab [8] and Telangana [9]. The lowest growth rate was observed in Tikamgarh, Bhind, Sehore and highest growth rate was observed in Datia, Raisen districts.

3.2 Cuddy-Della Valle Instability Index

The level of instability cannot be detected by focusing only on growth rates. The growth rate will only explain the rate of growth over time, whereas instability will determine whether the growth performance for the variable under study was stable or unstable over time. In this study, the level of instability in area, production and yield of rice was determined by using Cuddy-Della Valle Instability Index.

Table 1: District-wise growth rate of area, production and yield of rice in Madhya Pradesh for various periods

| District | Area | | | | Production | | | | Yield | | | |
|-------------|-------|--------|--------|-------|------------|--------|--------|-------|-------|-------|-------|-------|
| | P-I | P-II | P-III | P-IV | P-I | P-II | P-III | P-IV | P-I | P-II | P-III | P-IV |
| Jabalpur | 1.93 | -0.41 | 6.49 | 1.35 | 2.04 | -1.02 | 19.12 | 6.16 | 0.11 | -0.62 | 11.87 | 4.75 |
| Balaghat | 0.16 | 0.04 | 1.07 | 0.40 | 0.58 | 1.79 | 6.77 | 2.86 | 0.42 | 1.74 | 5.64 | 2.45 |
| Chhindwara | 1.40 | -2.74 | 5.74 | 0.40 | 0.88 | 2.47 | 4.66 | 4.64 | -0.52 | 5.37 | -1.02 | 4.23 |
| Seoni | - | 0.72 | 3.95 | 2.08 | - | 4.77 | 9.49 | 7.86 | - | 4.02 | 5.33 | 5.66 |
| Mandla | 1.50 | 0.95 | 2.80 | 1.36 | 3.66 | 1.96 | 18.03 | 5.27 | 2.13 | 1.01 | 14.82 | 3.86 |
| Narsinghpur | -4.28 | 3.14 | 20.92 | 5.45 | -3.38 | 2.99 | 34.48 | 9.27 | 0.95 | -0.14 | 11.21 | 3.62 |
| Sagar | -3.68 | -3.75 | 0.45 | -2.53 | -2.38 | -4.32 | 6.24 | -1.17 | 1.35 | -0.59 | 5.76 | 1.40 |
| Damoh | -0.12 | 0.21 | 5.31 | 1.11 | -0.84 | 1.55 | 12.64 | 4.23 | -0.72 | 1.34 | 6.96 | 3.08 |
| Panna | 0.65 | -1.65 | 1.77 | 0.25 | 1.22 | -0.37 | 13.29 | 4.62 | 0.57 | 1.30 | 11.33 | 4.36 |
| Tikamgarh | -2.94 | -6.72 | -15.62 | -5.94 | -4.57 | -14.49 | -15.43 | -8.40 | -1.69 | -8.34 | 0.22 | -2.62 |
| Chhatarpur | 0.04 | -6.03 | -6.95 | -5.47 | -0.72 | -10.63 | -5.49 | -6.11 | -0.76 | -4.89 | 1.57 | -0.68 |
| Rewa | 1.01 | -0.22 | 4.60 | 0.52 | 3.57 | -0.21 | 22.24 | 5.79 | 2.54 | 0.01 | 16.86 | 5.25 |
| Sidhi | 2.06 | 0.56 | 2.39 | 0.78 | 3.25 | 2.00 | 16.67 | 6.83 | 1.17 | 1.43 | 13.95 | 6.00 |
| Satna | 0.25 | -2.14 | 8.11 | -0.46 | 2.91 | -2.98 | 21.53 | 5.26 | 2.66 | -0.86 | 12.41 | 5.74 |
| Sahadol | 1.25 | 0.17 | 1.92 | 0.51 | 1.20 | 0.96 | 10.10 | 4.43 | -0.05 | 0.79 | 8.02 | 3.91 |
| Jhabua | -0.99 | -0.95 | -5.90 | -2.69 | 2.62 | 2.85 | 2.13 | -0.09 | 3.64 | 3.84 | 8.53 | 2.68 |
| Morena | 6.14 | -2.59 | 16.00 | 8.85 | 10.43 | 1.30 | 17.52 | 11.71 | 4.04 | 4.00 | 1.30 | 2.63 |
| Bhind | 5.12 | -19.31 | 10.37 | -3.00 | 4.37 | -19.04 | 14.70 | -0.33 | -0.72 | 0.34 | 3.93 | 2.75 |
| Gwalior | 12.74 | -8.50 | 7.32 | 3.79 | 12.46 | -7.03 | 6.87 | 4.87 | -0.25 | 1.61 | -0.42 | 1.04 |
| Shivpuri | 1.62 | 1.20 | 10.89 | 5.67 | 4.36 | 1.31 | 18.33 | 8.90 | 2.70 | 0.11 | 6.71 | 3.06 |
| Datia | 1.74 | 6.59 | 25.00 | 18.64 | 20.53 | 5.70 | 28.92 | 22.45 | 18.47 | -0.84 | 3.14 | 3.21 |
| Sehore | -4.62 | 6.33 | 21.18 | 12.47 | -5.74 | 7.53 | 34.55 | 19.60 | -1.17 | 1.13 | 11.03 | 6.34 |
| Raisen | -2.60 | 19.57 | 27.12 | 16.26 | -1.53 | 24.11 | 39.40 | 21.22 | 1.10 | 3.80 | 9.66 | 4.26 |
| Betul | -0.89 | 1.55 | -1.27 | 1.62 | 3.49 | 3.82 | 1.27 | 5.16 | 4.41 | 2.23 | 2.58 | 3.49 |
| Hosangabad | -1.08 | 6.71 | 28.22 | 10.42 | 0.24 | 9.19 | 28.81 | 14.45 | 1.33 | 2.32 | 0.46 | 3.65 |
| State | 1.10 | -0.80 | 4.23 | 1.31 | 2.42 | 0.90 | 12.82 | 5.03 | 1.31 | 1.71 | 8.24 | 3.67 |

P-I: Period I (1992-93 to 1999-2000); P-II: Period II (2000-01 to 2009-10); P-III: Period III (2010-11 to 2019-20);

P-IV: Period IV (1992-93 to 2019-20)

Table 2: District-wise instability of area, production and yield of rice in Madhya Pradesh for various periods

| District | Area | | | | Production | | | | Yield | | | |
|-------------|-------|-------|-------|--------|------------|-------|-------|--------|-------|-------|-------|-------|
| | P-I | P-II | P-III | P-IV | P-I | P-II | P-III | P-IV | P-I | P-II | P-III | P-IV |
| Jabalpur | 1.39 | 7.40 | 17.07 | 20.07 | 21.99 | 33.97 | 35.18 | 66.68 | 21.98 | 28.72 | 27.29 | 41.19 |
| Balaghat | 1.64 | 3.33 | 11.45 | 8.04 | 12.17 | 21.13 | 20.62 | 25.06 | 11.36 | 19.87 | 13.37 | 17.87 |
| Chhindwara | 8.77 | 3.09 | 43.51 | 26.20 | 24.24 | 21.34 | 38.88 | 46.72 | 18.98 | 18.68 | 23.57 | 44.08 |
| Seoni | - | 2.35 | 12.16 | 10.46 | - | 30.61 | 36.17 | 37.31 | - | 28.78 | 30.93 | 30.69 |
| Mandla | 0.71 | 3.30 | 8.82 | 8.09 | 16.46 | 21.51 | 25.80 | 49.04 | 15.97 | 19.66 | 32.77 | 43.71 |
| Narsinghpur | 10.53 | 5.99 | 24.55 | 56.93 | 18.76 | 18.36 | 49.34 | 101.72 | 11.71 | 15.22 | 43.80 | 44.49 |
| Sagar | 4.99 | 7.33 | 15.42 | 13.39 | 15.34 | 17.70 | 40.20 | 35.06 | 13.36 | 13.16 | 34.87 | 33.80 |
| Damoh | 0.31 | 2.38 | 6.57 | 10.96 | 14.09 | 23.84 | 29.05 | 41.89 | 13.99 | 23.50 | 21.77 | 29.47 |
| Panna | 3.23 | 5.43 | 14.55 | 10.89 | 18.44 | 26.56 | 46.45 | 59.27 | 17.60 | 24.78 | 47.34 | 54.74 |
| Tikamgarh | 3.40 | 9.53 | 24.71 | 16.33 | 24.78 | 40.87 | 50.41 | 35.41 | 22.89 | 35.32 | 50.24 | 36.93 |
| Chhatarpur | 1.75 | 6.81 | 36.33 | 13.59 | 23.79 | 35.19 | 58.80 | 36.05 | 23.49 | 30.01 | 37.92 | 33.84 |
| Rewa | 3.21 | 4.14 | 24.64 | 21.56 | 25.32 | 29.04 | 36.18 | 69.16 | 25.57 | 25.88 | 51.56 | 63.75 |
| Sidhi | 2.91 | 3.06 | 14.52 | 12.36 | 23.94 | 26.15 | 30.59 | 52.72 | 22.28 | 24.89 | 27.11 | 41.14 |
| Satna | 7.13 | 4.91 | 12.74 | 22.86 | 29.44 | 46.57 | 17.07 | 57.91 | 25.50 | 41.70 | 20.59 | 40.41 |
| Sahadol | 1.00 | 2.56 | 10.94 | 8.49 | 17.03 | 23.40 | 21.82 | 35.50 | 16.50 | 22.37 | 17.12 | 29.40 |
| Jhabua | 2.80 | 3.32 | 9.20 | 7.49 | 40.98 | 34.81 | 37.76 | 42.40 | 37.22 | 33.62 | 27.60 | 38.45 |
| Morena | 40.49 | 43.85 | 44.16 | 83.29 | 56.43 | 47.04 | 53.14 | 101.05 | 14.87 | 30.70 | 17.85 | 23.19 |
| Bhind | 8.25 | 91.47 | 66.94 | 72.77 | 19.47 | 95.70 | 83.15 | 101.45 | 21.19 | 23.29 | 29.49 | 28.69 |
| Gwalior | 10.44 | 47.71 | 42.89 | 50.38 | 23.91 | 60.98 | 50.60 | 61.46 | 14.76 | 25.96 | 21.08 | 20.49 |
| Shivpuri | 7.27 | 17.39 | 63.13 | 80.34 | 17.66 | 39.91 | 76.85 | 126.32 | 14.28 | 29.45 | 26.44 | 32.07 |
| Datia | 47.78 | 43.26 | 55.56 | 122.83 | 55.79 | 59.33 | 45.13 | 123.30 | 17.10 | 26.62 | 26.90 | 29.91 |
| Sehore | 7.13 | 46.12 | 22.32 | 69.29 | 26.41 | 59.28 | 40.83 | 102.48 | 24.58 | 13.52 | 30.89 | 50.55 |
| Raisen | 15.68 | 28.05 | 23.10 | 84.57 | 22.04 | 43.87 | 35.59 | 113.00 | 14.19 | 15.02 | 23.54 | 36.25 |
| Betul | 9.99 | 2.79 | 19.72 | 17.74 | 16.12 | 12.35 | 32.67 | 39.16 | 12.20 | 11.81 | 22.69 | 29.67 |
| Hosangabad | 5.80 | 5.76 | 18.31 | 73.48 | 15.12 | 15.41 | 24.16 | 86.45 | 13.52 | 11.22 | 54.70 | 50.92 |
| State | 3.77 | 3.78 | 8.23 | 9.00 | 14.43 | 19.44 | 5.93 | 30.70 | 11.28 | 17.07 | 6.29 | 22.57 |

P-I: Period I (1992-93 to 1999-2000); P-II: Period II (2000-01 to 2009-10); P-III: Period III (2010-11 to 2019-20);

P-IV: Period IV (1992-93 to 2019-20)

The Cuddy-Della Valle Index for area, production and yield of rice in Madhya Pradesh has been presented in Table 2. During period I (1992-93 to 1999-2000), the highest instability for the area (47.78), production (56.43) and yield (37.22) was found in Datia, Morena and Jhabua districts respectively whereas the lowest instability for the area (0.31) was observed in Damoh district, and for production (12.17) and yield (11.36) was observed Balaghat district.

During period II (2000-01 to 2009-10), the highest instability for the area (91.47) and production (95.70) was observed in Bhind district and for yield (41.70) observed in Satna district while the lowest instability for area (2.35), production (12.35) and yield (11.22) was observed in Seoni, Betul and Hosangabad districts respectively.

During period III(2010-11 to 2019-20), the highest instability for the area (66.94) and production (83.15) was observed in Bhindistrict and for yield (54.70) recorded in Hosangabad district whereas the lowest instability for area (6.57), production (17.07) and yield (13.37) was recorded inDamoh, Satna and Balaghatdistricts respectively.

During the overall period IV(2092-93 to 2019-20), the highest instability for the area (122.83), production (126.32) and yield (63.75) was observed in Datia, Shivpuri and Rewa districts respectively. The lowest instability for area (7.49) was recorded in Jhabua district whereas for production (25.06) and yield (17.87) it was recorded in Balaghat district.

The instability varies from 0.31 to 122.83 per cent for the area, 12.17 to 126.32 per cent for production and 11.22 to 63.75 per cent for yield across districts during the four periods. The instability for the state for area, production and yield vary from 3.77 to 9.00, 5.93 to 30.70 and 6.29 to 22.57per cent respectively.

The instability for area of the state as a whole is classified as very low and low instability; and for the production is classified as low instability and medium instability. Similar trend was observed in Jambhulkar et al [11].

3.3 Classification of Instability Index

All districts of Madhya Pradesh have been classified as very low instability, low instability, medium instability, high instability and very high instability based on the instability value of the area, production and yield of rice. The number of districts that fall under each class has been presented in Figure 2.

For area, most of the districts fall under the class very low instability and low instability during period 1 and period II. None of the districts fall under the class very low instability during period III and period IV, very high instability during period I and high instability during period IV. In period III, most of the districts fall under the class low instability and medium instability and during period IV most of the districts fall under the class low instability and very high instability.

For production, none of the districts showed very low instability in all the periods. This is in line with Jambhulkar et al [11]. None of the districts recorded low instability in period III and IV. Most of the

districts fall under medium instability in period I and II, high instability in period III and very high instability in period IV. Very few districts recorded low instability in period I and II and medium instability in period IV. For yield, none of the districts were classified as very low instability in all the periods. Similar result was observed in Jambhulkar et al. [11]. None of the districts were classified as very high instability in period I and II; and low instability in period IV. Most of the districts were classified as medium instability in period I, II and III and high instability in period IV. Only one district recorded low instability in period III and high instability in period I.




Figure 2: Number of districts under each instability class for area, production and yield of rice during four periods in Madhya Pradesh state

P-I: Period I (1992-93 to 1999-2000); P-II: Period II (2000-01 to 2009-10); P-III: Period III (2010-11 to 2019-20); P-IV: Period IV (1992-93 to 2019-20)

4. CONCLUSION

The study revealed that, for the state as a whole growth rate for production and yield was positive in all the periods. The growth rate varies from -0.80 to 4.23 for area, 0.90 to 12.82 for production and 1.31 to 8.24 per cent for yield. For the state as a whole, for the area, production and yield highest growth rate was recorded during period III for production whereas the lowest growth rate was recorded for area during period II. For the state as a whole, the instability for area ranges from 3.77 to 9.00 for production ranged from 5.93 to 30.70 and for yield ranges from 6.29 to 22.57. Lowest instability was observed for

area in period I however lowest instability was observed for production in period IV. None of the districts recorded very low instability for production and yield. Most of the districts recorded medium instability for yield in period I, II and III for production in period I and II. This study will be useful for the researchers, policy makers and other rice stakeholders for deciding their agricultural policy and effective implementation of agricultural policy in various districts of Madhya Pradesh state. Different policies can be modified and reoriented as per the requirement and need of the rice stakeholders of Madhya Pradesh state.

REFERENCES

1. Singh DV. A component analysis and value productivity growth of important crops in Himachal Pradesh. *Agricultural Situation in India*. 1981; 36(6): 479-484.
2. Cauvey R. Groundnut production in Tamil Nadu-A decomposition analysis. *Agricultural Situation in India*. 1991; 46(5): 321-324.
3. Ranade CG. Impact of cropping pattern on agricultural production. *Indian Journal of Agricultural Economics*. 1980; 35(2): 85-92.
4. Deosthali V, Chandrashekhar MN. Rice: Region wise growth trends in Maharashtra. *Economics and Political Weekly*. 2004; 39(3): 240-242.
5. Sikka BK, Vaidya CS. Growth rates and cropping pattern changes in agriculture in Himachal Pradesh. *Agricultural Situation in India*. 1985; 39(11): 843-846.
6. Samal P, Rout C, Repalli SK, Jambhulkar NN. State-wise Analysis of Growth in Production and Profitability of Rice in India. *Indian Journal of Economics and Development*. 2018; 14(3): 399-409.
7. Jambhulkar NN, Jena SS, Mondal B, Samal P. Estimation of Growth Rate and Instability Analysis of Area, Production and Yield of Rice in Odisha State of India. *International Journal of Current Microbiology and Applied Sciences*. 2020; 9(07): 3107-3115.
8. Jambhulkar NN, Panigrahi US, Bisen J, Mondal B, Mishra SK, Kumar GAK. Growth rate and instability analysis of rice area, production and yield in Punjab. *The Pharma Innovation Journal*. 2021; 10(9) (special issues): 352-355.
9. Akul M, Bandumula N and Rathod S. Rice production in Telangana: growth, instability and decomposition analysis. *Oryza*. 2022; 59(2): 232-240.
10. Jambhulkar NN, Mondal B, Bisen J, Mishra SK, Pradhan AK. Growth and instability analysis of rice production: A district level assessment in Uttar Pradesh and Uttarakhand states of India. *The Pharma Innovation Journal*. 2023a; 12(2): 813-819.
11. Jambhulkar NN, Mondal B, Pradhan AK. Growth and Instability in Rice production: A District Level Analysis of Bihar and Jharkhand States of India. *Wulfenia*. 2023b; 30(5): 20-31.
12. Jambhulkar NN, Mondal B, Bisen JP, Pradhan AK. Growth and instability of rice production: a district level analysis in West Bengal, India. *Oryza*. 2023c; 60(3): 487-494.

13. Kumar R, Jaulkar AM, Srivastava SC, Singh S. A study on growth and instability of Paddy and Wheat crops in Gwalior District (Madhya Pradesh). *The Pharma Innovation Journal*. 2021; SP-10(5): 745-748.
14. Mishra S, Balai HK, Malviya PK. Growth and Instability of Major *Kharif* Pulse crops in Madhya Pradesh State of India. *Agro Economist*. 2022; 9(04): 251-257.
15. Balai HK, Meena PC, Meena ML, Bairwa KC, Kumar S, Meena RP, Bairwa SK. Growth, Decomposition and Instability Analysis of Major Rabi Pulse Crops in Madhya Pradesh, India. *Asian Journal of Agricultural Extension, Economics & Sociology*. 2024; 42(7):79-88.
16. Dandekar VM. Introduction seminar on data and methodology for the study of growth rates in agriculture. *Indian Journal of Agricultural Economics*. 1980; 35(2): 1-12.
17. Minhas BS. Rapporteur's report on measurement of agricultural growth. *Indian Journal of Agricultural Economics*. 1966; 21(4): 165-182.
18. Singh IJ, Rai KN. Regional variations in agricultural performance in India. *Indian Journal of Agricultural Economics*. 1997; 52(3): 374-377.
19. Cuddy JDA, Della Valle PA. Measuring the instability of time series data. *Oxford Bulletins of Economics and Statistics*. 1978; 40(10): 79-84.