

# Synthesis, Characterization and Antimicrobial Activities of Pyrrolidin-2-ylidene-2-(4-chlorophenyl)semicarbazone and its Cd(II) complex

## Abstract

A novel Schiff base ligand has been prepared by condensation between 4-chlorophenylsemicarbazide and 2-pyrrolidone in the presence of glacial acetic acid. The ligand and its Cd(II) complex were characterized using GC-MS, UV-Vis., IR, XRD, conductivity, melting point measurements and magnetic studies. The results of conductivity tests revealed that Cd(II) complex is good electrolyte while the ligand exists as a neutral molecule. Single crystal x-ray studies revealed that the ligand and its Cd(II) complex are pure crystals with respective space group symmetry of  $P3_121$  and  $P6_3/mmc$ . The Cd(II) complex of the ligand is diamagnetic with square pyramidal geometry. The emergence of infra red absorption bands at  $427$  and  $411\text{cm}^{-1}$  in the complex revealed that the ligand is bidentate and coordinates the metal ion through pyrrolidone nitrogen and the carbonyl oxygen. Antimicrobial studies on the ligand and its Cd(II) complex revealed that the free ligand showed less inhibitory activities than the metal complex against gram positive (*S. aureus*, *B. subtilis*), gram negative (*E. coli*, *K. pneumonia*) bacteria and fungi (*C. albicans*, *A. flavus*).

**Keywords:** Schiff bases, Metal complexes, pyrrolidone moiety, Ant-bacterial activities, Ant-fungal activities.

## 1. Introduction

Semicarbazones, a class of Schiff bases with the presence of imine group which impart biological activity, have proved the efficiency and efficacy in combating various diseases (Singh *et al.*, 2010). It is of great interest because of their chemistry and potentially beneficial biological activities such as anticonvulsant, antiarrhythmic (Wu *et al.*, 2006), uterotrophic (Wafaet *et al.*, 1992), antiviral (Yang *et al.*, 2010), antimalarial (Pingaewet *et al.*, 2010), antitubercular, cytotoxic, antibacterial and antifungal (Pandeya *et al.*, 2000) activities. Recently, growing consideration have been paid to the preparation of N-heterocyclic compounds because of their beneficial pharmacological properties and their role as main intermediate and fine chemicals in pharmacological and organic chemistry (Zhang *et al.*, 2012) and are proper forerunner for unusual  $\beta$ -amino acids such as statin and its analogues. Many pyrrolidinones are the construction blocks of abundant natural products such as Pyrrocidine A, B (He *et al.*, 2002), Ypaoamide (Chen *et al.*, 2009), Lactacystin (Omura *et al.*, 2003), (-)- Azaspirene (Asamiet *et al.*, 2002) that is a new inhibitor of angiogenesis which was isolated from the fungus *Neosartorya* sp. Oteromycin, as a white powder, and also from two various strains of an unidentified fungus. Functionalized dihydro-2-oxypyrrroles ring available in some of the alkaloids possesses broad pharmaceutical effect and they are also utilized as optoelectronic materials (Wiedeman *et al.*, 2003), HIV integrase inhibitors (Kawasujiet *et al.*, 2017), receptors of vascular endothelial growth factor (Peifer *et al.*, 2008), antibacterial and antifungal, nootropic agents (Adam *et al.*, 2004), peptidomimetic (Raghuraman *et al.*, 2011). Cotinine is a common main 2-pyrrolidinone, an alkaloid reported in tobacco and is also the main metabolite of nicotine (Dwoskin *et al.*, 1999). Ethosuximide is a succinimide anticonvulsant, utilized principally in the absence of seizures (Patsalos, 2005). This work is designed to explore the synthesis, characterization and

antimicrobial activities of 4-chlorophenylsemicarbazone having pyrrolidone moiety and some metal complexes.

## 2. Experimental

All reagents used in this work are of analytical grade and were obtained from directly from Aldrich Chemical Co. Inc. The reagents were used as starting materials for synthesis of more complex compounds without further purification. Metal salts were obtained from E. Merk and distilled water that was purchased from Chifok Scientific Co. (Nig.). Culture broths, Muller Hinton Agar, yeast, mould were obtained from Microbiology Department of Nnamdi Azikiwe University, Awka, Nigeria. Melting points were determined using capillary tube, Molar conductivity was measured in deionized water at 25°C using a WTW conductivity meter. The infra red (FTIR) spectra were recorded using FTIR.8300 Shimadzu Spectrophotometer using CsI disc in the frequency range of 4000-400cm<sup>-1</sup>. The ultra violet-visible spectra of the ligand and complexes will be recorded by using ultrospec. 2100 pro. ultra violet Spectrophotometer in the range of 200-800nm. Mass Spectra were recorded using gas chromatography-mass spectrophotometer (QP2010 plus, shimadzu), crystallinity was measured using x-ray diffractometer (PANalytical X' pert PRO MPD), Magnetic susceptibility was measured using Sherwood scientific magnetic moment balance (MK1 model) and weights were measured using digital balance (Mettler Toledo PL 203).


### 2.1 Synthesis of pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L)

The ligand was synthesised according to the method used by Singh *et al.*, (2009) as follows:

**Step1:** 4-chloroaniline (0.1M, 1.2757g) was dissolved in 10cm<sup>3</sup> of glacial acetic acid and diluted to 100cm<sup>3</sup> with distilled water. Equimolar quantity (0.1M, 0.33g) of sodium cyanate in 50cm<sup>3</sup> of warm water will be added in the previous solution with stirring. The reaction mixture is allowed to stand for 30minutes, and crystals formed were filtered, washed, dried and recrystallized from boiling water.

**Step2:** 60g of 4-chlorophenylurea in 100ml of hydrazine hydrate refluxed for 3hrs. Vacuum distillate at 70°C, the product remaining in the flask is digested with 250ml methanol and refluxed for 45minutes, cooled room temperature and filter. The filtrate obtained is cooled in ice bath to 4-chlorophenylsemicarbazide which was further purified by recrystallization from ethanol.

**Step3:** A solution of the 4-chlorophenylsemicarbazide (0.1M, 2.22g) and equimolar quantity of pyrrolidone (0.1M) in 100cm<sup>3</sup> of ethanol will be refluxed at 70°C for 2hours in the presence of glacial acetic acid (1cm<sup>3</sup>). The product obtained after cooling was filtered and recrystallized from 95% ethanol to give pure Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) ligand. Yield: 51%, M.p.: 198°C, C<sub>11</sub>H<sub>13</sub>ON<sub>4</sub>Cl(L), Anal. Found: C, 51.50; H, 5.07; O, 6.24; N, 21.85; Cl, 13.85%. Calc. C, 52.23; H, 5.15; O, 6.34; N, 22.18; Cl, 14.06% IR (KBr, v/cm<sup>-1</sup>): v(C=O) 1650; v(C=N) 1540; v(N4-H) 3413; v(N1-H) 3310; v(C-N) 1248; v(C-Cl) 817. UV-Vis.(DMSO) λ<sub>max</sub>/cm<sup>-1</sup> 36,765, 31,447. XRD: cell dimension(Å) 4.91, 4.91, 5.43. interfacial angles (°) 90, 90, 120. Volume of unit cell (Å<sup>3</sup>), 130.907. GC-MS (m/z): cal., 252.5; found, 256.3




**Scheme 1: Synthesis of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone(L)**

### 2.2 Synthesis of Cd(II) Complex of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone(L)

Cadmium (II) Complex of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone were synthesized according to the method used by Yousif *et al.*, (2013) as follows: 50cm<sup>3</sup> of 0.25M ethanolic solution of Cadmium (II) trioxonitrate (V) tetrahydrate [Cd(NO<sub>3</sub>)<sub>2</sub>.4H<sub>2</sub>O]. was added to 50cm<sup>3</sup> of 0.5M ethanolic solution of pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone and heated under reflux at 70°C for 1 hour and cooled in ice. The product was washed with distilled water and recrystallized from 95% ethanol. Yield: 47%, M.p.: 260°C, CdL(NO<sub>3</sub>)<sub>3</sub>: Anal. Found: C, 26.98; H, 2.66; O, 32.71; N, 20.03; Cl, 7.26; Cd, 10.36%. calc. C, 27.00; H, 2.66; O, 32.73; N, 20.03; Cl, 7.26, Cd, 10.31% IR (KBr, v/cm<sup>-1</sup>): v(C=O) 1648; v(C=N) 1541; v(N<sup>4</sup>-H) 3413; v(N<sup>1</sup>-H) 3309; v(C-N) 1248; v(C-Cl) 817, v(Cd-N), 427; v(Cd-O), 411. UV-Vis.(DMSO) λ<sub>max</sub>/cm<sup>-1</sup>: 34,013. XRD: cell dimension (Å) 5.98, 5.98, 9.76. interfacial angles (°) 90, 90, 120.

Volume of unit cell ( $\text{\AA}^3$ ), 349.022. conductivity( $\mu\text{s/cm}$ ), 200; effective magnetic moment, diamagnetic. GC-MS (m/z): cal., 488.9; found, 489.2.



**Scheme 2: Synthesis of Cd(II) Complex of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone(L)**

### 3. Results and Discussion

The reaction of 4-chlorophenylsemicarbazide with 2-pyrrolidone in the presence of glacial acetic acid yielded the ligand (L); Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (Scheme1). The reaction of the ligand (L); Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone with cadmium trioxonitrate (V) tetrahydrate  $[\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}]$  in 2:1 ligand to metal ratio which yielded the corresponding cadmium (II) complex,  $\text{CdL}(\text{NO}_3)_3$ . The ligand and its cadmium (II) complex were isolated as air stable microcrystalline solids in good percentage yield of 51 and 47% respectively (Table 1). While the white colouration of the ligand is seen to result from less conjugated nature of the compound, white colour of the complex is attributed to completely filled d-orbital ( $d^{10}$ ) configuration in the complex. High molar conductivity value of the complex suggests that it can function as a good electrolyte, while the ligand showed no signal on conductivity. This signals the presence of metal ions in the complex which is indicative of complex formation, while the ligand on the basis of its conductivity value could be regarded as neutral molecule. The melting points of the ligand and complexes suggest that the synthesized compounds are air and moisture stable (Table 1). Observed increase in solubility in water and increase in the melting point of the ligand from 198 to  $260^\circ\text{C}$  upon complexation with Cd(II) ion is indicative of metal-ligand bond formation (Wissner *et al.*, 2000). The complex is a monometallic centred compound. This is deduced from the results of percentage elemental analysis which are in good agreement with the assigned formulations. The theoretically calculated percentage values of elements in the ligand and in complexes are in close agreement with experimental values obtained from the results of gas chromatography-mass spectroscopic analyses.


**Table 1: Physicochemical Properties pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cadmium (II) Complex**

S/ N	Compounds	Colours	Meltin g Points (°C)	Yiel d (%)	Mola r mass g/mol	% Element found (% calculated)					
						%C	%H	%O	%N	%CL	%M
1	C <sub>11</sub> H <sub>13</sub> ON <sub>4</sub> Cl(L)	White	198	51	256.3 (252.5)	51.50 (52.23)	5.07 (5.15)	6.24 (6.34)	21.85 (22.18)	1385 (14.06)	- -
2.	CdL(H <sub>2</sub> O) <sub>3</sub> NO <sub>3</sub>	White	260	41	489.2 (488.9)	26.98 (27.00)	2.66 (2.66)	32.71 (32.73)	20.03 (20.04)	7.26 (7.26)	10.36 (10.31)


The infra red spectra of the ligand are almost the same in the regions of 3413cm<sup>-1</sup>-1540cm<sup>-1</sup> to those of the corresponding metal complexes. This suggests that the molecular functional groups in both the ligand and the complexes are identical. In Cd(II) complex, the most feasible coordination option is bonding through N<sup>1</sup>-H and C=O. This is suggested by variations in the absorption frequencies in the complex (N<sup>1</sup>-H at 3309cm<sup>-1</sup>, C=O at 1648cm<sup>-1</sup>) compared to that of the ligand (N<sup>1</sup>-H at 3310cm<sup>-1</sup>, C=O at 1650cm<sup>-1</sup>). It is also inferred from the infra red results that approach of the ligand was electrostatic (Ali *et al.*, 2019), thereby compressing the crystal structures of the complexes directly through N<sup>1</sup>-H and C=O axes, shortening the bond length, strengthening and increasing the bond energies of N<sup>1</sup>-H and C=O bonds leading to the observed increase in the absorption frequencies as compared the that of the free ligand.

**Table 2 : Selected Infra Red Data of pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cadmium (II) Complex**

S/N	Compounds	(N <sup>1</sup> -H)	(N <sup>4</sup> -H)	(C=O)	(C=N)	(M-N)	(M-O)
1.	C <sub>11</sub> H <sub>13</sub> ON <sub>4</sub> Cl(L)	3413	3310	1650	1540	-	-
2.	CuL(H <sub>2</sub> O) <sub>3</sub> NO <sub>3</sub>	3413	3309	1648	1541	427	411




**Fig. 1: Infra Red Spectrum of pyrrolidin-2-ylidene-2-(4-chlorophenyl)semicarbazone (L)**



**Fig.2: Cadmium (II) Complex of pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L)**


Gas chromatogram of the ligand showed peaks at  $m/z$  429.2, 355.1, 284.3 and 252.5, with at  $m/z$  429.2 result from the interaction between the already formed ligand with 2-pyrrolidone dimer present in the reaction medium. This interaction was necessitated by high tendency of

intermolecular hydrogen bond formation between the ligand and coupled 2-pyrrolidone molecules (Yekeler, 2001). Upon fragmentation, the two pyrrole rings that make up the 2-pyrrolidone dimer are lost gradually in two step mechanisms. At first, one pyrrole ring is detached leaving behind a cation with  $m/z$  355.1, in the second step, another pyrrole ring fragments out resulting to a cation with  $m/z$  284.3. The fragment with  $m/z$  284.3 then loses an hydroxyl ion and an atom of oxygen to give the ligand peak  $m/z$  256.3. All the observed fragments possess  $m/z$  values are in agreement with the calculated values confirming the ligand formation (Fig.3).




**Figure 3: GC-MS Spectrum of The Ligand**

While for Cd(II) complex of the ligand,  $\text{CdL}(\text{NO}_3)_3$ , the observed  $M^+$  is  $m/z$  489.2 and the calculated molecular mass is  $488.9 \text{ g mol}^{-1}$ , this shows total agreement between the observed and the calculated molecular masses for the complex.



**Figure 4: GC-MS Spectrum of Cd(II) Complex of The Ligand**

Sharp peaks obtained from the single crystal x-ray analysis of the compounds confirmed that both the ligand and its Cd (II) complex exist as pure crystals. The spectra obtained in this work were interpreted by matching with standard libraries and similar materials in the literature (Liang and Hawthorne, 1996). The ligand was identified to be of  $p3_121$  symmetry and in trigonal crystal system while Cd (II) complexes of the ligand possessed a symmetry of  $p63/mmc$  and hexagonal in shape.



**Fig.5: XRD Spectra of The Ligand (A) and its Cd(II) Complex(B)**

**Table 3: XRD Data of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Copper(II) Complex**

Parameters	C <sub>11</sub> H <sub>13</sub> ON <sub>4</sub> Cl(L)	CdL(NO <sub>3</sub> ) <sub>3</sub>
Empirical formula	C <sub>11</sub> H <sub>13</sub> N <sub>4</sub> OCl	CdC <sub>11</sub> H <sub>13</sub> O <sub>10</sub> N <sub>7</sub> Cl
Formula weight (g/mol)	252.5	489.2
Temperature (K)	298	298
Wavelength(Mo K $\alpha$ )( Å)	0.71073	0.71073
Crystal system	Trigonal	Hexagonal
Space group	P3 <sub>1</sub> 21	P6 <sub>3</sub> /mmc
Lattice constant		
a(Å)	4.91	5.98
b(Å)	4.91	5.98
c(Å)	5.43	9.76
$\alpha$ (°)	90	90
$\beta$ (°)	90	90
$\gamma$ (°)	120	120
Volume(Å <sup>3</sup> )	130.907	349.022


Spectra of the ligand showed peaks at 36,765cm<sup>-1</sup> (272nm) and 31,447cm<sup>-1</sup> (318nm) assignable to  $\pi \rightarrow \pi^*$  and  $n \rightarrow \pi^*$  transitions respectively. The former transition band arose from the excitation of pi-electrons to the next higher energy level which is attributed to the presence of double bonds of the benzene ring that are in conjugation with lone pair of electrons of the anilinic nitrogen group, while the later transition band occurring at 31,447cm (318nm) revealed the excitation of non-bonding electrons to pi antibonding orbitals of the imine (C=N<sup>2</sup>) functional group. The observed variation in the transitions occurring in the free ligand and those of its metal complexes is indicative of metal-ligand bond formation resulting from complexation(Wissner *et al.*, 2000).High molar conductivity value of the complex suggests that it can function as a good electrolyte, while the ligand showed no signal on conductivity andcould be regarded as a neutral molecule. This is indicative of complex formation (Brown *et al.*, 1976).

**Table4: Electrical Data of Pyrrohdin-2-hydene-2-, (4-chlorophenyl) Semicarbonzone and Its Cd (II) Complex.**

S/N	Compounds	Conductivity $\mu$ S/CM	Magnetic Moment (BM)	Absorption Bands (CM <sup>-1</sup> )	Assigned Transitions	Geometry
1.	C <sub>11</sub> H <sub>13</sub> ON <sub>4</sub> Cl(L)	-	-	36,765, 31,447	$\pi \rightarrow \pi^*$ , $n \rightarrow \pi^*$	-
2.	CdL(NO <sub>3</sub> ) <sub>3</sub>	200	Diamagnetic	34,103	L→Cd(CT)	Square Pyramidal

**Proposed Structures of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Copper(II) Complex**


From the elemental analysis, observed magnetic moment and the spectral data, structures of the synthesized compounds are suggested as shown in Fig. 3 The structures assigned to these compounds are confirmed by the IR, GC-MS, XRD and UV-Vis. Spectroscopic data and by analysis of the analogous structures available in the literature (Turdo *et al.*, 2007).




**Fig.6:Proposed Structures of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cadmium(II) Complex**

#### **Antimicrobial Activities of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)-semicarbazone and its Cd(II) complex**

Pyrrolidin-2-ylidene-2,(4-chlorophenyl)-semicarbazone and its Cd(II) complex were tested in vitro growth the inhibitory activities two gram positive bacteria (*A. Aureus* and *B. Subtilis*), two gram negative bacteria (*E. coli* and *K. Pneumoniae*) and two fungi (*C. albicans* and *A.flavus*) by standard disc diffusion technique according to Bindu *et al.*, (1998) using 100% DMSO as solvent. Effectiveness of antimicrobial agent was based on the diameter of inhibition. Antimicrobial agents were considered effective when the diameter of inhibited zone is 9mm and above, but ineffective or negative when the diameter of inhibited zone is less than 9mm as was adopted from Joseph *et al.*, (2006).




**Fig.7: Antimicrobial Activities of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cd(II) Complex at 1000µg/ml Dosage**



**Fig.8: Antimicrobial Activities of Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cd(II) Complex at 500µg/ml Dosage**

The free ligand did not show appreciable inhibition activity against all bacteria tested while Cd(II) complex caused inhibition for *S.aureus*, *B. Subtilis*, *E.coli*, *K. Pneumoniae*. From the antibacterial activities data, complexes are more potent antimicrobials than the free ligand against one or more microorganisms. This is attributable to the hydrophobic nature of the Schiff base ligands, which restricts their permeation to the cells and tissues. In addition, chelation enhances biochemical potential of bioactive organic species (Nair *et al.*, 2012). Cd(II) complex showed high inhibition activity against Gram negative bacteria (*E. coli* and *K. Pneumoniae*) at both low and high doses. This suggests that apart from being a chelate, Cd(II) complex of the ligand possessed some sort of lipophilic characteristics in order to enhance membrane permeability into the outer lipopolysaccharides layers of Gram-negative bacteria (Raman *et al.*, 2009). The results of antifungal activities revealed that the ligand showed no observable activity against the test organisms (*C. albicans* and *A. flavus*), while Cd(II) complex of the ligand was only active against *A. flavus* at high dose. In the case of *A. flavus*, the observed resistance against the ligand and its Cd(II) complex (at low dose of 500mg) could be attributed to the development of biofilms which provide temporary antifungal drug resistance and protects the pathogen in the hostile environment, this is very common behavior among *Aspergillus* spp. (Paul *et al.*, 2017). The underlying mechanism of resistance in *C. albicans* as observed against the ligand complexes could be attributed to the ability of the organism to bring about alterations in drug targets, due to mutation in target which reduces binding of drug to the target (Sanglard, 2016). The high activities of Cd(II) complex against *C. albicans* and *A. flavus* over the free ligand could be attributed to the presence of metal ions, and its effect on the normal cell membrane. Metal chelates bear polar and nonpolar properties together; this makes them suitable for permeation to the cells and tissues. This is possible because the cell wall is essential to the survival of many organisms and some antibiotics are able to kill them by inhibiting a step in the synthesis of peptidoglycan (El-Sherif and Eldebss, 2011). Cd(II) complex showed positive activity at respective MIC and MBC of 62.5 and 125 against *S.aureus*, *E. coli*, *A. flavus*. In comparison with the ligand which is rated inactive against all organisms studied due to its higher MIC and MBC values, potent bactericidal behavior of these metal complex can be attributed to the fact that chelation increases solubility, conductivity and  $\pi$ -electron delocalization in metal complexes (Wissner *et al.*, 2000).



**Fig. 9:** Bacterial Activities of the ligand(A) and its Cd(II)(B) against *B. subtilis* by disc diffusion method

#### 4. Conclusion

The ligand Pyrrolidin-2-ylidene-2,(4-chlorophenyl)semicarbazone (L) and its Cd(II) complex were synthesized and fully characterized. The ligand coordinated to the Cadmium (II) ion through metal-oxygen and metal-nitrogen bond formation to afford the complex. The metal ion was five coordinated in the complex and possessed square pyramidal geometry. Antimicrobial studies on the ligand and its Cd(II) complex revealed that the free ligand showed less inhibitory activities than the metal complex against gram positive (*S. aureus*, *B. subtilis*), gram negative (*E. coli*, *K. pneumonia*) bacteria and fungi (*C. albicans*, *A. flavus*).

#### 5. References

- Adam, W., Zhang, A. (2004). High *pi*-facial selectivity through chelation of magnesium ions in the DMD epoxidation of  $\alpha$ ,  $\beta$ -unsaturated imides with chiral pyrrolidinone auxiliaries. *Eur. J. Org. Chem.* (1), 147-152.
- Ali, U., Taj, M.B., Raheel, A., Alelwani, W., Tirmizi, S.A. (2019). A Min. Review on mixed chelation Parameters. *Biomed. J Sci and Tech. Res.* 20(4), 15287-15292.
- Asami, Y., Kakeya, H., Onose, R., Yoshida, A., Matsuzaki, H., Osada, H. (2002). Azaspirene: a novel angiogenesis inhibitor containing a 1-oxa-7-azaspiro [4.4] non-2-ene-4, 6-dione skeleton produced by the fungus *Neosartorya* sp. *Org. Lett.*, 4(17), 2845-2848.

- Brown, R., Fischer, R., Berlin, K.D., Ramalingam, N.N. (1976). Biological activity and active group of novel pyrazole, thiosemicarbazone and substituted thiozoles. *Journal of Molecular Pathology of Diseases* 5(6), 15-17.
- Chen, J., Huang, P.Q., Queneau, Y. (2009). Enantioselective synthesis of the Renantiomer of the feeding deterrent (S)-Ypaoamide. *J. Org. Chem.*, 74(19), 7457-7463.
- Dwoskin, L.P., Teng, L., Buxton, S.T., Crooks, P.A. (1999). (S)-(-)-Cotinine, the major brain metabolite of nicotine, stimulates nicotinic receptors to evoke [3H] dopamine release from rat striatal slices in a calcium-dependent manner. *J. Pharmacol. Exp. Ther.*, 288(3), 905-911.
- El-Sherif, A.A., Eldebss, T.M.A. (2011). Synthesis, spectral characterization, solution equilibria, in vitro antibacterial and cytotoxic activities of Cu(II), Ni(II), Mn(II), Co(II) and Zn(II) complexes with Schiff base derived from 5-bromosalicylaldehyde and 2-aminomethylthiophene. *Spectrochim Acta*;79A:1803-14.
- He, H., Yang, H.Y., Bigelis, R., Solum, E.H., Greenstein, M., Carter, G.T. (2002). Pyrrocidines A and B, new antibiotics produced by a filamentous fungus. *Tetrahedron Lett.*43(9), 1633-1636.
- Joseph, M., Kuriakose, M., Kurup, M.R.P., Suresh, E., Kishore, A., Bhat, S.G. (2006). Structural, antimicrobial and spectral studies of copper(II) complexes of 2-benzoylpyridine N(4)-phenyl thiosemicarbazone. *Polyhedron* 25,61-70
- Kawasuji, T., Fuji, M., Yoshinaga, T., Sato, A., Fujiwara, T., Kiyama, R. (2007). 3-Hydroxy-1,5-dihydro-pyrrol-2-one derivatives as advanced inhibitors of HIV integrase. *Bioorg. Med. Chem.*, 15(16), 5487-5492.
- Patsalos, P.N. (2005) Properties of antiepileptic drugs in the treatment of idiopathic generalized epilepsies. *Epilepsia*, 46(s9), 140-148.
- Peifer, C., Selig, R., Kinkel, K., Ott, D., Totzke, F., Schachtele, C., Heidenreich, R., Röcken, M., Schollmeyer, D., Laufer, S. (2008). Design, synthesis, and biological evaluation of novel 3-aryl-4-(1H-indole-3-yl)-1,5-dihydro-2H-pyrrole-2-ones as vascular endothelial growth factor receptor (VEGF-R) inhibitors. *J. Med. Chem.*,51(13), 3814-3824.
- Pingaew, R., Prachayasittikul, S. and Ruchirawat, S. (2010). *Molecules*, 15, 988.
- Raghuraman, A., Ko, E., Perez, L.M., Ioerger, T.R., Burgess, K. (2011). Pyrrolinone-pyrrolidine oligomers as universal peptidomimetics. *JACS*,133(32), 12350-12353.
- Singh, S., Tiwari, A.K., Ojha, H., Kumar, N., Singh, B. and Mishra, A.K. (2010). *Cancel Biother, Radiopharm.*, 25, 117.
- Wafa, O.M.A., El-din, M.M.M. and Omar, A.M.M.E. (1992). *Steroid*, 57, 199.
- Wiedeman, P.E., Trevillyan, J. (2003). Dipeptidyl peptidase IV inhibitors for the treatment of impaired glucose tolerance and type 2 diabetes. *Curr. Opin. Investig. Drugs*,4(4), 412-420.
- Wissner, A., Berger, D.M., Boschelli, D.H. (2000). 4-anilino-6,7-dialkoxyquinoline-3-carbonitrile inhibitors of epidermal growth factor receptor kinase and

bisosteric relationship to the 4-anilino-6,7-dialkoxyquinoline-3-carbonitrile inhibitors. *Journal of Medicinal Chemistry*, vol.43, no.17, 3241-3256.

Wu, S., Fluxe, A., Janusz, J.M., Sheffer, J.B., Browning, G., Blass, B., Cobum, K., Hedges, R., Murawsky, M., Fang, B., Fadayeel, G.M., Hare, M. and Djandjighian, L. (2006). *Bioorg. Med. Chem. Lett.*, 16, 5859.

Yang, Y.J, Zhao, J.H, Pan, X.D. and Zhang, P.C. (2010). *Chem. Pharm. Bull.*, 58, 208.

Yekeler, H. (2001). Solvent effects on dimeric self association of 2-pyrrolidone: an ab initio study. *J Comput. Aided Mol. Des.*15(4):287-95.

Yousif, E., Majeed, A., Al-Sammarrae, K., Salih, N., Salimon, J. and Abdullah. (2013). Metal Complexes of Schiff Base: Preparation, Characterization and antibacterial activity. *Arabian Journal of Chemistry*. 06.006.

Zhang, N., Fan, Y., Zhang, Z., Zuo, J., Zhang, P., Wang, Q. (2012). Syntheses, crystal structures and anticancer activities of three novel transition metal complexes with Schiff base derived from 2-acetylpyridine and l-tryptophan. *Inorg Chem Commun*;22:68-72.

UNDER PEER REVIEW