

Original Research Article

Effect of selected botanicals and carbendazim against *Alternaria* blight disease (*Alternaria brassicae*) of mustard (*Brassica juncea* L.)

ABSTRACT

A field experiment was conducted at the research plot of the Department of Plant Pathology, SHIATS, Allahabad, U.P. during the *Rabi* season of 2023-24 to evaluate selected botanicals, and carbendazim for the management of *Alternaria* blight (*Alternaria brassicae*) of Indian mustard (*Brassica juncea* L.) by applying of foliar spray of botanical and seed treatment of carbendazim. The disease intensity(%), yield (q/ha), and cost benefit ratio were recorded. The minimum disease intensity (%) at 75 and 90 DAS was recorded of neem leaf extract + carbendazim (31.72) and (35.63) followed by lantana leaf extract+ carbendazim (33.54) and (38.72) as compared with the control (T_0). The maximum yield (q/ha) was recorded (13.07q/ha) and cost benefit (1:1.82) neem leaf extract followed by lantana leaf extract + carbendazim as compared to untreated check (T_0).

Key words: *Alternaria brassicae*, botanicals, carbendazim

1.1 INTRODUCTION

Mustard [*Brassica juncea* (L.) Czern and Cross] is important *Rabi* oilseed crop which belongs to family "Cruciferae". It is also growing in certain tropical and subtropical regions as cold weather crop. Indian mustard required annual temperature of 6 to 27°C, and pH of 4.3 to 8.3 (Kumar and Shete, 2021). The seed and oil of mustard have a peculiar pungency due to presence of glucosinolate and its hydrolysis products such as Allyl Isothiocyanate (0.30-0.35%). Mustard seed in general, contains 30- 33% oil, 17-25% proteins, 8-10% fibres, 6-10% moisture, and 10-12% extractable substances (Pandey *et al.*, 2023).

The productivity saw 7% jump from 1331 to 1419 kg/ha. The area under rapeseed & mustard increased by 28% from 68.56 in 2019-20 to 88.06 lakh ha in 2022-23 (Source: DA&FW National Conference for *Rabi* Campaign 2023-24).

Alternaria blight, sclerotinia stem rot, white rust and nutrient deficiency causing substantial yield loss. which of the following *alternaria* blight is a major concern. *Alternaria* blight disease caused by *Alternaria brassicae* have been reported from all the continents of the

Comment [H1]: Try to add general statement regarding *Alternaria* and mustard

Comment [H2]: Is it SHUATS?

Comment [H3]: Add this portion in materials and method section

Comment [H4]: Write the full form of abbreviation if used first time in sentence

Comment [H5]: Finish the abstract with line of importance of this study. Like How this study will helpful?

Comment [H6]: Add "mustard"

Comment [H7]: *Rabi*, should not italic

Comment [H8]: Full stop is needed before starting the new sentence

world affect most cruciferous crops and is one among the important diseases of rapeseed-mustard causing severe yield losses with no proven source of transferable resistance in any of the hosts.

Comment [H9]: Rephrasing of sentences needed with elaboration. Avoid repetition of sentence. Need citation. Try to add more introduction as it seems incomplete. Add aim and objective of the study

2.1 Materials and Methods

2.1.1 Isolation

Potato Dextrose Agar (PDA) was prepared and 80 mg of streptomycin, an antibiotic was added to each 500 ml preparation of the PDA to inhibit probable bacterial growth. The infected leaf parts were cut into small pieces of 2-3 mm dimension in a manner so that pieces may have some green portion also. Such leaf bits were surface sterilized with 0.1 per cent sodium hypochlorite (NaOCl) solution for 30 seconds and washed three times with sterile distilled water to remove any traces of sodium hypochlorite adhered with leaf bits (Tuite, 1969). Two to three leaf bits were transferred on PDA medium contained in petri plates aseptically with the help of sterilized forceps. These petri plates were incubated at $27\pm 2^{\circ}\text{C}$.

2.1.2 Morphological characters of *Alternariabrassicae*:

The mycelium of *Alternaria brassicae* is septate and it becomes brown to brownish grey in colour. The conidiophores are dark septate, measuring $14-74\mu \times 4-8\mu$. The structure of conidia is brownish black, singly borne or sparingly in chains with 2-4, Muriform along with beak (Ellis, 1971).

2.1.3 Preparation of plant extracts:

The fresh leaves of selected plants were gently washed under running tap water and finally in sterile distilled water. They were separately grinded in sterile water at the rate 1 ml/g of plant material in pestle and mortar. Then were filtered through double layer of muslin cloth and finally through sterilized Whatman no.1 filter paper. This forms 100 % standard plant extract solution. Further ~~its dilution~~ dilutions were performed of required concentration with sterilized water. By grinding 10 g of leaves in 100 ml of sterile water 10 % of leaf extract can be obtained.

Comment [H10]: Mention the plant names and number of plants selected for treatments

2.1.4 Per cent disease intensity

Per cent disease intensity was recorded at 45, 60, 75 and 90 days after sowing. The disease intensity was calculated using the following formula (Wheeler, 1969)

$$\text{Disease intensity (\%)} = \frac{\text{Sum all disease ratings}}{\text{Total number of ratings} \times \text{Maximum disease grade}} \times 100$$

2.1.5 Disease intensity scale

Five plants were randomly selected as grades in each plot at the interval of 15 days as per the scale given by Mayee and Datar (1986) which is given below:

List 1 : Disease intensity scale

| Grade | Leaf area infected |
|-------|--|
| 0 | No symptoms on the leaves |
| 1 | Small spots covering 1% or less area |
| 3 | Small spots (upto 5 mm in size) covering 1-10% of leaf area |
| 5 | Spots enlarging and covering 26-50% of leaf area |
| 7 | Spot coalescing to form big patches covering 26-50% of leaf area |
| 9 | Big spot covering 51% or more of leaf area |

2.1.6 Cost benefit ration

Gross return was calculated by multiplying total yield with the market price of the produce. Cost of cultivation and cost of treatment imposition will be deducted from the gross returns, to find out net returns and cost benefit ratio by following formula (Reddy and Reddi, 2004)

$$C : B \text{ ratio} = \frac{\text{Gross return}}{\text{Total cost of cultivation}}$$

3.1 Results and Discussion

The application of botanicals and seed treatment with carbendazim, such as neem leaf extract + carbendazim, lantana leaf extract + carbendazim, eucalyptus leaf extract + carbendazim, datura leaf extract + carbendazim, chenopodium leaf extract + carbendazim and mancozeb 75% WP (0.2%) + carbendazim.

The results revealed that the application of botanicals and seed treatment with carbendazim, showed minimum disease intensity (%) @ 75 DAS was recorded with neem leaf

Comment [H11]: Add this portion in materials and methods section and mention the control treatment. Also mention how to apply this treatments

extract + carbendazim (31.720%) followed by lantana leaf extract + carbendazim (33.547%) as compared to treated check mancozeb + carbendazim (26.760%) and control (47.597%). Minimum disease intensity (%) @ 90 DAS was recorded with neem leaf extract + carbendazim (35.630%) followed by lantana leaf extract + carbendazim (38.723%) as compared to treated check mancozeb + carbendazim (29.547%) and control (53.537%).

Comment [H12]: Write the percentage Up to 4 digit only

Comment [H13]: Mention control and check treatment in materials and method section

In the present study, the minimum disease intensity at 45, 60, 75 and 90 DAS was recorded with neem leaf extract + carbendazim followed by lantana leaf extract + carbendazim and found to be effective over other treatments. The probable reason for such a finding may be that neem possess fungicidal property. It might be due to the presence of azadirachtin, nimbin, and salannin that are the most important biologically active substances of neem, which exhibit antimicrobial, insecticidal, and antifungal properties. The antifungal properties of neem extract are may be attributed to its ability to inhibit the fungal growth through mechanisms disruption of fungal cell membranes, causing leakage of cellular contents. Inhibition of spore germination and mycelial growth, may reduce the spread of the pathogen and enhance the plant health. Similar findings have been reported by Shrivastava and Swarnkar (2014) and Thakur and Zacharia (2017).

Comment [H14]: Need citation

Comment [H15]: Briefly discuss the findings and correlate with study

Maximum yield (q/ha) was recorded with neem leaf extract + carbendazim (13.07q) followed by lantana leaf extract + carbendazim (12.42q) as compared to treated check mancozeb + carbendazim (14.00q) and control (9.14q). Maximum cost benefit ratio was recorded with neem leaf extract + carbendazim (1:1.82) followed by lantana leaf extract + carbendazim (1:1.74) as compared to treated check mancozeb + carbendazim (1:2.01) and control (1:1.41).

Comment [H16]: Not matching with Table data

Comment [H17]: Not matching with Table data

Comment [H18]: Not matching with Table data

Comment [H19]: Not matching with Table data

In this study, the increases in yield (q/ha) could potentially be attributed to the presence of neem leaf extract + carbendazim followed by lantana leaf extract + carbendazim and found to be effective over other treatments. It may have released growth-regulating substances and aided in the control of pathogen, also leading to the observed enhancement of yield. The antifungal properties of neem have suppressed fungal growth and increased in yield. Similar findings have been reported by Sailaja *et al.* (2017), Sharma *et al.* (2021) and Ravella *et al.* (2023).

Comment [H20]: Need to discuss the findings

Table 1 Effect of selected treatments on yield (q/ha) and per cent disease intensity (%) of alternaria leaf blight at 45, 60, 75, and 90 DAS

| Tr. No. | Treatment | Per cent disease intensity (%) | | | | Yield (q/ha) | C:B ratio |
|----------------|--|--------------------------------|--------|--------|--------|--------------|-----------|
| | | 45 DAS | 60 DAS | 75 DAS | 90 DAS | | |
| T ₀ | Control | 21.613 | 35.410 | 47.597 | 53.537 | 9.14 | 1:1.43 |
| T ₁ | Neem leaf extract @10% (F.S.) + Carbendazim 2g (S.T.) | 16.897 | 23.503 | 31.720 | 35.630 | 13.07 | 1:1.85 |
| T ₂ | Lantana leaf extract @10% (F.S.) + Carbendazim 2g (S.T.) | 18.360 | 25.100 | 33.547 | 38.723 | 12.42 | 1:1.76 |
| T ₃ | Eucalyptus leaf extract @10% (F.S.) + Carbendazim 2g (S.T.) | 18.620 | 30.820 | 38.340 | 43.263 | 10.76 | 1:1.53 |
| T ₄ | Datura leaf extract @ 10% (F.S.) + Carbendazim 2g (S.T.) | 19.413 | 31.160 | 37.820 | 42.697 | 11.10 | 1:1.58 |
| T ₅ | Chenopodium leaf extract @ 10% (F.S.) + Carbendazim 2g (S.T.) | 20.420 | 32.470 | 41.733 | 47.533 | 11.70 | 1:1.66 |
| T ₆ | Mancozeb 75 % WP (0.2%) (F.S.) + Carbendazim 2g (S.T.) | 15.563 | 21.603 | 26.760 | 29.547 | 14.00 | 1:2.04 |
| | C.D. (P=0.05%) | 0.554 | 0.538 | 0.526 | 0.588 | 0.498 | |
| | C.V. (%) | 1.664 | 1.058 | 0.804 | 0.795 | 2.386 | |

*F.S. Foliar Spray and *S.T. Seed Treatment




Figure1. Effect of selected treatments on yield (q/ha) and percent disease intensity (%) of alternaria leaf blight at 45, 60, 75, and 90DAS

Comment [H21]: Remove axis Title from graph

4.1 CONCLUSIONS

This study “Effect of selected botanical and carbendazim against *Alternaria* blight disease (*Alternaria brassicae*) of mustard (*Brassica juncea* L.)”. found that plant extracts can help manage the disease *alternaria* blight in mustard plants. It was found that botanical neem leaf extract significantly managed *Alternaria brassicae* which causes *Alternaria* blight disease on mustard. The findings also indicate that neem leaf extract (Foliar Spray) + carbendazim (Seed Treatment) to observe the minimum disease intensity (%), maximum plant height (cm), yield (q/ha) and cost benefit ratio. The study concludes that plant extracts play a key role to manage *alternaria* blight in mustard. This experiment proves that farmers can manage *alternaria* blight in mustard without chemicals by using various plant extracts. However, these findings are from just one growing season. To confirm these results, researchers should run more trials in the future.

Comment [H22]: Plant height data not given in manuscript

5.1 REFERENCES

- Ellis, M. B. (1971). *Dematiaceous Hyphomycetes*. Kew, UK; Commonwealth Mycology Institute.
- Kumar, R. and Shete, P. (2021). Symptomology, biology and management of *alternaria* leaf spot of mustard (*Brassica* spp.). *The Pharma Innovation Journal*. 10(6): 264-268.

Mayee, C. D. and Datar, V. V. (1986). Phyto-pathometry. tech. bulletin. Marathwada Agricultural University Parbhani, Maharashtra, India.(special bulletin 3), p: 95

Pandey, M. K., Ahmad, M. M., Siddiqui, S., Kumar, A. and Singh, S. K. (2023). Alternaria Blight – a serious affliction of rapeseed mustard: a review. *Biological Forum – An International Journal*. 15(3): 436-443.

Ravella, N. K., Udathala, D. and Iska, S. R. (2023). Efficacy of eco-friendly treatments on yield attributes in Indian mustard (*Brassica juncea* L.) against alternaria blight. *Environment Conservation Journal*. 24 (1):171-175.

Reddy, T.Y. and Reddi, G.H.S. (2004). Principles of Agronomy IInd edition. Kalyani Publisher. 527.

Shrivastava, K. D. and Swarnkar, K. (2014). Antifungal activity of leaf extract of neem (*Azadirachta indica* Linn). *International Journal of Current Microbiology and Applied Sciences*. 3(5): 305-308

Thakur, Y. and Zacharia, S. (2017). Efficacy of bio-agents and plant extracts against alternarial leaf blight of mustard (*Brassica juncea* L.). *International Journal of Chemical Studies*. 6(1): 1187-1189.

Tuite, J. (1969). Plant pathological methods: fungi and bacteria. *Burgess Publishing*. Minnesota. 239.

Wheeler, B. E. J. (1969). An Introduction to plant disease. *John Wiley and Sons Limited* London. 301.

Sailaja, S., Lal, A. A. and Simon, S. (2017). Comparative efficacy of bio agents, botanicals and fungicides against alternaria leaf spot (*Alternaria brassicae*) of cauliflower (*Brassica oleracea* var. *botrytis*). *Journal of Pharmacognosy and Phytochemistry*.6(5): 519-521.

Sharma, L. R., Ahir, R. R., Yadav, L. S., Sharma, P. and Ghasolia, P. R. (2021). Effect of nutrients and plant extracts on Alternaria blight of tomato caused by *Alternaria alternata*. *Journal of Plant Diseases and Protection*. 10(8): 165-175.