

Comparative *In Vitro* Assessment of Fungicides for Managing *Fusarium oxysporum* f. sp. *capsici* in Chilli

ABSTRACT

Chilli wilt disease, caused by *Fusarium oxysporum* f. sp. *capsici*, poses a significant threat to chilli cultivation, impacting yield and quality. This study ^{was} aimed to evaluate the fungitoxic effects of various fungicides against *Fusarium oxysporum* f. sp. *capsici* under *in vitro* conditions. Eight different fungicides viz., Tebuconazole 6.7 % + Captan 26.9 % w/w SC, Thiophenate Methyl 70% WP, Carbendazim 12% + Mancozeb 63% WP, Tebuconazole 50% + Trifloxystrobin 25% WG, Carbendazim 50% WP, Pyraclostrobin 20% WG, Chlorothalonil 75% WP and Azoxystrobin 18.2% + Difenoconazole 11.4% SC were tested at concentrations of 500 ppm and 1000 ppm using the Poison food technique. The experiment ^{was} designed using a Completely Randomized Design (CRD) and conducted at the Department of Plant Pathology, Dr. Sharadchandra Pawar College of Agriculture, Baramati, during the academic year 2023-24. All fungicides significantly ($P < 0.01$) inhibited the mycelial growth of *Fusarium oxysporum* f.sp. *capsici*, with Carbendazim 12% + Mancozeb 63% WP showing the highest efficacy, followed by Tebuconazole 50% + Trifloxystrobin 25% WG and Tebuconazole 6.7 % + Captan 26.9 % w/w SC across both the concentrations. This study suggests that the inhibition of pathogen growth increased with higher fungicide concentrations.

Key words: *Fusarium oxysporum* f. sp. *capsici*, fungicides, growth inhibition

1. INTRODUCTION

Chilli (*Capsicum annum* L.) is a valuable vegetable and spice crop, recognized globally for its aroma, taste, flavour and hotness. Chilli peppers are widespread in cultivation across tropical and subtropical areas around the globe. It is susceptible to numerous pathogens that can impair its growth and reduce yields. *Fusarium* wilt is some of severe biotic threats to chilli cultivation, affecting its marketability and export potential by reducing both the quality and quantity of the produce. It affects various commercial varieties and species within the *Capsicum* genus across different regions.

⇒ India stands as the world's leading producer, consumer and exporter of chilli peppers. India exports chilli to numerous international markets, including the USA, Canada, UK, Saudi Arabia, Germany, Malaysia, Singapore and many others. Major chilli cultivation regions in India include Andhra Pradesh, Maharashtra, Karnataka, Uttar Pradesh, Punjab, Tamil Nadu, Rajasthan, Odisha, West Bengal and Madhya Pradesh. Andhra Pradesh leads in dry chilli production with 700,000 tonnes, contributing 37.35 per cent of the total national production. Telangana follows with 433,120 tonnes (23.11%), while Madhya Pradesh is third, producing 296,690 tonnes (15.83%) of dry chillies. Other top dry chilli producing regions consist of Karnataka, Orissa, Maharashtra, Gujarat, Tamil Nadu, Assam and Punjab. In the context of production of green chillies in India, Karnataka stands as the top producer, contributing 1,006.96 thousand tonnes production. Following this, Madhya Pradesh produces 726.90 thousand tonnes, while Andhra Pradesh ranks third with 506.87 thousand tonnes production. Other major states involved in green chilli production are Bihar, Maharashtra and Jharkhand. (National Horticulture Board, 2021-22).

Anonymous,

The emergence of aggressive and invasive plant pathogens poses a serious risk to sustainable crop production and food security, significantly affecting both crop yield and quality. The agricultural sector, which depends heavily on fruit and vegetable cultivation, is particularly vulnerable. Chilli peppers are prone to various pathogens that hinder their growth and reduce yields. Among these, Fusarium wilt represents a major biotic threat, impacting the marketability and export potential of chilli by reducing the quality and quantity of the produce. This disease affects multiple commercial varieties and species within the Capsicum genus across diverse regions. Initially identified in New Mexico (Leonian, 1919), Fusarium wilt is now prevalent in India, where it leads to around 25% yield losses (Najar *et al.*, 2006). In the United States, the disease is observed in about 35% of states. It has been reported that *Fusarium oxysporum* can cause the death of up to 56% of chilli seedlings within a 30 days period following soil infection (Roberts *et al.*, 2004) Ragab *et al.*, 2012). Globally, Fusarium wilt has been associated with yield reductions ranging from 10% to 80% (Loganathan *et al.*, 2013). Disease management for wilt in chilli depends on the implementation of cultural, biological and chemical methods. In many cases, chemical treatments offer the most practical means for managing plant diseases, being both cost effective and more efficient than other available options. Therefore, this research was conducted to manage wilt disease using various fungicides at the Plant Pathology Department, Dr. Sharadchandra Pawar College of Agriculture, Baramati, during the 2023-24 academic year.

2. MATERIAL AND METHODS

2.1 isolation and identification of *Fusarium oxysporum* f. sp. *capsici*

The diseased plant samples were collected from wilt infected chilli plot, Malegaon, Baramati. The diseased chilli plants exhibiting symptoms of wilt, including yellowing of leaves, leaf drying and browning of the vascular tissues. The pathogen was isolated from infected roots/stems of chilli plant. The isolation was performed using tissue isolation method on Potato dextrose agar. After 3 to 4 days, the growth of pathogen was observed in petri dishes. A pure culture was subsequently obtained using the hyphal tip culture method under aseptic conditions and maintained through PDA slants in test tubes.

The pathogen was identified by its distinct cultural and morphological characteristics. The culture achieved maximum growth after incubating for 7 days at $27 \pm 1^\circ\text{C}$. The culture showed circular fluffy growth with a whitish and pinkish colour of the colony. The pigmentation of colony was pinkish to brownish. Microscopic observations like mycelium, size and shape of conidia, chlamydospores were examined and correlated with Booth's (1971) standard descriptions of *Fusarium* spp.

2.2 Pathogenicity of *Fusarium oxysporum* f. sp. *capsici*

The pathogenicity of *Fusarium oxysporum* f. sp. *capsici* was confirmed using the sick soil method, following Koch's postulates. Pots were inoculated with the pathogen and chilli seedlings (Pusa Jwala) were transplanted. After 10-12 days, inoculated plants showed characteristic symptoms such as leaf yellowing, browning, stem base browning and wilting. These symptoms matched those of naturally diseased plants, while uninoculated plants remained healthy. The pathogen was reisolated from diseased plants, cultured, and its characteristics matched those from the naturally wilted plants. This confirmed *Fusarium oxysporum* f. sp. *capsici* as the cause of chilli wilt, fulfilling Koch's postulates.

2.3 Evaluation of fungicides against *Fusarium oxysporum* f. sp. *capsici*

Eight different fungicides viz., Tebuconazole 6.7% + Captan 26.9% w/w SC, Thiophenate Methyl 70% WP, Carbendazim 12% + Mancozeb 63% WP, Tebuconazole 50% + Trifloxystrobin 25% WG, Carbendazim 50% WP, Pyraclostrobin 20% WG, Chlorothalonil 75% WP and Azoxystrobin 18.2% + Difenoconazole 11.4% SC were tested at concentrations of 500 ppm and 1000 ppm using the poison food technique (Nene and Thapliyal, 1993). The experiment was designed in a Complete Randomized Design (CRD), which included eight

missing in reference list

under in vitro

followed by

and drying

whether pathogen was mass multiplied (or) die off by the pathogen

treatments along with a control treatment and three replications for each treatment. For this experiment, Potato dextrose agar was used as the basal culture medium.

2.3.1 Experimental procedure:

Stock solutions of each fungicide were prepared by dissolving the appropriate amount of active ingredient in a conical flask. The required quantities of each fungicide were aseptically incorporated into molten PDA to achieve concentrations of 500 and 1000 ppm. To prevent bacterial contamination, streptomycin was added to each flask prior to pouring the medium into sterilized petri plates. A medium without any fungicide served as the control. This mixture was then aseptically dispensed at a volume of 20 ml per plate into sterile glass petri dishes with a diameter of 90 mm and allowed to solidify at room temperature. Using sterilized cork borer, 5 mm diameter culture discs were cut from 7 days old actively growing culture of the test pathogen. These discs were then inoculated in the center of the PDA medium in the petri dishes, positioned in reverse to ensure close contact between the test pathogen and the fungicide treated PDA. Three PDA plates per treatment were prepared with fungicide and inoculated accordingly, while an equal number of PDA plates without fungicide were inoculated separately with the test pathogen maintained as untreated control. All plates were then incubated at a temperature of $27 \pm 1^\circ\text{C}$ for 7 days. The inhibition percentage of the mycelial growth of pathogen was calculated using the formula proposed by Vincent (1927). The data collected were subjected to statistical analysis following the procedures described by Panse and Sukhatme (1967).

$$\text{Percent Inhibition} = \frac{C - T}{C} \times 100$$

Where, C = Radial growth of fungus on control plate
T = Radial growth of fungus on treated plate

3. RESULT AND DISCUSSION

The study confirmed that all fungicides tested at concentrations of 500 ppm and 1000 ppm effectively restricted the mycelial growth of *Fusarium oxysporum* f. sp. *capsici* compared to the control group. Inhibition levels increased with higher fungicide concentrations.

At 500 ppm concentration, the growth inhibition rates of the *Fusarium oxysporum* f. sp. *capsici* vary from 27.97 per cent to 100 per cent. Among the eight fungicides tested, Carbendazim 12% + Mancozeb 63% WP showed 100 per cent mycelial growth inhibition of the test pathogen. This treatment was superior compared to other treatments. The next best treatment was Tebuconazole 50% + Trifloxystrobin 25% WG with growth inhibition of 90.36 per cent. Following this, Tebuconazole 6.7% + Captan 26.9% w/w SC showed 87.78 per cent growth inhibition. Carbendazim 50% WP showed moderate efficacy with a mycelial growth inhibition of 76.30 per cent, which was followed by Azoxystrobin 18.2% + Difenconazole 11.4% SC with 60.19 per cent growth inhibition. The less effective treatments were Chlorothalonil 75% WP (51.11%) and Pyraclostrobin 20% WG (46.86%). Thiophenate Methyl 70% WP was the least effective among all tested fungicides with only 27.97 per cent growth inhibition rate. (Table 1; Fig. 1; Plate 1).

At 1000 ppm concentration, the test pathogen showed inhibition percentages ranging from 51.86 per cent to 100 per cent. The fungicide treatment with Carbendazim 12% + Mancozeb 63% WP again showed 100 per cent mycelial growth inhibition of test pathogen and was the treatment with the highest efficacy compared to others. Again, the second most effective treatment was Tebuconazole 50% + Trifloxystrobin 25% WG with 93.88 per cent growth inhibition. This was followed Tebuconazole 6.7% + Captan 26.9% w/w SC with 89.81 per cent growth inhibition. Carbendazim 50% WP also showed moderate effectiveness at 1000 ppm concentration with an 86.30 per cent growth inhibition. Other fungicides also found effective at this concentration include Azoxystrobin 18.2% + Difenconazole 11.4% SC (74.81%), Pyraclostrobin 20% WG (64.45%) and Chlorothalonil 75% WP (61.67%). The least inhibition was found in Thiophenate Methyl 70% WP, with a growth inhibition of 51.86 per cent.

Petri plates

increase in concentration of fungicides

least growth inhibition was observed with

Among fungicides tested

The results of the current study showed that, at both the concentrations (500 ppm and 1000 ppm) the mycelial growth was completely inhibited by Carbendazim 12% + Mancozeb 63% WP. Sanap *et al.* (2020) conducted similar experiment of evaluation of different fungicides at different concentrations against *Fusarium oxysporum* under *in vitro* conditions. They reported that, at all three concentrations (@ 1500, 2000 and 2500 ppm), Carbendazim 25% + Mancozeb 50% showed complete inhibition of mycelial growth followed by Tebuconazole 50% + Trifloxystrobin 25%. Ghimire *et al.* (2022) also reported that Saaf (Carbendazim 25% + Mancozeb 50%) inhibits 100% mycelial growth of *Fusarium* spp. followed by Nativo (Tebuconazole 50% + Trifloxystrobin 25%) with inhibition rate of 87.04 per cent. Bhujbal *et al.* (2021) conducted an *in vitro* study to evaluate fungicides against *Fusarium oxysporum*, finding that the combination of Carbendazim 12% + Mancozeb 63% WP completely inhibited the mycelial growth of pathogen. Gabrekiristos *et al.* (2020) reported that URGI 75% WP, which contains Carbendazim and Mancozeb, effectively suppressed the mycelial development of *Fusarium oxysporum* f. sp. *capsici*, suggesting its potential for further *in vivo* testing. Additionally, Pimpale *et al.* (2023) assessed various fungicides against *Fusarium oxysporum* f. sp. *capsici* and found that Carbendazim 12% + Mancozeb 63% WP had the highest efficacy, achieving a maximum growth reduction of 93.72 per cent. These findings are consistent with Golakiya *et al.* (2018), Rao *et al.* (2020), Sahane *et al.* (2021), Chand and Singh (2021), Yadav *et al.* (2022), Kartha *et al.* (2023) and Buttar *et al.* (2023).

Kindly include mechanism of action of Carbendazim + mancozeb

Table 1. Efficacy of various fungicides against *F. oxysporum* f. sp. *capsici* under *in vitro* conditions

Tr. No.	Treatments	Colony Diameter* (mm)		% Growth Inhibition	
		500 ppm	1000 ppm	500 ppm	1000 ppm
T ₁	Tebuconazole 6.7 % + Captan 26.9 % w/w SC	11.00	9.17	87.78	89.81
T ₂	Thiophenate Methyl 70%WP	64.83	43.33	27.97	51.86
T ₃	Carbendazim 12% + Mancozeb 63% WP	0.00	0.00	100	100
T ₄	Tebuconazole 50% + Trifloxystrobin 25% WG	8.67	5.50	90.36	93.88
T ₅	Carbendazim 50%WP	21.33	12.33	76.30	86.30
T ₆	Pyraclostrobin 20%WG	47.83	32.00	46.86	64.45
T ₇	Chlorothalonil 75%WP	44.00	34.50	51.11	61.67
T ₈	Azoxystrobin 18.2%+ Difenconazole 11.4% SC	35.83	22.67	60.19	74.81
T ₉	Control	90.00	90.00		
	S.E.(m)±	0.25	0.38		
	C.D. (0.01)	0.74	1.12		

*Mean of three replications




Fig. 1 Per cent Growth inhibition of *Fusarium oxysporum* f. sp. *capsici* under *in vitro* conditions




Plate 1: Evaluation of various fungicides against *Fusarium oxysporum* f. sp. *capsici* under *in vitro* conditions

4. CONCLUSION

The study demonstrated that all tested fungicides significantly inhibited the mycelial growth of *Fusarium oxysporum* f. sp. *capsici*, with higher concentrations yielding greater inhibition. Among the fungicides, Carbendazim 12% + Mancozeb 63% WP was the most effective, achieving 100% inhibition at both 500 ppm and 1000 ppm. Tebuconazole 50% + Trifloxystrobin 25% WG and Tebuconazole 6.7% + Captan 26.9% w/w SC also exhibited high efficacy. Thiophanate Methyl 70% WP was the least effective in both concentrations.

DISCLAIMER

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts. **COMPETING INTERESTS:** Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous, 2021-22. Indian Horticulture Database. National Horticulture Board, Gurgaon, Haryana, India
2. Bhujbal, M. M., Deokar, C. D., Bachar, D. A., Suryawanshi, J. S. and Daingade, N. M. (2021) *In vitro* efficacy of fungicides against *Fusarium oxysporum* f. sp. *lycopersici* causing Fusarium Wilt of Tomato. Int. J. Curr. Microbiol. App. Sci., 10(1): 623-627
3. Booth, C. 1971. The genus *Fusarium*. Commonwealth Mycological Institute, Kew, UK.
4. Buttar, H. S., Singh, A., Sirari, A., Anupam, Kaur, K., Kumar, A., Lal, M. K., Tiwari, R. K. and Kumar, R. (2023). Investigating the impact of fungicides and mungbean genotypes on the management of pod rot diseases caused by *Fusarium equiseti* and *Fusarium chlamydosporum*. *Frontiers in plant science*, 14: 1164245.
5. Chand, K. and Singh, S.K. (2021). Evaluation of Different Model Chemicals against Panama Wilt of Banana Incited by *Fusarium oxysporum* f. sp. *cubense* TR4. Int. J. Curr. Microbiol. App. Sci., 10(2): 951-957
6. Gabrekristos, E., Ayana, G. and Kabeto, E. (2020). *In vitro* Evaluation of Some Fungicides against *Fusarium oxysporum* the Causal of Wilt Disease of Hot Pepper (*Capsicum annum* L.) in Ethiopia. Adv Crop Sci Tech, 8(2) w d
7. Ghimire, M., Adhikari, N., Puri, S., Baral, S. and Basnet, B. (2022). Efficacy of chemical fungicides against the *Fusarium* rhizome rot of Ginger (*Zingiber officinale*). *Archives of Agriculture and Environmental Science*, 7(3): 402-406.
8. Golakiya, B. B., Bhimani, M. D. and Akbari, L. F. (2018). Efficacy of different fungicides for the management of chickpea wilt (*Fusarium oxysporum* f. sp. *ciceri*). *International Journal of Chemical Studies*, 6(2): 199-205
9. Kartha, A. S., Thara, S. S. and Divya, S. (2023). Evaluation of *in vitro* efficacy of fungicides against *Fusarium proliferatum* inciting root rot of cassava. *The Pharma Innovation Journal*, 12(5): 3261-3263.
10. Leonian, L. H. (1919). *Fusarium* wilt of Chile pepper. *New Mexico Agricultural Experimental Station Technical Bulletin*, (p. 121)
11. Loganathan, M., Venkataravanappa, V., Saha, S., Sharma, B.K., Tirupathi, S. and Verma, M.K. (2013). Morphological, cultural and molecular characterizations of *Fusarium* wilt infecting tomato and chilli. In: National Symposium on Abiotic and Biotic Stress Management in Vegetable Crops, Indian Society of Vegetable Science. IIVR. Varanasi., 12-14
12. Najar, A., M. Ahmed, M. Bhat and Z. Bhat. (2006). Effect of various soil amendments on chilli wilt (*Fusarium pallidoroseum* (Cooke) Sacc). *Sher-e-Kashmir University of Agricultural Sciences and Technology Journal of Research.*, 8:236-239
13. Nene, Y. L. and Thapliyal, R. N. (1993). Fungicides in Plant disease control. 3rd edition. IBH Pub. Co. New Delhi, India: 225
14. Panse, V. G. and Sukhatme, P. V. 1967. Statistical Methods for Agricultural Workers ICAR, New Delhi
15. Pimpale, K. B., Suryawanshi, J. S., Deokar, C. D. and Bansode, K. A. (2023). *In vitro* evaluation of different fungicides against *Fusarium oxysporum* f. sp. *capsici*. *The Pharma Innovation Journal.*, 12(2): 2935-2938
16. Ragab, M., Ashour, A., Abdelkader, M., el-mohamedy, R. and Abdel-Aziz, A. (2012). *In vitro* Evaluation of Some Fungicides Alternatives Against *Fusarium oxysporum* the Causal of Wilt Disease of Pepper (*Capsicum annum* L.). *International Journal of Agriculture and Forestry.*, 2(2) 70-77
17. Rao, V. G., Dhutraj, D. N., Navgire, K. D., Ambadkar, C. V. and Apet, K. T. (2020). *In vitro* efficacy of fungicides against *Fusarium oxysporum* f. sp. *melongenae* causing eggplant wilt. Chem Sci Rev Lett 9(36) 1074-1081
18. Roberts, P., Adkins, S., Pemeznny, K. and Jones, J. (2007). Diseases of Pepper and their Management. In book: Diseases of Fruits and Vegetables., 2:333-387

Page number is missing

need expanded form

need exact form

need expanded form

not found in the text

need expanded form

- ✓ 19. Sahane, P. A., Chavan, R. A., Brahmankar, R. G., Kolhe, D. B. and Udar, V. B. (2021). *In vitro* efficacy of fungicides against *Fusarium oxysporum* f. sp. *ciceri*. *Journal of Pharmacognosy and Phytochemistry*, **10**(1): 978-984
- ✓ 20. Sanap, S. B., Mete, V. S., Jaiswal, K. L., Sanap, S. B. and Mulekar, V. G. (2020). *In vitro* efficacy of non systemic and combi fungicides against wilt of tomato caused by *Fusarium oxysporum* f. sp. *lycopersici*. *Journal of Pharmacognosy and Phytochemistry*, **9**(5): 3280-3283
- ✓ 21. Vincent J. M. (1927). Distortion of fungal hyphae in presence of certain inhibitors. *Nature*. **59**: 850
- ✓ 22. Yadav, P., Ahir, R., Yadav, A., Yadav, V., Babu, S., Yadav, P. D., Yadav, P., Meena, R., Choudhary, S. and Kumar, V. (2022). Comparative efficacy of fungicides, biocontrol agents and plant extracts against wilt of cluster bean caused by *Fusarium solani*. *Agricultural mechanization in Asia*, **53**(2):5662-5672

UNDER PEER REVIEW