

# Original Research Article

## "Evaluation of Bioagents Against *Stemphylium vesicarium* Inducing Stemphylium Blight in Onions (*Allium cepa* L.)"

---

### ABSTRACT

Stemphylium blight of onion incited by *Stemphylium vesicarium* is one of the economically important diseases, causing severe yield losses in onion crop. The present investigation was carried out to evaluate efficacy of different bioagents against *Stemphylium vesicarium* under *in vitro* conditions. The experiment was laid out in a Completely Randomized Design (CRD) with three replications per treatment. Five fungal biocontrol agents viz., *Trichoderma harzianum*, *T. viride*, *T. hamatum*, *T. virens* and a fungal consortium from MPKV, Rahuri along with two bacterial biocontrol agents namely *Pseudomonas fluorescens* and *Bacillus subtilis* were evaluated for their effectiveness against *S. vesicarium*. The results show significant variation in the inhibitory effects of bioagents on the mycelial growth of the test pathogen with percent growth inhibition ranging between 53.70 per cent to 85.19 per cent. *Trichoderma harzianum* was reported to be most effective, showing a minimum colony diameter of 13.33 mm with the highest growth inhibition (85.19%) thus reported as superior to all other treatments whereas, *Pseudomonas fluorescens* was reported with a least percent growth inhibition (53.70%).

**Key words:** *Stemphylium vesicarium*, *Trichoderma*, Bioagents, Growth inhibition

### 1. INTRODUCTION

The onion (*Allium cepa* L.) is the important commercially cultivated vegetable crop all over the world belonging to family *Alliaceae*. In India, onions are extensively cultivated for both domestic consumption and export purpose. India is one of the largest producers of onions, with Maharashtra is the leading state in production, contributing about 43% of the nation's total onion output, producing 13,301.7 tons. Madhya Pradesh, Karnataka, and Gujarat follow as the next leading producers. In the 2022-23 season, India's onion production reached an estimated 318 lakh metric tons (LMT), slightly surpassing the previous year's 316.98 LMT. (Ministry of Farmer and Agriculture Welfare, 2023).

Onions are susceptible to a range of diseases incited by specific pathogens causing severe impact on health and yield of the crop. Stemphylium leaf blight caused by *Stemphylium vesicarium* is one of the economically important diseases causing severe losses in onion production. For the first time, Stemphylium blight of onion was documented in India by Rao and Pavgi (1973). They reported the yield of onion bulbs was reduced by nearly 90% due to disease incidence.

Spread of Stemphylium blight is occur by infected seed material and through infected soil. Initial symptoms of Stemphylium blight appear on the foliar plant parts. The symptoms of Stemphylium blight include small, water-soaked lesions that develop into dark brown to black spots on the leaves, often leading to complete blighting. (Tesfaendrias *et al.*, 2014). These lesions then develop into elongated spots, eventually turning dark olive brown

to black due to spore development. The center of the lesions turns brown and thus produces conidiophore and conidia which gives blighted appearance to infected plant. Severe infestation may result in complete blighting of the leaves.

Eco friendly, non chemical management techniques are becoming more prominent as an alternative to chemical management practices. It is feasible to control the *Stemphylium* blight of onions with the application of fungicides, but prolonged use of chemicals has a harmful environmental impact and has caused pathogens to become resistant to them. So, keeping in view the commercial importance of the crop in India and the magnitude of losses caused by *Stemphylium* blight in onion, the present study was planned to evaluate efficacy of different bioagents against *Stemphylium vesicarium* under *in vitro* conditions.

## **2.MATERIAL AND METHODS**

### **2.1 Isolation and identification of the pathogen associated with onion leaf blight**

Naturally infested onion leaves showing characteristic symptoms of *Stemphylium* blight were collected from the experimental plot located at Krishi Vigyan Kendra, Baramati. Collected samples were put in paper bag and brought to plant pathology laboratory. The infected onion leaves were washed thoroughly with distilled water. Leaf samples were blot dried and cut with sharp sterilized blade into small bits (5 mm), keeping half healthy and half diseased portion intact. These pieces were surface sterilized with 0.1% aqueous solution of sodium hypochlorite for two minutes followed by giving three consecutive washes with sterile distilled water to remove the residues of sodium hypochlorite. The surface sterilized diseased leaf bits were inoculated on the solidified and cooled PDA medium in petri plates under aseptic conditions of laminar air flow cabinet. Inoculated plates were incubated in BOD incubator at  $24\pm 1^{\circ}\text{C}$  temperature for 7 days and sub culture was carried out until the pure isolate of *Stemphylium vesicarium* was obtained and identified. The isolated test pathogen was identified based on colony characters, spore morphology referring to the description by Chowdhury *et al.* (2015).

### **2.2 Pathogenicity test**

Onion variety Agrifound Dark Red was used to perform the pathogenicity test. Potting mixture of soil: sand: FYM were sterilized and filled in earthen pots in 2:1:1 ratio. Seeds were sown in pots and healthy onion seedling per pot was maintained, watered regularly and kept in the polyhouse for further growth. The test pathogen (*Stemphylium vesicarium*) was mass multiplied on the culture medium of PDA in petri dishes. Spore cum mycelial suspension of the test pathogen was prepared from 7 to 8 days old culture in plates by flooding with 5-10 ml sterile distilled water. The resultant spore mycelial suspension was filtered through double layered muslin cloth and filtrate obtained was suitably diluted with sterile distilled water. Spore concentration was adjusted to  $5\times 10^4$  conidia per ml with hemocytometer. 2 months old seedlings of onion which were already grown in earthen pots were artificially inoculated by spraying spore suspension of the test pathogen with hand atomizer. Onion seedling of uninoculated control was maintained in earthen pots and sprayed with sterile water i.e., without inoculum. Both inoculated and uninoculated pots in the polyhouse were maintained with 80 - 90% humidity and  $24\pm 1^{\circ}\text{C}$  optimum temperature by covering them with clear polyethylene bags for 72 hours. After 3 days, polythene bag was removed and disease development was assessed until the development of characteristic symptoms of *Stemphylium* blight. Pathogen was reisolated from the developed symptoms on artificially inoculated leaves, and the resulting cultures were compared with the original inoculants to fulfil Koch's postulates.

### **2.3 Antimicrobial efficacy of bioagents**

The antimicrobial efficacy of various biocontrol agents, including *Trichoderma harzianum*, *Trichoderma hamatum*, *Trichoderma viride*, *Trichoderma virens*, *Pseudomonas fluorescens*,

*Bacillus subtilis*, and MPKV fungal consortium, was evaluated against *Stemphylium vesicarium* using the dual culture technique (Dennis and Webster, 1971) on PDA medium.

### 2.3.1 Dual Culture Technique

PDA medium was prepared and sterilized in an autoclave at 15 psi pressure at 121°C for 20 minutes. Sterilized media (20 ml) were poured into sterilized Petri dishes (90 mm diameter). After solidification, each Petri dish was inoculated with a 5 mm mycelial disc from an actively growing 7-day-old culture of *S. vesicarium* placed at the periphery of one side of the plate. The biocontrol agent disc was inoculated on the opposite side of the same plate, equidistant from the pathogen. For bacterial bioagents, the inoculum was streaked on opposite sides of the pathogen. PDA plates inoculated solely with *S. vesicarium* served as untreated controls.

### 2.3.2 Experimental Procedure

All inoculated petri plates were incubated at 24 ± 1°C for seven days. Each treatment was replicated three times. After the incubation period, observations were carried out when the fungal growth in the control plates reached a maximum diameter of 90 mm. Radial growth of the test pathogen was measured, and the percentage of growth inhibition relative to the control was calculated using Vincent's (1947) formula,

$$\text{Percent Inhibition} = \frac{C - T}{C} \times 100$$

Where, C = Radial growth of fungus on control plate

T = Radial growth of fungus on treated plate

## 3. RESULT AND DISCUSSION

The results show significant variation in the inhibitory effects of bioagents on the mycelial growth of the test pathogen with per cent growth inhibition ranging between 53.70 and 85.19 per cent. *Trichoderma harzianum* was reported to be most effective, showing a minimum colony diameter of 13.33 mm with the highest growth inhibition of 85.19 per cent and being reported as superior to all other treatments. The mechanism behind *T. harzianum*'s effectiveness can be attributed to the secretion of cell-wall degrading enzymes such as chitinases and glucanases, alongside its ability to outcompete pathogens for space and nutrients. (Contreras-Cornejo *et al.*, 2020).

The second most effective treatment was *Trichoderma viride*, with a colony diameter of 16.83 mm and growth inhibition of 81.30 per cent. *Trichoderma hamatum* and *Trichoderma virens* were also notably effective, with colony diameters of 22.50 mm and 29.17 mm with inhibition rates of 75.00 per cent and 67.59 per cent respectively. The *Trichoderma* species causes enzyme production and secondary metabolite synthesis thus have competitive abilities against other pathogens in the rhizosphere. (Vinale *et al.*, 2021).

The MPKV fungal consortium showed significant effectiveness, with a colony diameter of 27.50 mm and growth inhibition of 69.44 per cent. The use of fungal consortia has been shown to produce synergistic effects, enhancing overall biocontrol potential against diverse pathogens (Vaid *et al.*, 2022). This suggests that combining multiple strains may offer a broader spectrum of protection and resilience in integrated pest management (IPM) system.

On the other hand, the bacterial bioagents, *Bacillus subtilis* and *Pseudomonas fluorescens*, showed comparatively moderate inhibition rates. *Bacillus subtilis* was found to be comparatively less effective with colony diameter of 36.33 mm and 59.63 per cent growth inhibition. whereas, *Pseudomonas fluorescens* was reported with a least per cent growth inhibition (53.70%) and a colony diameter of 41.67 mm. Although *Bacillus* and *Pseudomonas* species are widely recognized for their antimicrobial activities producing secondary metabolites such as lipopeptides and siderophores. The moderate performance of these bacterial agents suggests they may be better suited for suppressing bacterial pathogens or used in combination with fungal bioagents for enhanced its control.

**Table 1. Antimicrobial efficacy of different bioagents against the *Stemphylium vesicarium* under *in vitro* condition**


Tr. No.	Treatments	Colony diameter of Pathogen* (mm)	Per cent growth inhibition *
T <sub>1</sub>	<i>Trichodermaharzianum</i>	13.33	85.19
T <sub>2</sub>	<i>Trichoderma viride</i>	16.83	81.30
T <sub>3</sub>	<i>Trichoderma hamatum</i>	22.50	75.00
T <sub>4</sub>	<i>Trichoderma virens</i>	29.17	67.59
T <sub>5</sub>	<i>Pseudomonas fluorescens</i>	41.67	53.70
T <sub>6</sub>	<i>Bacillus subtilis</i>	36.33	59.63
T <sub>7</sub>	MPKV fungal consortium	27.50	69.44
T <sub>8</sub>	Control	90.00	-
	<b>SE(m)±</b>	<b>0.46</b>	
	<b>C.D. at 1%</b>	<b>1.92</b>	

\*Mean of three replications



**Fig.1** *In vitro* efficacy of various bioagents against *Stemphylium vesicarium*

**Plate 1.** *In vitro* efficacy of various bioagents against *Stemphylium vesicarium*



The effectiveness of *Trichoderma harzianum* and *Trichoderma viride* had been reported earlier by Kumar *et al.* (2012), Mishra and Gupta (2012), Shahnaz *et al.* (2013), Nainwal and Vishunavat (2015), Abo Elyousr (2016), Kamal *et al.* (2017), Gondaliya *et al.* (2020), Devi *et al.* (2022) that is in accordance with present findings.

The findings are exactly matching with the report of Kumar *et al.* (2012) who conducted an *in vitro* experiment to evaluate the efficacy of fungal bioagents *viz.*, *Trichoderma harzianum* and *T. viride* against the *Stemphylium* pathogen. Among the bioagents tested, *T. harzianum* showed the highest inhibition of colony growth (81.2%) followed by *T. viride* (74.5%)

The results obtained of present investigation are in close conformity with Mishra and Gupta (2012) who tested the efficacy of fungal biocontrol agents *viz.*, *T. viride*, *T. harzianum*, *T. hamatum*, *A. niger*, *T. koningii*, *T. virens* and bacterial antagonists *Pseudomonas fluorescens* and *Bacillus subtilis*. All antagonistic organisms demonstrated inhibitory effects on the growth of *S. vesicarium* with inhibition ranging from 19.20 per cent to 55.95 per cent. *T. viride* exhibited the highest level of inhibition about 56.15 per cent against *S. vesicarium*. This was followed by *T. harzianum* with inhibition of about 51.95 per cent and *T. koningii* with inhibition of 45.25 per cent, respectively. Conversely, *P. fluorescens* displayed the least effectiveness with inhibition rate 20.17 per cent.

Gondaliya *et al.* (2020) reported the *in vitro* efficacy of various isolates of bioagents against the *Stemphylium vesicarium* and *Alternaria porii* and resulted that *Trichoderma viride* shows the highest growth inhibition followed by *T. harzianum*, *longibrachiatum*, *P. fluorescens* and *Bacillus subtilis*. Furthermore, Hussein *et al.* (2007) also evaluated the *in vitro* efficacy of different bioagents against the *S. vesicarium* and reported that *Trichoderma harzianum* shows 78.1 per cent mycelial growth inhibition when dual cultured with *Stemphylium vesicarium* under *in vitro* condition.

#### **DECLARATION OF ARTIFICIAL INTELLIGENCE**

The authors hereby declare that no generative AI technologies, including but not limited to Large Language Models (such as ChatGPT, Copilot, etc.) or text-to-image generators, were utilized in the writing or editing of this manuscript.

#### **References**

1. Abo-Elyousr, K. A., Abdel-Hafez, O. and Abdel-Rahim, I. (2016). Control of *Stemphylium* leaf blight disease of onion and elevation of seed production using certain bioagents. *International Journal of Plant Pathology.*, **8**(1):1-7.
2. Chowdhury, H.A., Islam, N., Hossain, B., Ahmed, M., Mohsin, S. and Islam, R. (2015). A comparative analysis of culture media for optimizing the mycelial growth and sporulation of *Stemphylium vesicarium* cause of white blotch of onion. *Journal of Agricultural Science and Technology.*, **5**: 440-448.

3. Contreras-Cornejo, H. A., Macías-Rodríguez, L., del-Val, E. and Larsen, J. (2020). *Trichoderma* species: Versatile plant symbionts. *Phytochemistry Reviews*, **19**(3), 577–593.
4. Dennis, C. and Webster, J., (1971). Antagonistic properties of species-groups of *Trichoderma*: II. Production of volatile antibiotics. *Transactions of the British Mycological Society*, **57**(1): 41-44.
5. Devi, M., Banyal, D. K., Anudeep, B., and Sinha, D. (2022). Management of gray leaf spot of tomato caused by *Stemphylium lycopersici* under protected cultivation. *Plant Disease Research*, **36**(2):154–160
6. Gondaliya, K.K., Patel, S. I. and Thumbadiya, N.K. (2020) Efficacy of different Phyto extracts and biocontrol agents against foliar blight complex of onion *in vitro*. *Journal of Pharmacognosy and Phytochemistry*, **9**(4): 800-803
7. Hussein, M. A. M., Hasan, M. H. A., Allam, A. D. A. and Abo-Elyousr, K. A. M. (2007). Management of *Stemphylium* blight of onion by using biological agents and resistance inducers. *Egypt J. Phytopathology*, **35**(1): 49-60.
8. Kamal A. M., Abo Elyousr, Sobhy I. I., Abdel-Hafez and Ismail, R. Abdel-Rahim, (2017). Control of *Stemphylium* leaf blight disease of onion and elevation of seed production using certain bioagents. *International journal of plant pathology*, **8**:(1)384-391.
9. Kumar U., Naresh P. and Biswas S.K. (2012). ecofriendly management of stemphylium blight (*Stemphylium botryosum*) of garlic by plant extract and bioagents. *Hortflora Research Spectrum*, **1**(1): 42-45.
10. Ministry of Farmer and Agriculture Welfare. (2023). *Annual report on onion production and statistics*.
11. Mishra, K. R. and Gupta, P. R. (2012). *In vitro* evaluation of plant extracts, bioagents and fungicides against purple blotch and *Stemphylium* blight of onion. *Journal of Medicinal Plants Research*, **6**(30): 5658-5661.
12. Nainwal, D. and Karuna, V. (2016) Management of purple blotch and *Stemphylium* blight of onion in Tarai and Bhabar regions of Uttarakhand, India. *J. Applied and Natural Sci.*, **8**(1): 150-153.
13. Rao, N.N.R. and Pavgi, M.S. (1973). *Stemphylium* blight of onion. *Mycopathology*, **56**(2): 113- 118.
14. Shahnaz, E., Razdan, V. K., Rizvi, S. E., Rather T. R., Gupta, S. and Andrabi, M. (2013) Integrated Disease Management of Foliar Blight Disease of Onion: A Case Study of Application of Confounded Factorials. *Journal of Agricultural Science*, **5**(1) 1916- 9760
15. Tesfaendrias, M. T., Paibomesai, M., Celetti, M., and McDonald, M. R. (2014). The battle against *Stemphylium* leaf blight of onion in Ontario, Canada.
16. Vaid, S. K., Dubey, S. C. and Kumar, A. (2022). Biocontrol potential of *Trichoderma* spp. against fungal diseases: A review. *Biological Control*, **165**, 104776.
17. Vinale, F., Sivasithamparam, K., Ghisalberti, E. L., Marra, R., Barbetti, M. J., Li, H. and Lorito, M. (2021). *Trichoderma harzianum* as a biocontrol agent against *Stemphylium* blight: Mechanisms and application strategies. *Journal of Plant Pathology*, **103**(2), 345-356.
18. Vinale, F., Sivasithamparam, K., Ghisalberti, E. L., Marra, R., Barbetti, M. J., Li, H. and Lorito, M. (2021). *Trichoderma harzianum* as a biocontrol agent against *Stemphylium* blight: Mechanisms and application strategies. *Journal of Plant Pathology*, **103**(2), 345-356.
19. Vincent, J. M. (1927). Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*: 159-180.