

Original Research Article

Hilbert Graceful Labeling on Complete Multipartite Graph

Abstract: Let G be a simple, finite, connected, undirected, non-trivial graph with p vertices and q edges. $V(G)$ be the vertex set and $E(G)$ be the edge set of G . The n^{th} Hilbert number is denoted by H_n and is defined by $H_n = 4(n - 1) + 1$ where $n \geq 1$. A Hilbert graceful labeling is an injective function \mathcal{H} from the vertex set $V(G)$ to a set of Hilbert number $\{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\}$ which induces a bijective function \mathcal{H}^* from the set $E(G)$ to the set of number $\{1, 2, 3, 4, \dots, q\}$, where for each edge $uv \in E(G)$ with $u, v \in V(G)$ applies $\mathcal{H}^*(uv) = \frac{1}{4}|\mathcal{H}(u) - \mathcal{H}(v)|$. A graph with Hilbert graceful labeling is called a Hilbert graceful graph. This research aims to prove some complete multipartite graphs are Hilbert graceful.

AMS Subject Classification – 05C78

Keywords: Hilbert numbers, Graceful labeling, Hilbert graceful labeling, Hilbert graceful graph.

1. Introduction

A graph G consists of a finite set of vertices $V(G)$ and a set of edges c consisting of distinct, unordered pairs of vertices [1]. The graph discussed in this paper is a simple, undirected, and finite graph. $|V(G)|$ represents the number of vertices on graph G , and the number of edges on graph G is represented by $|E(G)|$. Graph labeling has been studied since the 60s. Graph labeling is a branch of graph theory that continues to develop. Labeling on a graph is the assignment of an integer value to the elements of the graph, usually a positive integer. Alex Rosa first discovered graceful labeling in 1967 [2]. Since this discovery, many researchers have been interested in looking for graceful labeling constructs and their variations. Several graphs with graceful labeling include tree graphs with vertices less or equal to 35, circle graph C_n for $n = 0 \pmod{4}$ or $n = 3 \pmod{4}$, and wheel graph W_n . Another class of graphs known to have graceful labeling can be seen in the survey conducted by Gallian [3].

The following shows some relevant research: graceful labeling of paths 4], graceful labeling of pendant graphs 5, 6, 7], vertex graceful labeling of caterpillar graphs 8], graceful labeling on torch graph 9], counting graceful labelings of trees 10], and other results on super graceful labeling of graphs 11]. Motivated by the above articles, the new type of graceful labeling called Hilbert gracefullabeling is introduced in this paper and Hilbert gracefullabeling of some complete bipartite graph is studied.

2. Definition

Definition 2.1: The graph $K_{1,m,n}$ is obtained by connecting all the vertices of complete bipartite graph $K_{m,n}$ to a vertex v .

Definition 2.2: The graph $K_{2,m,n}$ is obtained by connecting all the vertices of complete bipartite graph $K_{m,n}$ to a vertices u and v .

Definition 2.3: The graph $K_{1,1,m,n}$ is obtained by connecting all the vertices of complete bipartite graph $K_{m,n}$ to the end points of an edge uv .

Definition 2.4: A graph G with the vertex set $V(G) = \{u_i: 1 \leq i \leq m\} \cup \{v_j: 1 \leq j \leq n\} \cup \{w_k: 1 \leq k \leq l\}$ and the edge set $E(G) = \{u_i v_j: 1 \leq i \leq m, 1 \leq j \leq n\} \cup \{v_j w_k: 1 \leq j \leq n, 1 \leq k \leq l\}$ such that $|V(G)| = m + n + l$ and $|E(G)| = m(n + l)$ is denoted by $K_{m,n,l}$.

3. Main Result

Theorem 3.1: The graph $K_{1,m,n}$ admits Hilbert graceful labeling.

Proof: Let G be a $K_{1,m,n}$ graph.

$$V(G) = \{u_i: 1 \leq i \leq m\} \cup \{v_j: 1 \leq j \leq n\} \cup \{v\} \text{ and}$$

$$E(G) = \{u_i v_j: 1 \leq i \leq m; 1 \leq j \leq n\} \cup \{v u_i: 1 \leq i \leq m\} \cup \{v v_j: 1 \leq j \leq n\}.$$

$$|V(G)| = m + n + 1 \text{ and } |E(G)| = m n + m + n$$

We define a function $\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\}$

The vertex labeling is as follows:

$$\mathcal{H}(u_i) = 4[i] + 1 \quad 1 \leq i \leq m$$

$$\mathcal{H}(v_j) = 4[m + (m + 1) j] + 1 \quad 1 \leq j \leq n$$

$$\mathcal{H}(v) = 1 \qquad 1 \leq k \leq l$$

By above labeling pattern, we observed that function

$$\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\} \text{ is } 1 - 1.$$


From the induced function $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$, we get the edge labels as follows.

Table 1: Edge labels of the graph $K_{1,m,n}$

\mathcal{H}^*	Edge Labels	Value of i, j and k
$ \mathcal{H}(u_i) - \mathcal{H}(v_j) $	$m - i + (m + 1)j$	$1 \leq i \leq m, 1 \leq j \leq n$
$ \mathcal{H}(u_i) - \mathcal{H}(v) $	i	$1 \leq i \leq m$
$ \mathcal{H}(v_j) - \mathcal{H}(v) $	$m + (m + 1)j$	$1 \leq j \leq n$

From the above table 1, we observe that $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$ defined by $\mathcal{H}^*(uv) = \frac{1}{4}|\mathcal{H}(u) - \mathcal{H}(v)|$ is a bijective. Hence, \mathcal{H} is Hilbert graceful labeling and the graph $K_{1,m,n}$ is Hilbert graceful graph.

Example 3.1: Hilbert graceful labeling of the graph $K_{1,3,4}$



Theorem 3.2: The graph $K_{1,3,4}$ admits Hilbert graceful labeling.

Proof: Let G be a $K_{2,m,n}$ graph.

$$V(G) = \{u_i: 1 \leq i \leq m\} \cup \{v_j: 1 \leq j \leq n\} \cup \{u, v\} \text{ and}$$

$$E(G) = \{u_i v_j: 1 \leq i \leq m; 1 \leq j \leq n\} \cup \{u u_i: 1 \leq i \leq m\} \cup \{u v_j: 1 \leq j \leq n\} \cup \{v u_i: 1 \leq i \leq m\} \cup \{v v_j: 1 \leq j \leq n\}.$$

$$|V(G)| = m + n + 2 \text{ and } |E(G)| = mn + 2m + 2n$$

We define a function $\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\}$

The vertex labeling is as follows:

$$\mathcal{H}(u_i) = 4i + 1 \quad 1 \leq i \leq m$$

$$\mathcal{H}(v_1) = 1$$

$$\mathcal{H}(v_j) = 4[(m + 2)j - 2] + 1 \quad 2 \leq j \leq n$$

$$\mathcal{H}(u) = 4[mn + 2m + 2n] + 1$$

$$\mathcal{H}(v) = 4[mn + m + 2n - 1] + 1$$

By above labeling pattern, we observed that function

$$\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\} \text{ is } 1 - 1.$$


From the induced function $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$, we get the edge labels as follows.

Table 2: Edge labels of the graph $K_{2,m,n}$

\mathcal{H}^*	Edge Labels	Value of i, j and k
$ \mathcal{H}(u_i) - \mathcal{H}(v_j) $	i	$1 \leq i \leq m, j = 1$
$ \mathcal{H}(u_i) - \mathcal{H}(v_j) $	$(m + 2)j - i - 2$	$2 \leq j \leq n; 1 \leq i \leq m$
$ \mathcal{H}(u_i) - \mathcal{H}(u) $	$mn + 2m + 2n - i$	$1 \leq i \leq m$
$ \mathcal{H}(v_j) - \mathcal{H}(u) $	m	$j = 1$
$ \mathcal{H}(v_j) - \mathcal{H}(u) $	$mn + 2m + 2n + 2 - (m + 2)j$	$2 \leq j \leq n$
$ \mathcal{H}(u_i) - \mathcal{H}(v) $	$mn + m + 2n - 1 - i$	$1 \leq i \leq m$
$ \mathcal{H}(v_j) - \mathcal{H}(v) $	$mn + m + 2n - 1$	$j = 1$
$ \mathcal{H}(v_j) - \mathcal{H}(v) $	$mn + m + 2n + 1 - (m + 2)j$	$2 \leq j \leq n$

From the above table 2, we observe that $\mathcal{H}^*: E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$ defined by $\mathcal{H}^*(uv) = \frac{1}{4}|\mathcal{H}(u) - \mathcal{H}(v)|$ is a bijective. Hence, \mathcal{H} is Hilbert graceful labeling and the graph $K_{2,m,n}$ is Hilbert graceful graph.

Example 3.2: Hilbert graceful labeling of the graph $K_{2,m,n}$



Theorem 3.3: The graph $K_{1,1,m,n}$ admits Hilbert graceful labeling.

Proof: Let G be a $K_{1,1,m,n}$ graph.

$$V(G) = \{u_i: 1 \leq i \leq m\} \cup \{v_j: 1 \leq j \leq n\} \cup \{u, v\} \text{ and}$$

$$E(G) = \{u_i v_j: 1 \leq i \leq m; 1 \leq j \leq n\} \cup \{u u_i: 1 \leq i \leq m\} \cup \{u v_j: 1 \leq j \leq n\} \cup \{v u_i: 1 \leq i \leq m\} \cup \{v v_j: 1 \leq j \leq n\} \cup \{u v\}.$$

$$|V(G)| = m + n + 2$$

$$|E(G)| = m n + 2m + 2n + 1$$

We define a function $\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\}$

The vertex labeling is as follows:

$$\mathcal{H}(u_i) = 4i d + 1 \quad 1 \leq i \leq m$$

$$\mathcal{H}(v_j) = 4[m + (m + 2)j] + 1 \quad 1 \leq j \leq n$$

$$\mathcal{H}(u) = 4[mn + 2m + 2n + 1] + 1$$

$$\mathcal{H}(v) = 1$$

By above labeling pattern, we observed that function

$$\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\} \text{ is } 1 - 1.$$


From the induced function $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$, we get the edge labels as follows.

Table 3: Edge labels of the graph $K_{1,1,m,n}$.

\mathcal{H}^*	Edge Labels	Value of i, j and k
$ \mathcal{H}(u_i) - \mathcal{H}(v_j) $	$m - i + (m + 2)j$	$1 \leq i \leq m, 1 \leq j \leq n$
$ \mathcal{H}(u_i) - \mathcal{H}(u) $	$mn + 2m + 2n + 1 - i$	$1 \leq i \leq m$
$ \mathcal{H}(v_j) - \mathcal{H}(u) $	$mn + m + 2n + 1 - (m + 2)j$	$1 \leq j \leq n$
$ \mathcal{H}(u_i) - \mathcal{H}(v) $	i	$1 \leq i \leq m$
$ \mathcal{H}(v_j) - \mathcal{H}(v) $	$m + (m + 2)j$	$1 \leq j \leq n$
$ \mathcal{H}(u) - \mathcal{H}(v) $	$mn + 2m + 2n + 1$	

From the above table 3, we observe that $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$ defined by $\mathcal{H}^*(uv) = \frac{1}{4}|\mathcal{H}(u) - \mathcal{H}(v)|$ is a bijective. Hence, \mathcal{H} is Hilbert graceful labeling and the graph $K_{1,1,m,n}$ is Hilbert graceful graph.

Example 3.3: Hilbert graceful labeling of the graph $K_{1,1,3,4}$.



Theorem 3.4: The graph $K_{m,n,l}$ admits Hilbert graceful labeling.

Proof: Let G be a $K_{m,n,l}$ graph.

$$V(G) = \{u_i : 1 \leq i \leq m\} \cup \{v_j : 1 \leq j \leq n\} \cup \{w_k : 1 \leq k \leq l\} \text{ and}$$

$$E(G) = \{u_i v_j : 1 \leq i \leq m, 1 \leq j \leq n\} \cup \{u_i w_k : 1 \leq i \leq m, 1 \leq k \leq l\}.$$

$$|V(G)| = m + n + l$$

$$|E(G)| = m(n + l)$$

We define a function $\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\}$

The vertex labeling is as follows:

$$\mathcal{H}(u_i) = 4[i - 1] + 1 \quad 1 \leq i \leq m$$

$$\mathcal{H}(v_j) = 4[mj] + 1 \quad 1 \leq j \leq n$$

$$\mathcal{H}(w_k) = 4[mn + mk] + 1 \quad 1 \leq k \leq l$$

By above labeling pattern, we observed that function

$$\mathcal{H} : V(G) \rightarrow \{x : x = 4(i - 1) + 1, 1 \leq i \leq 2q\} \text{ is } 1 - 1.$$


From the induced function $\mathcal{H}^* : E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$, we get the edge labels as follows.

Table 4: Edge labels of the graph $K_{m,n,l}$.

\mathcal{H}^*	Edge Labels	Value of i, j and k
$ \mathcal{H}(u_i) - \mathcal{H}(v_j) $	$1 - i + mj$	$1 \leq i \leq m, 1 \leq j \leq n$
$ \mathcal{H}(u_i) - \mathcal{H}(w_k) $	$mn + 1 - i + mk$	$1 \leq i \leq m, 1 \leq k \leq l$

From the above table 4, we observe that $\mathcal{H}^*: E(G) \rightarrow \{1, 2, 3, 4, \dots, q\}$ defined by $\mathcal{H}^*(uv) = \frac{1}{4}|\mathcal{H}(u) - \mathcal{H}(v)|$ is a bijective. Hence, \mathcal{H} is Hilbert graceful labeling and the graph $K_{m,n,l}$ is Hilbert graceful graph.

Example 3.4: Hilbert graceful labeling of the graph $K_{3,4,4}$.



4. Conclusion

In this paper, we proved some complete bipartite graph with pendant edges are Hilbert graceful graph. Labeling pattern is demonstrated by means of illustrations, which provide better understanding of derived results. Analysing Hilbert graceful on other families of graph are our future work.

References

- 1] Bapat, R. B. (2010). *Graphs and matrices* (Vol. 27). New York: Springer.
- 2] Rosa, A. (1966, July). On certain valuations of the vertices of a graph. In *Theory of Graphs (Internat. Symposium, Rome)* (pp. 349-355).
- 3] Gallian, J. A. (2022). A dynamic survey of graph labeling. *Electronic Journal of combinatorics*, 6(25), 4-623.
- 4] Cattell, R. (2007). Graceful labellings of paths. *Discrete Mathematics*, 307(24), 3161-3176.
- 5] Graf, A. (2014). Graceful labelings of pendant graphs. *Rose-Hulman Undergraduate Mathematics Journal*, 15(1), 10.
- 6] Barrientos, C. (2005). Graceful graphs with pendant edges. *Australasian Journal of Combinatorics*, 33.
- 7] Akerina, A., & Sugeng, K. A. (2021, February). Graceful labeling on a multiple-fan graph with pendants. In *AIP Conference Proceedings* (Vol. 2326, No. 1). AIP Publishing.
- 8] Santhakumaran, A. P., & Balaganesan, P. (2018). Vertex graceful labeling of some classes of graphs. *Proyecciones (Antofagasta)*, 37(1), 19-43.
- 9] Manulang, J. M., & Sugeng, K. A. (2018). Graceful labeling on torch graph. *Indonesian Journal of Combinatorics*, 2(1), 14-19.
- 10] Anick, D. (2016). Counting graceful labelings of trees: A theoretical and empirical study. *Discrete Applied Mathematics*, 198, 65-81.
- 11] Lau, G. C., Shiu, W. C., & Ng, H. K. (2016). Further results on super graceful labeling of graphs. *AKCE International Journal of Graphs and Combinatorics*, 13(2), 200-209.