

Minireview Article

Cervical Cancer in Burkina Faso: Current Status, Government Strategies, and the Role of Traditional Medicine in Treatment

ABSTRACT

In Burkina Faso, cervical cancer is the second most common cancer among women, and the second leading cause of cancer-related death. Both incidence and mortality rates have risen steadily in recent years. Its natural history is linked to persistent infection with an oncogenic human papillomavirus, a necessary but not sufficient factor in its development. However, cervical cancer is said to be preventable, due to its slow evolution. To date, it is the only cancer that can be prevented by vaccination against human papillomaviruses (HPV). Systematic and regular screening enables early detection of precancerous lesions so that they can be treated, thereby preventing the onset of cervical cancer. Unfortunately, access to these preventive measures remains limited for the majority of the population due to cost. The government of Burkina Faso, through its Ministry of Health, has put in place strategies to facilitate access to screening and care. Burkina Faso's researchers and teacher-researchers are also collaborating with traditional health practitioners in the search for better remedies and active substances with no side effects, through the use of medicinal plants. This review aims to take stock of cervical cancer in Burkina Faso: achievements, shortcomings, strategies implemented by the government and the contribution of scientific research to the fight against this scourge.

Keywords: cervical cancer, human papillomavirus, screening, medicinal plants, Burkina Faso

1. INTRODUCTION

Cervical cancer is a major public health problem. It is a disease associated with persistent human papillomavirus (HPV) infection [1]. Around 89% of the 565,541 cases of cervical cancer reported worldwide in 2019 occurred in low- and middle-income countries[2]. According to GLOBOCAN estimates, in 2022, in Burkina Faso, 988 new cases of cervical cancer were diagnosed with 775 deaths[3]. Cervical cancer is the second leading cause of cancer mortality among women in Burkina Faso, with a mortality rate of 23.8 per 100,000 women [4]. In Burkina Faso, there is a lack of technical platforms and specialist healthcare staff in our hospital structures, particularly in rural areas, as well as a healthcare system that does not allow regular gynecological screening and care as it does in developed countries [4]. All these factors mean that screening is generally late, resulting in deaths in our countries. These results are all the more worrying given that these deaths are avoidable, and early diagnosis would enable rapid management and therefore cure [5]. Moreover, medical treatment is extremely difficult and expensive; hence its limited accessibility to the vast majority of the population suffering from this cancer [6].

Recent years have seen a surge in therapeutic research into cervical cancer, through the use of local medicinal plants derived from traditional medicine. Indeed, medicinal plants are a source of secondary metabolites with therapeutic properties. Several molecules used in cancer treatment originate from medicinal plants[7], [8]. Several ethnobotanical surveys have been carried out to identify the plants used by traditional practitioners to treat tumours[9]. The aim of this review is to take stock of cervical cancer in Burkina Faso: achievements, shortcomings, strategies implemented by the government and the contribution of scientific research to the fight against this scourge.

2. NATURAL HISTORY OF CERVICAL CANCER

Cervical cancer is a slowly evolving disease of infectious origin, which can take more than fifteen years to develop, from primary infection with an oncogenic human papillomavirus to the various precancerous histological lesions accompanying persistent infection [10]. Human papillomavirus (HPV) infection is a very common sexually transmitted infection[11]. Almost all cervical cancers begin in the epithelium. These cancers can appear either on the outside of the cervix (ectocervix) in 75-80% of cases (squamous cell carcinomas) or on the inside of the cervix (endocervix) in 20-25% of cases (adenocarcinomas) [12].

Persistent infection with high-risk oncogenic HPV (HR HPV) is recognized as the main cause of cervical cancer [1], and HPV infection can disappear (clearance) within a few months. In fact, some 80-90% of HPV infections are transient, disappearing spontaneously within 24 months of first detection [13]. It can also persist for several years (1 to 10 years), progressing to precancerous lesions or cervical intraepithelial neoplasia (CIN), which can be mild, moderate or severe (CIN 1, CIN 2, CIN 3). CIN 3 is a necessary stage for the development of invasive cancer, unlike CIN 1 and CIN 2. Cervical precancerous lesions may regress to normal epithelium, with a probability depending on lesion severity, accompanying viral clearance. They may also persist or progress to a more advanced stage, including CIN 3. The persistence of these lesions leads to the invasion of infected cells and thus to invasive cervical cancer [10].

Rarely, lesions may affect the glandular epithelium of the cervix, corresponding mainly to adenocarcinomas in situ, the only lesion preceding invasion. This lesion, already cancerous, is also due to persistent infection with a high-risk oncogenic HPV.

3. CAUSAL RELATIONSHIP BETWEEN HUMAN PAPILLOMAVIRUS (HPV) AND CERVICAL CANCER

Cervical cancer is caused by many factors. However, the primary and necessary cause is human papillomavirus infection [1]. Human papillomaviruses or HPVs (Human PapillomaVirus) are small (around 60 nm in diameter) naked viruses (without envelopes) of the *Papillomaviridae* family. In 2016, 205 different types of HPV were identified [14]. This virus is common, and the infection it causes is the most frequently diagnosed sexually transmitted infection (STI). The sexual route is the main route of HPV acquisition, although there are other routes [15], [16]. Almost all sexually active women have already been infected with HPV. But in the majority of cases, the infection disappears within six months to two years without symptoms of disease. However, in cases where the infection does not

clear and persists, the likelihood of developing precancerous lesions and subsequent cervical cancer is higher [15]. Precancerous and cancerous lesions of the cervix are caused by so-called high-risk oncogenic HPV types (HR-HPV): HPV 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59 and 68 [17]. Thus, persistent HR-HPV infection is the cause of nearly 100% of invasive cervical cancer cases, with HPV DNA detected in 90-100% of specimens from cervical lesions [18]. HR-HPVs are responsible for 7.7% of cancers in developing countries (DCs), mainly cervical cancer [19]. HPV types 16 and 18 are responsible for 50-60% and 10-20% respectively of cervical cancer cases encountered in most countries followed by HPV 45 and 31 with 4-8% and 1-5% respectively [18], [20]. Detection of HR-HPV in cervical samples is a highly sensitive tool for identifying women at risk of precancerous or cancerous cervical lesions. The prevalence is just as high in squamous cell carcinomas as in adenocarcinomas, although the distribution of HPV types varies slightly between these 2 histological forms [21]. Indeed, a worldwide study found 90.9% of HPV-positive women with squamous cell carcinoma, compared with 82.0% of those with adenocarcinoma. In contrast, the prevalence of multiple infections did not differ significantly primarily by histological type [22].

4. EPIDEMIOLOGY, RISK FACTORS AND HPV PREVALENCE IN CERVICAL CANCER IN BURKINA FASO

4.1 Epidemiology

Very few data are available on the prevalence, incidence and mortality due to cervical cancer in Burkina Faso due to the absence of a functional cancer registry. Nevertheless, it is estimated to be the second most common cancer (11%) after breast cancer (15.3%) among women. Between 1986 and 2006, it was estimated that cervical cancer accounted for 23% of cancers encountered in women [23]. According to GLOBOCAN data, cervical cancer represents 6.8 % of all cancers cases diagnosed in 2022 in Burkina Faso [3]. In 2022, 988 cases of cervical cancer were recorded with 775 deaths, which is 78.4% of diagnosed cases [3]. This shows that cervical cancer is a major public health problem in Burkina Faso.

4.2 Risk factors

Although HR-HPV infection is necessary, it is not sufficient to induce precancerous or cancerous lesions. This infection must persist for several years to lead to the appearance of precancerous lesions, and this persistence can be influenced by several factors [1]. These include environmental factors, use of oral contraceptives, co-infections (with HIV or several HPV types, for example) that weaken the immune system, smoking, frequency of re-infections...[24]. There are very few data on association between these different factors and the occurrence of cervical cancer in Burkina Faso. However, a study found a significant association ($p=0.01$) between the number of pregnancies and the occurrence of precancerous lesions and cervical cancer [25]. Another study on precancerous lesions found association between malphigian intraepithelial lesions of the cervix, HIV 1 co-infection and HPV multiple infection positivity [26]. Other cofactors such as age at first intercourse, smoking and the presence of other sexually transmitted infections were not associated with precancerous cervical lesions in this study[26]. In addition, some of these factors increase the risk of HPV infection, and therefore promote the development of cervical cancer. In Burkina Faso, these are: age, age at first sexual intercourse, marital status, level of education, occupation, number of sexual partners, frequency of sexual intercourse, non-use of condoms and history of sexually transmitted infections, use of contraceptives, HIV infection [27]–[29]. In contrast, other studies found no significant association between HPV positivity and polygamy, female genital mutilation, hormonal contraceptives, multiparity, level of education or number of sexual partners in the last 12 years [28], [30].

4.3 Prevalence of HPV infection among Burkinabe suffering from cervical cancer

Most studies in Burkina have focused on HPV prevalence in the general population in many regions of the country [31]–[34], in HIV-positive individuals [35]–[37], in pregnant women [38], in hepatitis infected women [39], precancerous lesions [26], [40], in cervical cancer [41], and in otorhinolaryngological (ORL) cancers[42]. Very few studies of women diagnosed with cervical cancer have been carried out in Burkina. One involved 112 formalin-fixed, paraffin-embedded archival cervical tissue blocks dated from 2009 to 2015 from women aged between 21 and 84 years. After molecular characterization of the 112 samples, 58.04% had a valid result (beta-globin detected). Of the valid results, 47 samples (72.31%) were HPV-positive. Overall, 11 of the 14 genotypes tested were identified in this study [41]. Of the genotypes found, the most common, in descending order, were HPV18 (25.71%), HPV31 (15.71%), HPV39 (12.86%), HPV16 (12.86%) and HPV45 (12.86%), HPV35 (7.14%) and HPV58 (5.71%) [41]. Among HR-HPV-positive samples, 65.96% of isolated infections and 34.04% of multiple infections were detected, with the number of genotypes ranging from 1 to 4 per individual [41]. Another, which concerns 100 samples (including 83 cervical cancer and 17 precancerous lesions) found 75.9% as HPV prevalence among women with cervical cancer[25]. A third study (review article) [43]synthetizes results of these two studies. The distribution of HR-HPV depending on the histological type of cervical cancer in this study is presented in **table I**. These prevalences of HPV in cervical cancer (72.31% and 75.9%) are lower than those found in Ghana (85.4%)[44] and Ivory Coast (89.4%) [45] and far from the 99.7% of HPV positives found in cervical cancer worldwide [46]. These geographical differences in HPV prevalence could be explained by differences in the number of samples or the method used to detect HPV.

Table I: distribution of HR-HPV depending on the histological type of cervical cancer[43]

Histological type	High risk HPV genotypes					Total
	HPV16	HPV18	HPV31	HPV33	Others*	
Squamous cell carcinoma	7 (5.15%)	28 (20.59%)	11 (8.10%)	20 (14.71%)	55 (40.44%)	121
Invasive adenocarcinoma	2 (1.47%)	3 (2.21%)	1 (0.74%)	4 (2.94%)	5 (3.68%)	15
Total	9	31	12	24	60	136

*Others: HPV 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.

5. NATIONAL STRATEGIES FOR THE PREVENTION AND MANAGEMENT OF CERVICAL CANCER IN BURKINA FASO

In Burkina Faso, cervical cancer is one of the major public health challenges. In response to this threat, the Burkina Faso government has put in place policies for the prevention, screening and treatment of cervical cancer. Cervical cancer screening plays an important

role in controlling this disease and reducing the number of deaths linked to it. To this end, the ministry of health has set up awareness-raising and free screening campaigns aimed at informing the population of the risks and measures to take to prevent cervical cancer. In addition, the government applies a policy of free care for women and children, which, since 2016, incorporates a free examination to detect the most common cancers including cervical and breast, and free treatment for precancerous cervical lesions [6]. In 2020, the World Health Organization (WHO) adopted a strategy to eliminate cervical cancer, based on three pillars:

- Vaccinate 90% of girls against HPV before the age of 15;
- Screen 70% of women with an effective test between the ages of 35 and 45;
- Treat 90% of women with precancerous lesions and 90% of women with invasive cancer[47]. In order to achieve the first objective, which is to vaccinate 90% of young girls, Burkina Faso, through the ministry of health, has introduced Gardasil 4 (a quadrivalent vaccine that protects against HPV 6, 11, 16 and 18) into the expanded vaccination program for girls aged 9 to 14 since April 26, 2022. Prior to this, a pilot project had been conducted between 2015 and 2017 in two health districts of the country (Baskuy and Solenzo) and 8487 9-year-old girls had been vaccinated [48]. In addition, in 2021, a radiotherapy center was opened at the Centre Hospitalier Universitaire (CHU) de Bogodogo in Ouagadougou to provide better care for cancer patients. Although this center was temporarily closed, the government again launched its reopening in March 2023 [49]. In addition to public sector efforts to improve cervical cancer screening and treatment, private initiatives and community health programs are also being implemented to support the government in the fight against cervical cancer. These include JHPIEGO (Johns Hopkins Program for International Education in Gynecology and Obstetrics) [50]. JHPIEGO is a non-profit Non-Governmental Organization (NGO) that works in several fields, notably in maternal and child health, including cervical cancer screening and treatment. Also, several Burkinabes associations such as KIMI, Espoir Cancer Féminin, ZONTA Club, etc, organize cervical cancer awareness and screening campaigns [4].

6. CERVICAL CANCER SCREENING AND TREATMENT IN BURKINA FASO

Cervical cancer screening in Burkina Faso is based mainly on visual inspection with acetic acid (VIA) with immediate management of precancerous lesions in CHU's, regional hospitals, medical centers and district hospitals since 2012 [51]. This method is an alternative to the Papanicolaou test (PAP test) which is not technically and financially feasible in Burkina Faso [52]. It is also the most recommended test in resource-limited settings [53]. Cervical cancer screening is also based on the cytological study of smears or biopsies of lesions and/or the HPV test, which uses a molecular biology technique to detect the presence of the virus in cervical cells. However, HPV test is not as commonly used as VIA test because of its cost (around 30 US dollars) [54]. It is mainly used in research and pilot projects like PARACAO (Partnership for Action and Research against Cervical Cancer in West Africa) [55], SUCCESS (Scale Up Cervical Cancer Elimination with Secondary Prevention Strategy project) [56], HARP (The HPV in Africa Research Partnership) [57] studies... Cervical cancer screening rates in Burkina Faso are currently low. A cross-sectional secondary analysis (data from all 13 regions in Burkina Faso) revealed that only 6.2% of women are being screened. The highest rates were observed in the "Hautsbassins" (19.2%) and "Centre" regions (14.7%) while the "Plateau central" (0%) and "Est" regions had the lowest rates (0 and 0.6% respectively). The screening frequency was 18.5% for urban areas and 2.8% for rural areas [58]. In 2020, it was estimated that only 6% of women in Western/Central Africa had

been screened for cervical cancer in their lifetime. Burkina Faso's screening coverage (4%) was higher than Benin (1%), Cote d'Ivoire (2%), and Ghana (3%), but lower than Senegal (8%). To achieve the target of screening 70% of women aged 35-45 years, Burkina Faso needs to screen at least 926,691 women [53]. The most common barriers to screening in Burkina Faso are poor knowledge[59], mainly lack of awareness about the disease, its screening, not knowing where to get screening, fear of being diagnosed with the disease, distance from the hospital, and financial difficulties[52], particularly for rural women.

Managing of positive screening results passed by cryotherapy, thermocoagulation or Diathermic Loop Resection (DLR) for precancerous lesions. There are also other types of treatment for cancer in general: surgery, which is a loop electrosurgical excision procedure (depending on the progress of the cancer, hysterectomy extended to the peripheral tissues of the uterus), chemotherapy (drug treatment) and radiotherapy (radiation treatment). These treatments are not accessible to all due to their high cost [6]. Cervical cancer treatment is very complex and expensive in Burkina Faso. For example, chemotherapy price ranges from 172 to 689 US\$ (depending on drugs and protocol used) every three weeks, carried out 6 times, with the possibility of further treatment. However, most people monthly salary is less than 85 US dollars[6]. Consequences are treatment refusals or treatment breaks. Radiotherapy is available in Burkina Faso but is centralized. Moreover, as in most countries in Sub Saharan Africa, most women are diagnosed in advanced stage, and the only suitable treatment options are palliative treatments. These factors are barriers to effective treatment of cervical cancer in Burkina Faso.

7. RESEARCH OF TREATMENTS USING MEDICINAL PLANTS IN BURKINA FASO

Many efforts are being made through research by scientists in Burkina Faso to enable the contribution of traditional medicine using medicinal plants in the fight against cancer in general and cervical cancer in particular [6]. Indeed, some plants have shown real anti-cervical cancer potential. Extracts of *Nelsoniaca nescens* showed quite interesting results on HeLa and SiHa cervical cancer cell lines, with the best activity observed for methanolic extracts whose half maximal inhibitory concentration (IC_{50}) were $21.72 \pm 0.08 \mu\text{g/ml}$ and $25.30 \pm 0.44 \mu\text{g/ml}$ respectively for the two cell types. In addition, these extracts showed high selectivity on HeLa and SiHa cancerous cells compared to normal L929 cells with a selectivity index of 7.93 ± 0.03 for methanolic extracts for HeLa cells and 7.48 ± 0.05 for ethyl acetate extracts for SiHa cells [60].

Similarly, another study showed that *Hyptissuaveolens* essential oil exerts a dose-dependent antiproliferative activity and acts on the cell cycle of HeLa cells by inducing G0/G1 cell cycle arrest and a decrease in the G2/M phase. However, cytometric analysis did not detect any changes in the subG1 population (linked to apoptosis) in the cancer cells examined and treated with essential oil. However, a significant peak close to the G0/G1 zone most likely related to necrotic cells on HeLa cells was observed [61]. The essential oils of *Cymbopogon nardus*, *Cymbopogon schoenanthus*, and *Lantana camara* also showed antiproliferative activities on cervical cancer HeLa cell lines with IC_{50} s of 142 ± 6.0 ; 46.17 ± 11.83 ; $229.27 \pm 11.25 \mu\text{g/ml}$ respectively for each essential oil [62]–[64]. Traditional recipes based on blends of several medicinal plants are also used in cervical cancer treatment research. We have, for example, the *Acti-plus* ($IC_{50}=362 \pm 27 \mu\text{g/ml}$)[65], *Cancerol S* ($EC_{50}=84.6 \pm 5.2 \mu\text{g/ml}$)[66], *paanfetiben 1* ($IC_{50}=112.02 \pm 0.03$), and *paanfetiben 2* ($IC_{50}>1000 \mu\text{g/ml}$)[67], recipes which have cytotoxic effect on HeLa cell lines. Figures 1 and 2 summarise effect of medicinal plants and traditional recipes on cervical cancer cell lines. The search for treatments also involves finding plants with antioxidant potential (cancer being linked to oxidative stress) and rich in anti-cancer substances. Thus, two plants,

Diospyros mespiliformis and *Daniellia oliveri*, used in traditional medicine for the treatment of tumors in two (2) towns in Burkina (Bobo Dioulasso and Fada N'Gourma) showed antioxidant potential by ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)), FRAP (Ferric ion reducing antioxidant power) and DPPH (2,2-diphenyl-1-picrylhydrazyl) methods and contained anticancer substances such as flavonoids and total phenolics [68].




Figure 1: Effect of medicinal plants on cervical cancer cell lines HeLa and SiHa




Figure 2: Effect of traditional recipes on cervical cancer cell lines HeLa

8. CONCLUSION

HPV-induced cervical cancer is a so-called preventable cancer. Indeed, HPV vaccination, early detection and treatment of cervical cancer are crucial to prevent the development and progression of the disease. To this end, Burkina Faso, through its Ministry of Health and in collaboration with non-governmental organizations, has set up awareness-raising strategies, free vaccination for girls' aged 9, regular screening campaigns and a radiotherapy center for cancer treatment. However, much remains to be done to combat this disease in this country, especially in terms of public acceptance of the vaccine. It is therefore essential to continue raising women's awareness of cervical cancer. In addition, the search for substances from local medicinal plants used in Burkina Faso to treat tumors could also be an alternative treatment for cervical cancer in this context of limited financial resources and lack of access to anti-cancer molecules.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

REFERENCES

- [1] Cubie HA. Diseases associated with human papillomavirus infection. *Virology*. 2013; 445(1–2): 21–34. doi: 10.1016/j.virol.2013.06.007
- [2] Momenimovahed Z, Mazidimoradi A, Maroofi P, Allahqoli L, Salehiniya H, Alkatout I. Global, regional and national burden, incidence, and mortality of cervical cancer. *Cancer Rep*. 2023;6(3):1–10. doi: 10.1002/cnr2.1756.
- [3] IARC/WHO. Cancer today, cancer incidence and mortality in Burkina Faso for 2022. 2024. Accessed: May 20, 2024. Available: <https://gco.iarc.fr/today/data/factsheets/populations/854-burkina-faso-fact-sheets.pdf>
- [4] Ministry of health. Strategic plan to fight cancer 2013 - 2017. 2013. Accessed 15 April 2022. Available: [https://www.iccp-portal.org/system/files/plans/Burkina-Faso_Plan strat%C3%A9gique de lutte contre le cancer 2013-2017.pdf](https://www.iccp-portal.org/system/files/plans/Burkina-Faso_Plan_strat%C3%A9gique_de_lutte_contre_le_cancer_2013-2017.pdf). French
- [5] Nour NM. Cervical Cancer: A Preventable Death. *Rev Obstet Gynecol*. 2009;2(4):240–4. doi: 10.3909/riog0100.
- [6] Couliati TH, Eric A, Sombié D, Bangou J, Kiendrébéogo M. Burden of cancer and role of traditional medicine in Burkina Faso. *Int J Complement Altern Med*. 2019;12(5):194–201. doi: 10.15406/ijcam.2019.12.00473.
- [7] Cragg GM, Grothaus GP, Newman DJ. Impact of Natural Products on Developing New Anti-Cancer Agents. *Chem Rev*. 2009;109:3012–43.
- [8] Newman DJ, Cragg GM. Natural Products As Sources of New Drugs over the 30 Years from 1981 to 2010. *J Nat Prod*. 2012;75:311–35. doi: dx.doi.org/10.1021/np200906s.
- [9] Thiombiano HM, Bangou MJ, Nacoulma AP, Ouoba B, Sawadogo M, Lema A, et al. Ethnobotanical Survey on Medicinal Plants Used in Burkina Faso in the Treatment of Breast Cancer, Phytochemistry and Antioxidant Activities: *Euphorbia poissonii* Pax and *Flueggea virosa* (Willd.) Voigt. (Euphorbiaceae). *African J Biol Med Res*. 2022;5(1):1–16. doi: 10.52589/ajbmr-udc9chlg.

- [10] Moscicki A barbara, Schiffman M, Kjaer S, Villa LL. Chapter 5 : Updating the natural history of HPV and anogenital cancer. *Vaccine*. 2006;24S3:S3/42-S351. doi: 10.1016/j.vaccine.2006.06.018.
- [11] Usyk M, Zolnik CP, Castle PE, Porras C, Herrero R, Gradissimo A, et al. Cervicovaginal microbiome and natural history of HPV in a longitudinal study. *PLoS Pathog*. 2020;16(3):1–20. doi: 10.1371/journal.ppat.1008376.
- [12] Smith HO, Tiffany MF, Qualls CR, Key CR. The rising incidence of adenocarcinoma relative to squamous cell carcinoma of the uterine cervix in the United States - A 24-year population-based study. *Gynecol Oncol*. 2000;78(2):97–105. doi: 10.1006/gyno.2000.5826.
- [13] Gravitt PE. The known unknowns of HPV natural history. *J Clin Invest*. 2011;121(12):4593–9. doi: 10.1172/JCI57149.
- [14] Harden ME, Munger K. HUMAN PAPILLOMAVIRUS MOLECULAR BIOLOGY. *Mutat Res*. 2017;772:3–12. doi: 10.1016/j.mrrev.2016.07.002.
- [15] Cutts FT, Franceschi S, Goldie S, Castellsague X, De Sanjose S, Garnett G, et al. Human papillomavirus and HPV vaccines: A review. *Bull World Health Organ*. 2007;85(9):719–26 doi: 10.2471/BLT.06.038414.
- [16] Krzowska-Firych J, Lucas G, Lucas C, Lucas N, Pietrzyk Ł. An overview of Human Papillomavirus (HPV) as an etiological factor of the anal cancer. *J Infect Public Health*. 2019;12(1):1–6. Available from: <https://doi.org/10.1016/j.jiph.2018.06.005> doi: 10.1016/j.jiph.2018.06.005.
- [17] Gamboa-Hoil SI. Human papillomavirus in men. *Rev Int Androl*. 2023;21(1):100325. doi: 10.1016/j.androl.2021.09.001.
- [18] Huber J, Mueller A, Sailer M, Regidor PA. Human papillomavirus persistence or clearance after infection in reproductive age. What is the status? Review of the literature and new data of a vaginal gel containing silicate dioxide, citric acid, and selenite. *Women's Heal*. 2021;17. doi: 10.1177/17455065211020702.
- [19] Aubry PP, Gaüzère DB alex. Les cancers dans les pays en développement Actualités 2017. *Med Trop*. 2018;1–7.
- [20] Bosch FX, De Sanjosé S. The epidemiology of human papillomavirus infection and cervical cancer. *Dis Markers*. 2007;23:213–27.
- [21] Bosch FX, Qiao Y lin, Castellsagué X. The epidemiology of human papillomavirus infection and its association with cervical cancer. 2006;94:8–21
- [22] Li N, Franceschi S, Howell-Jones R, Snijders PJF, Clifford GM. Human papillomavirus type distribution in 30,848 invasive cervical cancers worldwide: Variation by geographical region, histological type and year of publication. *Int J Cancer*. 2011;128(4):927–35. doi: 10.1002/ijc.25396.
- [23] Goumbri/Lompo OM, Domagni OE, Sanou AM, Konsegré V, Soudre RB. Aspects épidémiologiques et histopathologiques des cancers au Burkina Faso Epidemiological and histopathological aspects of cancers in Burkina Faso. 2009;(1):207–11. doi: 10.1007/s12558-009-0052-x. French
- [24] Gómez DT, Santos JL. Human papillomavirus infection and cervical cancer: pathogenesis and epidemiology. *Commun Curr Res Educ Top Trends Appl Microbiol*. 2007;680–8.
- [25] Sanou-Lamien A, Ki R, Zohoncon-Kone TM, Ouedraogo AS, Konsegré V, Ido FAHA, et al. Aspects socio-démographiques et histopathologiques des lésions précancéreuses de haut grade et cancéreuses du col utérin et caractérisation moléculaire des HPV oncogènes impliqués à Ouagadougou (Burkina Faso). *Rev. Africaine Pathol*. 2018; 17(2): 48–55. French.
- [26] Didelot-Rousseau MN, Nagot N, Costes-Martineau V, Vallès X, Ouedraogo A, Konate I, et al. Human papillomavirus genotype distribution and cervical squamous intraepithelial lesions among high-risk women with and without HIV-1 infection in Burkina Faso. *Br J Cancer*. 2006;95(3):355–62.

- [27] Zohoncon TM, Djigma WF, Karim A, Traore IMA, Ouedraogo RA, Traoré EMA, et al. Mapping of fourteen high-risk human papillomavirus genotypes by molecular detection in sexually active women in the West African sub-region. 2020;12(June):11–21. doi: 10.5897/IJGMB2019.0193.
- [28] Hatzipanagiotou M, Holzinger D, Millogo F, Pawlita M, Wacker J. Human papillomaviruses in Western Africa: prevalences and risk factors in Burkina Faso. *Arch Gynecol Obstet.* 2018;298:789–96. doi: 10.1007/s00404-018-4860-z.
- [29] Ouedraogo CMR, Djigma FW, Bisseye C, Sagna T, Zeba M, Ouermi D et al. Épidémiologie et caractérisation des génotypes de papillomavirus humain dans une population de femmes Ouagadougou. *J. Gynecol. Obstet. Biol. la Reprod.* 2011; 40(7): 633–638. doi: 10.1016/j.jgyn.2011.05.012. French
- [30] Ouédraogo CMR, Rahimy RML, Zohoncon TM, Djigma FW, Yonli AT, Ouermi D, et al. Épidémiologie et caractérisation des génotypes à haut risque de Papillomavirus humain dans une population d'adolescentes sexuellement actives à Ouagadougou. *J Gynecol Obstet Biol la Reprod.* 2015;44(8):715–22. doi: 10.1016/j.jgyn.2014.12.021. French
- [31] Ouedraogo RA, Zohoncon TM, Ouattara AK, and Simpore J. Prédominance du papillomavirus humain 56 dans une sous-population de femmes sexuellement actives à Garango, Centre-Est, Burkina Faso. *J. Appl. Biosci.* 2020; 150: 15499–15509. doi: 10.35759/jabs.150.10. French.
- [32] Traore IMA, Zohoncon TM, Djigma FW, Compaore TR, Traore Y, Simpore J. Association of TNF- α -308G/A and IL-18 Polymorphisms with risk of HPV infection among sexually active women in Burkina Faso. *Biomol Concepts.* 2020;11(1):97–101. doi: 10.1515/bmc-2020-0008.
- [33] Ouedraogo RA, Zohoncon TM, Guigma SP, Angèle Traore IM, Ouattara AK, Ouedraogo M, et al. Oncogenic human papillomavirus infection and genotypes characterization among sexually active women in Tenkodogo at Burkina Faso, West Africa. *Papillomavirus Res.* 2018;6(May):22–6. doi: 10.1016/j.pvr.2018.09.001.
- [34] Traore IMA, Zohoncon TM, Dembele A, Djigma FW, Obiri-Yeboah D, Traore G, et al. Molecular Characterization of High-Risk Human Papillomavirus in Women in Bobo-Dioulasso, Burkina Faso. *Biomed Res Int.* 2016;2016. doi: 10.1155/2016/7092583.
- [35] Djigma FW, ZOHONCON TM, DOUAMBA Z, SORGHO PA, Obiri-Yeboah D, OUATTARA AK, et al. Molecular Genotyping of Human Papillomavirus among HIV-infected and HIV- uninfected Women in Ouagadougou , Burkina Faso. *J Med Biomed App Sci.* 2020;8(1):324–33. doi: 10.15520/jmbas.v.
- [36] Djigma FW, Ouédraogo C, Karou DS, Sagna T, Bisseye C, Zeba M, et al. Prevalence and genotype characterization of Human Papillomaviruses among HIV-seropositive in Ouagadougou, Burkina Faso. *Acta Trop.* 2011;117(3):202–6. doi: 10.1016/j.actatropica.2010.12.007.
- [37] Sagna T, Djigma F, Zeba M, Bisseye C, Karou S, Ouermi D, et al. Human papillomaviruses prevalence and genital co-infections in HIV-seropositive women in Ouagadougou (Burkina Faso). *Pakistan J Biol Sci.* 2010;13(19):951–5. doi: 10.3923/pjbs.2010.951.955.
- [38] Kabre KM, Ouermi D, Zohoncon TM, Traore FPW, Gnoumou OP, Ouedraogo RA et al. doi: 10.1515/bmc-2022-0026.
- [39] Ouédraogo E, Zohoncon TM, Bayala B, Bado P, Ido-Da RPC, Ouedraogo RA et al. HPV / HBV or HPV / HCV Co-Infections in Women Treated for Chronic Hepatitis at Hôpital Saint Camille in Ouagadougou , Burkina Faso. *Am. J. Mol. Biol.* 2023. 14:1-12. doi: 10.4236/ajmb.2024.141001.
- [40] Ouédraogo C, Zohoncon TM, Traoré EMA, Ouattara S, Bado P, Ouedraogo CT, et al. Distribution of High-Risk Human Papillomavirus Genotypes in Precancerous Cervical Lesions in Ouagadougou , Burkina Faso. 2016;39(March):196–204.

- [41] Zohoncon TM, Bado P, Ouermi D, Traoré EMA, Djigma FW, Traore IMA, et al. Molecular Characterization of High-Risk Human Papillomavirus Genotypes Involved in Invasive Cervical Cancer From Formalin-Fixed , Formalin Paraffin-Embedded Tissues in Ouagadougou , Burkina Faso. *Int J Curr Res.* 2016;8(09):pp.39314-39318.
- [42] Ilboudo M, Zohoncon TM, Traore EMA, Traore IMA, Kande A, Djigma FW, et al. Characterization of high-risk oncogenic human papillomavirus genotypes in histologically confirmed ear, nose and throat (Ent) cancers in burkina faso. *Asian Pacific J Cancer Prev.* 2019;20(11):3429–35. doi: 10.31557/APJCP.2019.20.11.3429.
- [43] Zohoncon TM, Saoura EFT, Konsegré V, Ouedraogo E, Ouedraogo RA, Ouedraogo S et al. Génotypes HPV à haut risque impliqués dans les lésions précancéreuses et cancéreuses du col de l'utérus au Burkina Faso : revue systématique de littérature. *Sciences et Technique.* 2023; 46(1):125–141. French
- [44] Akakpo PK, Imbeah EG, Ulzen-Appiah K, Darkwa-Abrahams A, Adjei E, Amo-Antwi K et al. The distribution of hrHPV genotypes among cervical cancer cases diagnosed across Ghana: a cross-sectional study. *BMC Infect. Dis.* 2024;24(1):1–11. doi: 10.1186/s12879-024-09166-7
- [45] Boni SP, Tenet V, Horo A, Heideman DAM, Bleeker MCG, Tanon A. High-risk human papillomavirus distribution according to human immunodeficiency virus status among women with cervical cancer in Abidjan, Côte d'Ivoire, 2018 to 2020. *International Journal of Cancer.* 2024; 154(6):962-968. doi: 10.1002/ijc.34774.
- [46] Walboomers JMM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, et al. HUMAN PAPILLOMAVIRUS IS A NECESSARY CAUSE OF INVASIVE CERVICAL CANCER WORLDWIDE. *Thyroid Disord Basic Sci Clin Pract.* 1999;19(May):87–116. doi: 10.1007/978-3-319-25871-3_6.
- [47] World Health Organization, Global strategy to accelerate the elimination of cervical cancer as a public health problem and its associated goals and targets for the period 2020 – 2030, Vol. 2, United Nations General Assembly. 2020. 1–56.
- [48] WHO Burkina. Burkina Faso introduces vaccine against HPV (papillomavirus Humain) responsible of cervical cancer. 2022. Accessed 2023, Aug 22. Available from: <https://www.afro.who.int/fr/countries/burkina-faso/news/le-burkina-faso-introduit-le-vaccin-contre-le-hpv-papillomavirus-humain-responsable-du-cancer-du-col>. French
- [49] Government information service. Health infrastructure _ Bogodogo radiotherapy centre back in operation. 2023. Accessed 2023 Aug 22. Available from: <https://www.sig.bf/2023/03/infrastructures-sanitaires-le-centre-de-radiotherapie-de-bogodogo-reprend-du-service/>. French
- [50] Sawadogo B, Sheba GN, Rutebemberwa E, Sawadogo M, Meda N. Knowledge and beliefs on cervical cancer and practices on cervical cancer screening among women aged 20 to 50 years in Ouagadougou, Burkina Faso, 2012: A cross-sectional study. *Pan Afr Med J.* 2014;18:1–10. doi: 10.11604/pamj.2014.18.175.3866.
- [51] Ministry of health. National strategy for fight against cancer 2021-2025. 2020. Available from: <https://gco.iarc.fr/today/data/factsheets/populations/854-burkina-faso-fact-sheets.pdf>. French
- [52] Compaore S, Ouedraogo CMR, Koanda S, Haynatzki G, Chamberlain RM, Soliman AS. Barriers to cervical cancer screening in Burkina Faso: needs for patient and professional education. *J Cancer Educ.* 2016, 31(4):760-766. doi: 10.1007/s13187-015-0898-9.
- [53] Bruni L, Serrano B, Roura E, Alemany L, Cowan M, Herrero R et al. Cervical cancer screening programmes and age-specific coverage estimates for 202 countries and territories worldwide: a review and synthetic analysis. *Lancet Glob Heal.* 2022;10(8):e1115-e1127. doi: 10.1016/S2214-109X(22)00241-8.
- [54] Louie KS, De Sanjose S, Mayaud P. Review epidemiology and prevention of human

- papillomavirus and cervical cancer in sub-Saharan Africa : a comprehensive review. *Tropical Medicine and International Health* volume 2009;14(10):1287-1302. doi: 10.1111/j.1365-3156.2009.02372.x.
- [55] Mensah K, Kaboré C, Zeba S, Bouchon M, Duchesne V, Pourette D et al. Implementation of HPV-based screening in Burkina Faso: lessons learned from the PARACAO hybrid-effectiveness study. *BMC Women's Health*. 2021;21(1):1-16. doi: 10.1186/s12905-021-01392-4.
- [56] Kabue M, Gauvreau CL, Dacene N, Bertram MM, Shissler T, Reis V et al. Understanding integrated HPV testing and treatment of pre-cancerous cervical cancer in Burkina Faso, Cote d'Ivoire, Guatemala and Philippines: study protocol. *Reprod. Health*. 2023; 20(1):1–19. doi: 10.1186/s12978-023-01696-8.
- [57] Devine A, Vahanian A, Sawadogo B, Zan S, Bocoum FY, Kelly H et al. Costs and cost-effectiveness of cervical cancer screening strategies in women living with HIV in Burkina Faso: The HPV in Africa Research Partnership (HARP) study. *PLoS One*. 2021; 16(3): 31–41. doi: 10.1371/journal.pone.0248832.
- [58] Diendéré J, Kiemtoré S, Coulibaly A, Tougri G, Ily NI, and Kouanda S. Faible niveau de dépistage du cancer du col de l'utérus, disparités géographiques et déterminants sociodémographiques du dépistage chez les femmes adultes au Burkina Faso: résultats de la première enquête nationale en population générale. *Rev. Epidemiol. Sante Publique*. 2023; 71(4) :1-7. doi: 10.1016/j.respe.2023.101845. French
- [59] Tassemedo S, Winter CH, Traore IT, Ouattara A, Sawadogo M, and Meda N. Cervical pre-cancer screening by visual inspection of the cervix after application of acetic acid in rural Burkina Faso: evaluation of women's knowledge, screening practice habits, acceptability and prevalence of risk factors and lesions in Boussé health. *Pan Afr Med J*. 2023;45(135): 1-11. doi: 10.11604/pamj.2023.45.135.36933.
- [60] Ouattara N, Ramdé-Tiendrébeogo A, Guenné S, Sumera Z, Meda NTR, Konaté K, et al. Cytotoxicity Potential of *Nelsonia canescens* (Lam .) Spreng Extracts against Cervical Cancer Cell Lines. 2019;3214:439–43. doi: 10.36348/sjbr.2019.v04i12.008.
- [61] Bayala B, Nadembega C, Guenné S, Buñay J, Zohoncon TM, Djigma FW, et al. Chemical Composition , Antioxidant and Cytotoxic Activities of *Hyptis suaveolens* (L .) Poit . Essential Oil on Prostate and Cervical Cancers Cells. *Pakistan J Biol Sci*. 2020;23(9):1184–92. doi: 10.3923/pjbs.2020.1184.1192.
- [62] Bayala B, Coulibaly LL, Djigma F, Bunay J, Yonli A, Traore L, et al. Chemical Composition of Essential Oil of *Cymbopogon schoenanthus* (L.) Spreng from Burkina Faso, and Effects against Prostate and Cervical Cancer Cell Lines. *Molecules*. 2023 Jun 1; 28(11). PMC10254514
- [63] Coulibaly LL, Bayala B, Zongo L, Zongo PFI, Ouedraogo E, Djigma FW, et al. Chemical composition and antiproliferative activity on prostate and cervical cancer cell lines of *Lantana camara* Linn. essential oil. *Int J Biol Chem Sci*. 2023;17(2):293–303. doi: 10.4314/ijbcs.v17i2.1.
- [64] Bayala B, Coulibaly AY, Djigma FW, Nagalo BM, Baron S, Figueredo G, et al. Chemical composition, antioxidant, anti-inflammatory and antiproliferative activities of the essential oil of *Cymbopogon nardus*, a plant used in traditional medicine. *Biomol Concepts*. 2020 Jan 1;11(1):86–96. doi: 10.3390/MOLECULES28114561.
- [65] Adico MDW, Bayala B, Bazié BVEJT, Traoré L, Yonli AT, Simporé J. Cytotoxic Properties on Prostate and Cervical Cancer Cells in Culture of « Acti-plus », a Recipe Based on Medicinal Plants from Burkina Faso. *J. Dis. Med. Plants*. 2023; 9(4): 148–155. doi: 10.11648/j.jdmp.20230904.14.
- [66] Adico MDW, Bayala B, DjigmaFW, Bazié BVEJT, Darga P, Lobaccaro J-Met al. Antioxidant , Anti-inflammatory and Cytotoxic Properties of Cancerol S , A Blend of Medicinal Plants , on Cancerous Cells in Culture. *Annu. Res. Rev. Biol*. 2024;39(8): 87–98. doi: 10.9734/arrb/2024/v39i82112.
- [67] Ouédraogo E, Bayala B, Zohoncon TM, Ouédraogo S, Lompo OM, Dovo EE et

al. Cytotoxic Properties on Cervical and Liver Cancer Cells of Two Plant Recipes from Burkina Faso. *J. Biosci. Med.* 2024;12(06): 216–229. doi: 10.4236/jbm.2024.126019.

- [68] Ouoba B, Sessouma B, Bangou MJ, Thiombiano MH, Lema A, Sawadogo M, et al. Ethnobotanical survey on Medicinal Plants used in the treatment of tumors in two cities of Burkina Faso: Phytochemistry and Antioxidant activities. *Ijcm.* 2023;3(1):106–17. doi: 10.58349/IJCM.1.3.2023.00116.

UNDER PEER REVIEW