

Prevalence of *Acinetobacter baumannii* in Diyala governorate from Different Clinical Source

Comment [L1]: Add (s) to be sources, Then add to the title: and its Antibiotic profile analysis

Abstract

Acinetobacter baumannii bacteria from various clinical sources (Blood, urine, CSF, Ear swab, burn swab, wound swab, sputum and vaginal swab) were isolated from patients were attending to Baquba Teaching Hospital and Al-Betool Teaching Hospital. A total of (27) isolates of *A. baumannii* bacteria out of (200) samples were isolated from various clinical sources and the maximum isolates were obtained from blood samples 10 (37.03%), urine samples 4 (14.8%), CSF samples 4 (14.81%), burn swab (14.8%), sputum swab 2 (7.4%), wound swab 1 (3.7), Ear swab 1 (3.7) and vaginal swab 1 (3.7). All isolates were tested for their resistance to (23) different antibiotics and the results showed that resistant to Amoxicillin, Amoxicillin/Clavulanic acid and Trimethoprim were (96.2%) but lesser to Ticarcillin, Ticarcillin / Clavulanic Acid, Piperacillin/Tazobactam, Ceftriaxone, Imipenem, Ciprofloxacin and Levofloxacin were 88.8%.

Comment [L2]: delete

Comment [L3]: add: the diagnosis was achieved by vitek-2 apparatus

Comment [L4]: complete name

Comment [L5]: combine the percentage of urine, csf and burns in the same sentence.

Keywords: *Acinetobacter baumannii*, AST and Vitek 2 system

Comment [L6]: Complete term

Comment [L7]: Add Antibiotics

Introduction

Acinetobacter baumannii is a Gram-negative bacterium, aerobic and non-motile rod. This opportunistic bacterium has emerged as a significant concern in human health, responsible for a variety of infections, including pneumonia, meningitis, septicemia, and urinary tract infections. It possesses several virulence factors, such as the ability to form biofilms, adhere to surfaces, invade host cells, acquire iron, and induce host cell death [1]. *A. baumannii* easily attaches to both biological and non-biological surfaces, where it can form biofilms. This characteristic is significant for many pathogens as it aids in the colonization of prosthetic materials and plays a role in drug resistance and evasion of the host immune system in vivo [2]. *A. baumannii* has developed resistance to many of the most effective antimicrobial agents, leading to a significant increase in morbidity and mortality rates, particularly in intensive care units across various countries [3]. *Acinetobacter baumannii* is a significant multidrug-resistant (MDR) opportunistic pathogen commonly associated with nosocomial infections, largely due to its remarkable ability to develop resistance to various groups of antibiotics. Multiple mechanisms contribute to the acquisition of this multidrug resistance [4]. Resistance in *Acinetobacter baumannii* strains exhibiting the MDR phenotype is attributed to a diverse array of genes responsible for antibiotic resistance, encompassing both intrinsic and acquired mechanisms [5]. There are three mechanisms of antibiotic resistance in *A. baumannii*: First, resistance can arise by reducing membrane permeability or increasing antibiotic efflux, thereby preventing access to the target. Additionally, bacteria may employ genetic changes or post-translational modifications to protect the drug target. Finally, antibiotics can be directly inactivated through hydrolysis or modification processes [6]. This pathogen has developed multidrug resistance (MDR) in recent years, mainly due to the widespread overuse of antibiotics and inadequate antibiotic management. Prolonged hospital stays, use of catheters, and mechanical ventilation are associated with MDR isolates, and invasive infections are more likely to occur in immunocompromised individuals and those who are critically ill [7].

Comment [L8]: Put the abbreviation (*A. baumannii*)

Comment [L9]: Abbreviate and italic

Comment [L10]: Abbreviate and italic

Comment [L11]: italic

Comment [L12]: write down the aim of the study here

Materials and Methods

In the current study, 27 *A. baumannii* isolates were employed. Samples were gathered between January and July of 2024. Blood, urine, CSF, burns, wounds, blood cultures, and sputum were among the items that were taken from four Baghdad hospitals. All obtained samples, with the exception of blood, were grown immediately on blood agar and MacConkey agar and incubated for twenty-four hours at 37°C. Blood samples are cultured immediately on blood agar and MacConkey agar and then incubated for further 24 hours at 37°C. The blood specimen must be placed in blood culture bottles that include nutrients that encourage the growth of aerobes bacteria. All bacterial isolates were tested by Vitek 2 system. Bacterial identification based on the biochemical reactions between the bacterial suspension and the culture media existed in the GN-ID Cards loaded manually into the VITEK system. According to the manufacturer's instructions (BioMerieux, France), the following steps are performed automatically in the software of the system. The susceptibility of *Acinetobacter baumannii* isolates against 22 antibiotics were determined by Vitek 2 system by using Antibiotics Susceptibility Gram-negative (AST-GN).

Comment [L13]: italic

Comment [L14]: in the abstract was baquba

Comment [L15]: delete, the blood did not culture on agar immediately!!

Comment [L16]: delete

Comment [L17]: add: then were cultured on blood and MacConkey agar..... continue

Comment [L18]: which steps?????

Comment [L19]: Abbreviate and italic

Results

In this study a total of (200) samples were obtained from from (Blood, urine, CSF, Ear swab, burn swab, wound swab, sputum and vaginal swab as shown in table (1).

Table (1): Distribution of samples and number of *Acinetobacter baumannii* isolates

	Type of specimen	Total specimen	Number of isolates	%
<i>Acinetobacter baumannii</i> bacteria (27 isolated)	blood	30	10	37.03
	Urine	55	4	14.81
	CSF	15	4	14.81
	Ear swab	5	1	3.7
	Brun swab	30	4	14.81
	Wound swab	45	1	3.7
	Sputum swab	15	2	7.4
	vaginal swab	5	1	3.7
	Total	200	27	100

Comment [L20]: Delete the first column




Figure (1): Acinetobacter baumannii isolate distribution

Comment [L21]: Delete this figure, all the same data was written in the table

In this study most of *A. baumannii* isolates were highly resistant to the tested antibiotics as shown in table (2) and Figure (2).

Table (2): AST test on *A. baumannii* isolates

Comment [L22]: Write the complete term without abbreviation

AB	<i>Acinetobacter baumannii</i> bacteria (27 isolated)	
	S(100%)	R(100%)
Amoxicillin	1(3.7)	26(96.2)
Amoxicillin/Clavulanic acid	1(3.7)	26(96.2)
Ticarcillin	3(11.1)	24(88.8)
Ticarcillin / Clavulanic Acid	3(11.1)	24(88.8)
Piperacillin	6(22.2)	19(70.3)
Piperacillin/Tazobactam	3(11.1)	24(88.8)
Cefuroxime	2(7.4)	25(92.5)
Cefixime	2(7.4)	25(92.5)
Cefotaxime	2(7.4)	25(92.5)
Ceftriaxone	3(11.1)	24(88.8)
Ceftazidime	2(7.4)	25(92.5)
Cefepime	4(14.8)	23(85.1)
Imipenem	4(14.8)	24(88.8)
Meropenem	4(14.8)	23(85.1)
Gentamicin	5(18.5)	22(81.4)
Ciprofloxacin	3(11.1)	24(88.8)
Levofloxacin	3(11.1)	24(88.8)

Comment [L23]: Check the numbers, something wrong

Comment [L24]: Check the numbers, something wrong

Tigecycline	11(40.7)	16(59.2)
Minocycline	9(33.3)	18(66.6)
Colistin	9(33.3)	18(66.6)
Tobramycin	6(22.2)	21(77.7)
Trimethoprim	1(3.7)	26(96.2)
Trimethoprim/Sulfamethoxazole	10(37.03)	17(62.9)




Figure (2): AST test on *A. baumannii* isolates

Comment [L25]: Delete this figure, all the same data was written in the table




Figure (3): Increase Imipenem resistance of *A. baumannii* clinical isolates in Iraq

Comment [L26]: 1-Where did you get this information? This information should be referred to in the methodology!
2-It is supposed to indicate the fig. previously in the text of the results and talking about it , then display the fig.




Figure (4): Increase Colistin resistance of *A. baumannii* clinical isolates in Iraq

Discussion

The results shown in table (1) appeared that from the total 27 isolates 10(37.03%) from blood, 4(14.81%) isolated from Urine, CSF, Brun ,2(7.4%) from sputum respectively. The lowest number and percentage of *A.baumannii* was Wound, Ear, vaginal swabs 1(3.7%). According to this study, blood is where this bacterium most commonly causes infections. The results of this study agree with previous study [8]and this result differed with what the researcher reached [9]urine is often only detected when an indwelling urinary catheter is present. While it is infrequently linked to pyelonephritis and urosepsis, it is seldom invasive and typically just affects the lower urinary tract[10,11]. This result was compared to the result obtained by [12,13] and [14] have isolated *Acinetobacter baumannii* from urine rate reaching to (16,10% and 11.5%)respectively. The results of this study shows that *A. baumannii* bacteria can be isolated from burn and this result agreed withprevious study [15], who has isolated these bacteria from burn samples at rate of (17%).Around the world, burns are a serious public health problem and one of the most prevalent and destructive types of trauma[15,16]. The presence of an ample amount of oxygen in the respiratory tract, especially the lower part of it, encourages bacteria to settle and invade the region because it is a forced wind [17]. A bacteriological study stated that more than 60% of hospital-acquired pneumonia infections are caused by Negative bacteria including *Acinetobacter baumannii* the reason for this may be attributed to their ability to blind adhesion to epithelial cells in the respiratory tract as a result of inhibitory substances such as bacteriocins, was as the presence of mucous matter [18].

Because *A. baumannii* may form biofilms in the skin and cause infections in soft tissues, it is present in wounds and occlusive dressings [7]. Because many bacterial species have particular receptors for these chemicals, bacteria can readily colonize the surfaces of burn wounds.The reason for the difference in the isolation ratio is due to the number of samples taken the time of collection of the samples, the environment from which the samples were isolated, the health conditions in which the patients lived, and the length of time they stayed in Hospitalization, indiscriminate use of antibiotics excessively and the difference in the number of samples taken for a study and variance.Numerous dangerous nosocomial infections, including as urinary tract

Comment [L27]: 3-Where did you get this information? This information should be referred to in the methodology!
It is supposed to indicate the fig, previously in the text of the results and talking about it , then display the fig.

Comment [L28]: How are different? Write information

Comment [L29]: Write more about the isolation this bacteria from blood

Comment [L30]: Abbreviate and italic

Comment [L31]: italic

Comment [L32]: Abbreviate and italic

infections, meningitis, wound infections, bloodstream infections, and ventilator-associated pneumonia, can be caused by *A. baumannii*. Individuals in the high-risk groups include individuals who are mechanically ventilated, have indwelling foreign devices, are debilitating patients, and are hospitalized to intensive care units (ICUs) [19][20].

Comment [L33]: italic

Resistance to broad-spectrum antimicrobial agents in *A. baumannii* is now an emerging issue globally. In the present study, Tigecycline was found the most effective antimicrobial against isolated *A. baumannii* strains, with a 40.7% sensitivity rate, followed by trimethoprim-sulfamethoxazole by 37.03%, Minocycline and Colistin with a 33.3% while showed a high resistance rate to Amoxicillin, Amoxicillin/Clavulanic acid and Trimethoprim by (96.2%), Cefuroxime, Cefixime, Cefotaxime and Ceftazidime by (92.5%), Ticarcillin, Ticarcillin / Clavulanic Acid, Piperacillin/Tazobactam, Ceftriaxone, Ciprofloxacin and Levofloxacin (88.8%), and Cefepime, Imipenem, Meropenem, Gentamicin, Piperacillin, Tobramycin which exhibited the low level of antimicrobials effectiveness against *A. baumannii* strains.

Comment [L34]: italic

Comment [L35]: delete. these are results, must put in the results not in the discussion

The results of the current study demonstrated that the highest resistance to almost all β -lactam antibiotic classes under study was as follows cephalosporins (Amoxicillin, Amoxicillin/Clavulanic acid, Cefuroxime, Cefixime, Cefotaxime, Ceftriaxone, Ceftazidime, Cefepime, Piperacillin, Oxacillin). These findings were close to local studies related to *A. baumannii* isolates by [21] [22], who found that (100%) of *A. baumannii* isolates in Iraqi hospital environment resisted Oxacillin, Ceftazidime and Cefepime. Another study in Iran hospital by [23] found (100% resistance rate to (Cefoxitin, Cefoxitin, Oxacillin, Ceftazidime, Cefepime). The three main types of resistance mechanisms are: (1) enzymes that deactivate antimicrobials; (2) bacterial targets are less accessible; or (3) mutations that alter targets or cellular activities. Regarding the first group, a variety of β -lactamases found in *Acinetobacter* species hydrolyze and provide resistance to cephalosporins, carbapenems, and penicillins. Chromosome-encoded AmpC cephalosporinases provide resistance to broad-spectrum cephalosporins [24].

As the isolates of bacteria *A. baumannii* resistance to Gentamicin in the percentage (81.4%). The results are consistent with [25], which was reached by the resistance rate (62%). While the resistance to isolates of the antibiotic Imipenem (88.8%), Meropenem (85.1%). This close with the results of the current study, which gave a 100% resistance to this antibiotic.

The results of Ciprofloxacin and Levofloxacin are agreed with [26] research, which was reached by the resistance rate (85%), and also agreed with [27] research. The resistance rate was (80%). Another local study revealed that *A. baumannii* clinical isolates were 100% resistant to Ciprofloxacin [28]. resistance of *A. baumannii* for Tobramycin, were (77.7%) consistent with the study of [29], that showed 100% resistance for Tobramycin.

Because of its low rates of resistance, tigecycline could be the treatment of choice for managing infections at this facility. Although The results of resistance to trimethoprim /sulphamethoxazole also had a low resistance rate in our present study, were close to the results reached by the researcher in the research [27] and also close to the results of the research of [30] who recorded that the *Acinetobacter baumannii* resist to a wide range of antibiotic including trimethoprim / sulphamethoxazole in the percentage (81%). fewer patients were treated by this antibiotic, as its side effects of nephrotoxicity and neurotoxicity [31].

These bacteria are resistant to nearly all antibiotics for a variety of reasons. Among these, the most significant one is the existence of efflux pump families [32]. There are several families of efflux pumps in *Acinetobacter baumannii*, but the most significant family is the existence of Resistance. These families allow us to exclude antibiotics, heavy metals, and a host of other substances. The three pump groups that make up the nodulation cell division (RND) family are each in charge of resisting a particular kind of antibiotics. Clinical isolates of *A. baumannii* were reported to be 37% resistant to colistin in a previous investigation conducted in Sudan in 2015 (33). Previous study in Iraq in 2016 found that *A. baumannii* clinical isolates were 20% resistant to colistin (34). Previous study has reported that *A. baumannii* from clinical isolates were 96% resistant to colistin (35). Colistin-resistant *A. baumannii* may become resistant to colistin due to modifications of the outer membrane which may increase the permeability to other cell wall antimicrobial agents. Previous studies have reported that colistin-resistant *A. baumannii* strains were more susceptible to other antimicrobial agents than colistin-susceptible strain (36).

Previous study in Iraq in 2012 shown that resistance to Imipenem was 57.1% (37). Previous study in Iraq in 2014 shown that low resistance to Imipenem 30% (38). Other study also in Iraq shown that One (10%) isolate was observed to be imipenem and meropenem resistant (39). Iraqi study in 2017 found that *A. baumannii* clinical isolates were 68.7% and 50% resistant to Ceftazidime and Imipenem respectively (40). *A. baumannii* clinical isolates in previous study were 68.7% and 90.9% resistant to Imipenem (41,42).

In present study appeared to have resistance to Cefotaxime (92.5%) compare to previous study shown that all isolates appeared to have resistance to Cefotaxime (60%) (Jabur, 2014). Previous study have reported that most isolates of *A. baumannii* are considered to be resistant to Cefotaxime in rate reach to (80%) and (60%) respectively (43,44). Other previous study found that (12.2%) of isolates were resistant to Cefotaxime (45).

Conclusion

A. baumannii has demonstrated a high level of resistance to the majority of tested antibiotics, earning it the reputation of being the most therapeutically significant *Acinetobacter* species globally. It has become a significant nosocomial opportunistic pathogen in hospital infection epidemics.

Reference

- 1- Kunz, A and Brook, J. (2010) Emerging resistant gram-negative aerobic bacilli in hospital acquired infections. *Chemotherapy*; 56: 492–500.
- 2- Lee, H.W., Kah, Y.M., Kim, J., Lee, J.C., Seol, S.Y., Cho, D.T. (2008). Capacity of multidrug-resistant clinical isolates of *Acinetobacter baumannii* to form biofilm and adhere to epithelial cell surfaces. *Clin. Microbiol. Infect.* 14:49–54.
- 3- Moulana, Z., Babazadeh, A., Eslamdost, Z., Shokri, M., & Ebrahimpour, S. (2020). Phenotypic and genotypic detection of metallo-beta-lactamases in Carbapenem resistant *Acinetobacter baumannii*. *Caspian journal of internal medicine*, 11(2), 171.

Comment [L36]: 1-it was written in many pattern, so you have to follow the journal instructions
2- Write down recent references and delete some old references.

- 4- Da Silva, G. J., & Domingues, S. (2016). Insights on the horizontal gene transfer of carbapenemase determinants in the opportunistic pathogen *Acinetobacter baumannii*. *Microorganisms*, 4(3), 29.
- 5- Fallah, A., Rezaee, M. A., Hasani, A., Barhaghi, M. H. S., & Kafil, H. S. (2017). Frequency of *bap* and *cpaA* virulence genes in drug resistant clinical isolates of *Acinetobacter baumannii* and their role in biofilm formation. *Iranian journal of basic medical sciences*, 20(8), 849.
- 6- Wong, D., Nielsen, T. B., Bonomo, R. A., Pantapalangkoor, P., Luna, B., & Spellberg, B. (2017). Clinical and pathophysiological overview of *Acinetobacter* infections: a century of challenges. *Clinical microbiology reviews*, 30(1), 409-447.
- 7- Pamu, J. (2019). *Prevalence and Antimicrobial Susceptibility Pattern of Methicillin-resistant Staphylococcus Aureus at Port Moresby General Hospital, Papua New Guinea: A Retrospective Study* (Doctoral dissertation, Auckland University of Technology).
- 8- GÜLEN, T. A., Ayfer, İ. M. R. E., ÖDEMİŞ, İ., & KAYABAŞ, Ü. (2020). *Acinetobacter baumannii* Infections and Antibiotic Resistance in Hospitalized Patients in an Education and Research Hospital: A Six-Year Analysis.
- 9- S. M. Kadom, I. N. ABID, Detection of *bla* OXA-51-like and *bla* VIM Carbapenemase Genes in *Acinetobacter baumannii* isolated from burn patients, *International Journal of Pharmaceutical Research*, 12(2), (2020)
10. Kb, L. (2002). Incidence and risk factors for acquiring nosocomial urinary tract infection in the critically ill. *J Crit Care*, 17, 50-57.
11. Avecillas JF, Mazzone P, Arroliga AC.(2003) A rational approach to the evaluation and treatment of the infected patient in the intensive care unit. *Clin Chest Med*; 24:645-69.
12. Rodríguez-Baño, J., Cisneros, J. M., Fernández-Cuenca, F., Ribera, A., Vila, J., Pascual, A., ... & Grupo de Estudio de Infección Hospitalaria. (2004). Clinical features and epidemiology of *Acinetobacter baumannii* colonization and infection in Spanish hospitals. *Infection Control & Hospital Epidemiology*, 25(10), 819-824.
13. Rungruanghiranya, S., Somboonwit, C., & Kanchanapoom, T. (2005). *Acinetobacter* infection in the intensive care unit. *Journal of Infectious Diseases and Antimicrobial Agents*, 22(2), 77-92.
14. Yaman, A. (1998) Antibiotic resistance in *Acinetobacter* species in nosocomial and outpatient infections. *Ann. Med. Sci.* 7 : 31 – 34.
15. Qader, AR., Muhamad, JA. (2010) Nosocomial infection in sulaimani burn hospital, Iraq. *Ann Burns Fire Disasters*.(4):177-81.
16. Ezzatollah, R., Hosain, S., Mahbobe, N. and Hassan, A. (2011) Common pathogens in burn wound and changes in their drug sensitivity. Volume 37, Issue 5, Pages 805-807.

17. Lopes, B. S., Gallego, L., & Amyes, S. G. B. (2013). Multi-drug resistance profiles and the genetic features of *Acinetobacter baumannii* isolates from Bolivia. *The Journal of Infection in Developing Countries*, 7(04), 323-328.
18. MacFaddin, J. F. (2000). Biochemical tests for identification of medical bacteria, Williams and Wilkins. *Philadelphia, PA*, 113(7).
19. Akrami, F., & Namvar, A. E. (2019). *Acinetobacter baumannii* as nosocomial pathogenic bacteria. *Molecular Genetics, Microbiology and Virology*, 34, 84-96.
20. Lerner, A. O., Abu-Hanna, J., Carmeli, Y., & Schechner, V. (2020). Environmental contamination by carbapenem-resistant *Acinetobacter baumannii*: the effects of room type and cleaning methods. *Infection Control & Hospital Epidemiology*, 41(2), 166-171.
21. Adnan, S. (2024). Identification of *Acinetobacter baumannii* Isolated from Different Infections and Study the Prevalence of Antibiotic Resistance in Patients of Baquba City. *Academic Science Journal*, 2(2), 91–104. <https://doi.org/10.24237/ASJ.02.02.712A>.
22. Al-Kadmy, I. M. S., Ali, A. N. M., Salman, I. M. A., & Khazaal, S. S. (2018). Molecular characterization of *Acinetobacter baumannii* isolated from Iraqi hospital environment. *New microbes and new infections*, 21, 51-57.
23. R. Maryam, Y. Z. Golnaz, O. Mojgan, T. Malihe, A. Nour, Identification of five phylogenetic groups of carbapenemase (blaOXA-23,24,51,58,143) in *Acinetobacter baumannii* strains isolated from clinical samples in Iran by multiplex PCR, *Der Pharma Chem*, 7(7), 11-6(2015).
24. Tien, H.C., Battad, A., Bryce, E.A. (2007) Multi-drug resistant *Acinetobacter* infections in critically injured Canadian Forces soldiers. *BMC Infect. Dis* ;7:95.
25. Lin, M. F., & Lan, C. Y. (2014). Antimicrobial resistance in *Acinetobacter baumannii*: From bench to bedside. *World Journal of Clinical Cases: WJCC*, 2(12), 787.
26. Al-Rifai, S.B.; Mohamad, W.S. and Jasem, N.H. (2022) Surveillance of *Acinetobacter* isolated from urinary tract infection at a Baghdad urology center. *Journal of Medicinal and Chemical Sciences*. JMCS-2201-1409.
27. Ghaima, K. K. (2016). Study of OXA beta-lactamase Genes in Clinical Isolates of Multidrug Resistant *Acinetobacter baumannii*.
28. Al-Khafaji, S.M.S. (2006). Study on capsule of *Acinetobacter baumannii* and its effect on Immune Response. Ph.D. Thesis. Biology department. College of Sciences. AL-Mustansiriyah University.
29. Al-Kadmy, I. M. S., Ali, A. N. M., Salman, I. M. A., & Khazaal, S. S. (2018). Molecular characterization of *Acinetobacter baumannii* isolated from Iraqi hospital environment. *New microbes and new infections*, 21, 51-57.
30. Angoti, G., Bandehpour, M., Goudarzi, H., Hajizadeh, M., Zarringhalam Moghaddam, M., and Kouchaki, A. (2016). Detection of Efflux Pump Genes (adeA, adeB, adeC and abeM) in

Acinetobacter baumannii Isolated from Hospitalize Patients, North-west of Iran. *Infection, Epidemiology and Microbiology*, 2(4), 8-11.

31- Loho, T., & Dharmayanti, A. (2015). Colistin: an antibiotic and its role in multiresistant Gram-negative infections. *Acta Medica Indonesiana*, 47(2).

32- Monem, S., Furmanek-Blaszczak, B., Łupkowska, A., Kuczyńska-Wiśnik, D., Stojowska-Swędrzyńska, K., & Laskowska, E. (2020). Mechanisms protecting *Acinetobacter baumannii* against multiple stresses triggered by the host immune response, antibiotics and outside-host environment. *International Journal of Molecular Sciences*, 21(15), 5498.

33- Omer, M. I., Gumaa, S. A., Hassan, A. A., Idris, K. H., Ali, O. A., Osman, M. M., ... & Khaled, M. M. (2015). Prevalence and resistance profile of *Acinetobacter baumannii* clinical isolates from a private hospital in Khartoum, Sudan. *Am J Microbiol Res*, 3(2), 76-9.

34-Al-Samaree, M. Y., & Al-Khafaji, Z. M. (2016). Antibiogram of *Acinetobacter baumannii* isolated from Baghdad Hospitals. *Int. J. Adv. Res. Biol. Sci*, 3(4), 238-242.

35-Narjis, M. A., & Mahdi, M. S. (2023). Isolation and identification of multi-drug resistance *Acinetobacter baumannii* isolated from clinical samples at Baghdad, Iraq. *Journal of Applied and Natural Science*, 15(2), 663-671.

36-Li, J., Nation, R. L., Owen, R. J., Wong, S., Spelman, D., & Franklin, C. (2007). Antibigrams of multidrug-resistant clinical *Acinetobacter baumannii*: promising therapeutic options for treatment of infection with colistin-resistant strains. *Clinical infectious diseases*, 45(5), 594-598.

37-Shali, A. A. (2012). Identification of Multidrug-Resistant Genes in " *Acinetobacter baumannii*" in Sulaimani City-Kurdistan Regional Government of Iraq. *Asian Journal of Medical Sciences*, 4(5), 179-183.

38-Jabur, M. H. (2014). Isolation of *Acinetobacter baumannii* from different clinical source and study some antibiotic resistant and β -lactamase production. *Med J Babylon*, 11(2), 456-464.

39-AL-Harmoosh, R. A., Jarallah, E. M., & AL-Shamari, A. M. (2016). Coexistence of the blaIMP and blaSIM genes in clinical isolates of *Acinetobacter baumannii* IN Babylon Hospitals-Iraq. *International Journal of PharmTech Research*, 9(7), 257-264.

40- Al-Dulaimi, A. A. F., Al-Taai, H. R. R., & Al-Bajlany, S. M. M. (2017). Virulence Factors of *Acinetobacter baumannii* isolated from different clinical specimens in Baquba.

41-Ibrahim ME. Prevalence of *Acinetobacter baumannii* in Saudi Arabia: Risk factors, antimicrobial resistance patterns and mechanism of carbapenem resistance. *Ann Clin Microbiol Antimicrob* 2019; 18:1.

42-Mahmood, N. H., & Al-Brefkani, A. M. T. (2023). Detection and Characterization of Carbapenem Resistant *Acinetobacter baumannii* Isolated from Different Clinical Specimens in Duhok Province-Iraq. *History of Medicine*, 9(1), 783-794.

43-Madanan, A., Yadav, D., Madhavan, R.(2009) Imipenem resistance and biofilm production in Acinetobacter. Department of Microbiology, SRM Medical College, Tamil Nadu, India.

44-Nazmul, M.H., Jamal, H., Fazlul, M.K.(2012) Acinetobacter species associated infections and their antibiotic susceptibility profiles in Malaysia. Department of Microbiology, Faculty of Medicine, Universiti Teknologi MARA, Shah Alam, Malaysia. Biomed Res- India., 23 (4): 571-575.

45-Shareek, P.S., Sureshkumar, D., Ramgopalakrishnan, V., Ramasubramanian, K.A.(2012) Antibiotic sensitivity pattern of blood isolates of Acinetobacter species in a tertiary care hospital: A retrospective analysis. Am. J. Infect. Dis., 8: 65-69.

UNDER PEER REVIEW