

## Original Research Article

# Variability and Character Relationships in F<sub>4</sub> progenies for Seed Yield in Black Gram [*Vigna mungo* (L.) Hepper]

### Abstract

Black gram is a vital pulse crop, used for its high protein content and nutritional value. It plays a crucial role in diverse culinary traditions and contributes significantly to food security. The objective of this research was to assess the variability resulting from hybridisation for F<sub>4</sub> generation by employing multiple parameters, comprising variability, correlation and path analysis. In this study, 100 F<sub>4</sub> progenies resulting from the cross between NUK-15-02 and NUK-15-07 were assessed using Augmented Block Design, alongside four standard check varieties. High GCV and PCV values observed for traits such as branches per plant, clusters per plant, pods per plant, and seed yield per plant (g). High heritability coupled with high genetic advance as per cent of mean was recorded for branches per plant, clusters per plant, pods per plant, pod length (cm), 100 seed weight (g), days to maturity, harvest index (%) and seed yield per plant (g). Seed yield per plant (g) exhibited strong positive associations with plant height (cm), pod length (cm), seeds per pod, clusters per plant, pods per plant, and harvest index (%). Path analysis highlighted that the highest positive direct effects on seed yield per plant were attributed to pods per plant, followed by clusters per plant and seeds per pod. The study highlights significant variability among black gram progenies, which can guide breeding programs to enhance yield and traits through targeted selection.

**Keywords:** variability, correlation analysis and path analysis, plant breeders, Black gram

### Introduction

Pulses, often referred to as the "Poor Man's Meat" (Nunes *et al.*, 2006), play a crucial role as a primary source of dietary protein for vegetarians. The growing awareness of health and nutrition among people has drawn the attention of plant breeders toward enhancing pulse production rates while emphasizing nutritional aspects. "In India, the total pulse production for the year 2021-22 is estimated at 27.69 million tonnes" (Anonymous, 2022). "Legumes, including pulses, are well-suited to agronomic cropping systems, providing nutritionally significant crops with high economic value, and they also contribute to soil fertility improvement by fixing atmospheric nitrogen. Among the pulse crops, black gram [*Vigna mungo* (L.) Hepper, 2n = 22] holds a prominent position and is cultivated worldwide. Black gram is a self-pollinated, short-duration, fast-growing warm-weather crop, primarily grown in rice fallow areas under rainfed conditions. The major challenges in achieving higher productivity in black gram cultivation include the lack of suitable crop types, limited genetic variability, a narrow genetic base (resulting in a higher degree of relatedness among varieties), and susceptibility to both biotic and abiotic stresses" (Souframanien *et al.*, 2004). "Genetic variability is a crucial resource for improving yield and other complex polygenic traits" (Swamy *et al.*, 2004). "The success of selection depends on the extent and nature of genetic variability within the population" (Johnson *et al.*, 1995). Correlation coefficients provide valuable insights into the direction and strength of the associations and interactions between yield and its components (Vyas *et al.*, 2018), facilitating the selection of the most suitable combinations of traits for improving black gram yield (Kumar *et al.*, 2003). "Path coefficient analysis is designed to quantify the interrelationships between yield components and their direct and indirect effects on seed yield" (Sushmitharaj *et al.*, 2018). "For crop improvement, understanding genetic variability for specific economic traits, their heritability, and genetic advances are crucial in developing high-yielding black gram genotypes in breeding programs" (Rao *et al.*, 2006). Given the current status of black gram cultivation and future prospects, plant breeders aim to develop superior germplasm with stable and high yield

potential, along with broader adaptability to different cultivation regions. In pursuit of characterizing superior germplasm, the present research project, titled "Variability and Character Relationships in F<sub>4</sub> progenies for Seed Yield in Black Gram [*Vigna mungo* (L.) Hepper]," was designed to assess the significant variability of agro-morphological traits that can be harnessed for future agricultural improvement.

### Materials and Methods

The experiment was conducted at the Research Farm of the Department of Genetics and Plant Breeding, Navsari Agricultural University, Navsari, during the late *khari*f season of 2022-2023. A total of hundred F<sub>4</sub> progenies, resulting from the cross between NUK-15-02 and NUK-15-07, were used for the study. The experimental design employed was the Augmented Block Design, with four standard checks, namely NUK-15-02, NUK-15-07, GU-1, and GU-3, included in the experiment. These checks were replicated in five blocks such that in each row, there were 20 plants of a single progeny, with a spacing of 20 cm between rows and 10 cm between individual plants within the row. Standard agronomic practices were meticulously followed to cultivate healthy and uniform crops. Observations were recorded for ten randomly selected plants from each progeny, excluding the border plants. For the traits days to 50% flowering and days to maturity, data were collected on a population basis. A total of eleven quantitative traits were assessed, including days to 50% flowering, plant height (cm), branches per plant, clusters per plant, pods per plant, seeds per pod, pod length (cm), 100 seed weight (g), seed yield per plant (g), days to maturity and harvest index (%). The collected data were subjected to standard statistical analyses to compute various parameters related to genetic variability. These parameters included the genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV) (Burton, 1952), heritability (Allard, 1960), genetic advance (Johnson, 1955), and correlation and path analysis (Dewey and Lu, 1959). This research aimed to provide valuable insights into the genetic diversity and interrelationships between various traits in black gram progenies, contributing to the development of improved varieties with higher yield potential and adaptability for different regions of cultivation.

### Results and Discussion

The variability parameters for various traits in F<sub>4</sub> progenies of black gram was presented in (Table 1, Fig 1 and Fig 2). The genetic coefficient of variation (GCV) ranged from 3.94 (for days to 50% flowering) to 37.79 (for pods per plant), while the phenotypic coefficient of variation (PCV) ranged from 4.09 (for days to 50% flowering) to 39.05 (for single plant yield). High GCV and PCV values were observed for branches per plant (32.29 and 32.64), clusters per plant (36.08 and 37.26), pods per plant (37.79 and 38.17), and seed yield per plant (34.70 and 39.05), respectively. These findings align with previous studies by Panigrahi *et al.*, (2014), Gowsalya *et al.*, (2016) and Patel and Madhu Bala (2020b). Conversely, days to 50% flowering (3.94 and 4.09) and seeds per pod (7.97 and 9.86) exhibited low GCV and PCV values, similar with observations of Panigrahi *et al.*, (2014), Gowsalya *et al.*, (2016) and Bishnoi *et al.*, (2017). Pod length displayed moderate values for GCV (10.31) and PCV (10.83), while plant height (9.48 and 14.72) and days to maturity (9.97 and 10.0) exhibited low GCV and moderate PCV, consistent with the findings of Kiran and Lal (2021). For 100 seed weight (19.33 and 20.25) and harvest index (15.93 and 20.34), moderate GCV and high PCV were observed, which are in par with results reported by Panda *et al.*, (2017). The relatively high genetic advance as percent of mean in conjunction with heritability, presents a promising opportunity for the improvement of all yield-related traits.

The outcomes of this study demonstrate the significance of heritability and its implications for genetic improvement in black gram. The heritability values ranged from 41.48 % for plant height to 99.29 % for days to maturity. Notably, traits like pods per plant (98.03% and 43.85%), pod length (90.62% and 20.20%), branches per plant (97.83% and 65.88%), clusters per plant (93.75% and 72.07%), 100 seed weight (91.07% and 38.05%),

days to maturity (99.29% and 20.50%), harvest index (61.35% and 25.74%), and seed yield per plant (78.95% and 63.60%) displayed both high heritability and high genetic advance as percent of mean, suggesting a substantial role of additive gene effects and making them effective for selection. Similar results have been reported by Panigrahi *et al.*, (2014), Gowsalya *et al.*, (2016), and Kondagari *et al.*, (2017). In contrast, traits such as seed per pod exhibited high heritability (65.32% and 13.28%) coupled with moderate genetic advance as percent of mean, indicating that these traits are primarily under genetic control and can be improved through phenotypic selection. This pattern aligns with findings reported by Panigrahi *et al.*, (2014) and Kondagari *et al.*, (2017). Conversely, days to 50% flowering showed high heritability (92.88%) but low genetic advance as percent of mean (7.83%), suggesting less environmental influence on this trait. Plant height exhibited moderate heritability (41.48%) and genetic advance as per cent of mean (12.59%), indicating the potential involvement of non-additive gene action. Similarly, high heritability coupled with low genetic advance as per cent of mean implies the presence of non-additive gene action.

Correlation analysis reveals the nature and strength of relationships between yield-related traits. Phenotypic correlation coefficients (Table 2) were calculated for 11 traits in F<sub>4</sub> progenies of black gram to assess their associations with seed yield per plant. Seed yield per plant displayed significant and positive correlations with plant height (0.281), clusters per plant (0.273), and pods per plant (0.526), consistent with findings from studies by Kumar *et al.*, (2014), Panigrahi *et al.*, (2014), Gowsalya *et al.*, (2016) and Patel and Madhu Bala (2020a). Additionally, positive and significant correlations were observed with pod length, seeds per plant, and harvest index similar to the correlation patterns identified by Panigrahi *et al.*, (2014) for pod length. These correlation findings suggest that improving seed yield in black gram would benefit from selecting plant types with high plant height, more clusters per plant, and more pods per plant. Furthermore, traits such as pod length, seed per pod, and harvest index can also serve as selection criteria for enhancing seed yield in black gram.

Path coefficient analysis provides insights into their direct and indirect effects on yield. The results (Table 3 and Fig.3) of path coefficient analysis revealed that days to 50% flowering exhibited a non-significant but negative correlation with seed yield per plant (-0.078) and had a negative direct effect on seed yield per plant (-0.1376). However, it exerted positive indirect effects on clusters per plant, pods per plant, pod length, seeds per pod, 100 seed weight, days to maturity, and harvest index. Similar negative and direct effects were reported in previous studies by Sathees *et al.*, (2019) and Rana *et al.*, (2022). Plant height showed a positive and highly significant correlation with seed yield per plant (0.281) and had a positive direct effect on seed yield per plant (0.0655), similar with the positive and direct effects reported by Kumar *et al.*, (2014), Partap *et al.*, (2019) and Sathees *et al.*, (2019). Branches per plant exhibited a positive but non-significant correlation with seed yield per plant (0.179) and had a negative direct effect on seed yield per plant (-0.2496). It exerted positive indirect effects on plant height, clusters per plant, pods per plant, days to maturity and harvest index. This pattern agreed with previous findings by Kumar *et al.*, (2014) and Sathees *et al.*, (2019). Clusters per plant displayed a positive and highly significant correlation with seed yield per plant (0.273) and had a positive direct effect on seed yield per plant (0.2231), consistent with positive direct effects reported by Kumar *et al.*, (2014), Sathees *et al.*, (2019), Kiran and Lal (2021) and Rana *et al.*, (2022). Pods per plant exhibited a positive and highly significant correlation with seed yield per plant (0.526) and had a positive direct effect on seed yield per plant (0.4869). Similarly, pod length showed a positive and significant correlation with seed yield per plant (0.252) and had a positive direct effect on seed yield per plant (0.1352). These results were in par with findings reported by Partap *et al.*, (2019), Sathees *et al.*, (2019), Kiran and Lal (2021) and Rana *et al.*, (2022) for pods per plant and pod length. 100 seed weight displayed a positive but statistically non-significant

correlation with seed yield per plant (0.094) and exhibited a positive direct effect on seed yield per plant (0.0597). In contrast, days to maturity showed a significant negative correlation with seed yield per plant (-0.164) and had a negative direct effect on seed yield per plant (-0.1844). Harvest index demonstrated a positive and significant correlation with seed yield per plant (0.208) and had a positive direct effect on seed yield per plant (0.0177). This aligns with similar positive and direct effects of harvest index (%) on seed yield per plant reported by Kumar *et al.*, (2014), Partap *et al.*, (2019), Kiran and Lal (2021) and Rana *et al.*, (2022). Among the traits analyzed, branches per plant exhibited the highest negative direct effects on seed yield per plant, followed by days to maturity and days to 50% flowering, indicating their relatively lower significance in selection for higher seed yield in black gram. Conversely, the traits with the highest positive direct effects on seed yield per plant were pods per plant, followed by clusters per plant and seeds per pod. Therefore, special attention should be given to these three characters during the selection process to enhance seed yield per plant in black gram. Path coefficient analysis for seed yield revealed a residual effect of 0.57, suggesting that future studies should consider incorporating additional traits to capture a broader spectrum of available variation.

### **Conclusion**

In summary, the findings of this study indicate that there is a significant amount of variability within the studied black gram progenies. Traits with high heritability and high genetic advance, suggesting the significant role of additive gene effects and show effective selection and improvement. Seed yield per plant (g) is positively correlated with yield-related components such as plant height (cm), pod length (cm), seeds per pod, clusters per plant, pods per plant, and harvest index (%). This indicates that these traits can be simultaneously improved with seed yield per plant through direct selection. Path analysis highlights that the most influential direct effects on seed yield per plant are exhibited by pods per plant, followed by clusters per plant and seeds per pod. In conclusion, the results of this study provide valuable insights into the genetic variability, heritability, and interrelationships among various traits in black gram  $F_4$  progenies. These findings can guide breeding programs aimed at enhancing the yield and agronomic characteristics of black gram varieties through effective selection strategies.

### **Disclaimer (Artificial intelligence)**

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. ChatGPT 40 mini
- 2.
- 3.

## References

- Allard RW. *Principles of plant breeding*. John Wiley and Sons Inc., New York. pp: 1960;145-147.
- Anonymous. Ministry of Agricultural and Farmers Welfare, Govt. of India, 2022. Retrieved from <https://eands.dacnet.nic.in/> Accessed 30 April, 2023.
- Bishnoi A, Gupta P, Lal GM. Evaluation of genetic variability and heritability in black gram [*Vigna mungo* (L.) Hepper] genotypes. *Journal of pharmacognosy and phytochemistry*.2017;6(4):493-496.
- Burton GW, De Vane DE. Estimating heritability in tall fescue (*Festucaarundinacea*) from replicated clonal material. *Agronomy Journal*,1952;45(10):478-481.
- Dewey OR, Lu KH. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*,1959;57:515-518.
- Gowsalya P, Kumaresan D, Packiaraj D, Bapu JR. Genetic variability and character association for biometrical traits in black gram [*Vigna mungo* (L.) Hepper]. *Electronic Journal of Plant Breeding*,2016;7(2):317-324.
- Johnson HW, Robinson HF, Comstock RE. Genotypic and phenotypic correlation in soybean and their implications in selection. *Agronomy Journal*,1955;47:477-483.
- Kiran MS, Lal G. Studies on the component approach of selection in black gram [*Vigna mungo* (L.) Hepper]. *The bioscan*, 2021;16(1):61-67.
- Kondagari H, Sapna SL, Lal GM. Study on genetic variability and correlation in black gram [*Vigna mungo* (L.) Hepper]. *Journal of Pharmacognosy and Phytochemistry*, 2017;6(4):674-676.
- Kumar BS, Padmavathi S, Prakash M, Genesan J. Correlation and path analysis in blackgram (*Vigna mungo* (L.) Hepper). *Legume Research*, 2003;26(1):75-76.
- Kumar GV, Vanaja M, Sathish P,Vagheera P,Shishodia SS,Razak A. Correlation and path analysis of seed yield and yield contributing components of black gram [*Vigna mungo* (L.) Hepper] under rainfed condition from Andhra Pradesh, India. *International journal of applied biology and pharmaceutical technology*,2014;5(3):137-140.
- Nunes MC, Raymundo A, Sousa I. Gelled vegetable desserts containing pea protein, κ-carrageenan and starch. *European Food Research and Technology*, 2006;222(5):622-628.
- Panda DP, Lenka D, Dash AP, Tripathy SK. Genetic variability and heritability studies in relation to seed yield and its component traits in black gram [*Vigna mungo* (L.) Hepper]. *Trends in Biosciences*, 2017;10:1411-1414.
- Panigrahi KK, Mohanty A, Baisakh B. Genetic divergence, variability and character association in landraces of black gram [*Vigna mungo* (L.) Hepper] from Odisha. *Journal of crop and weed*,2014;10(2):155-165.
- Partap B, Kumar M, Kumar V, Kumar A. Genetic variability and correlation studies of seed yield and its components in black gram [*Vigna mungo* (L.) Hepper]. *Journal of Pharmacognosy and Phytochemistry*, 2019;8(3):2035-2040.
- Patel RN, Madhu Bala. Correlation studies for yield and its components in black gram [*Vigna mungo* (L.) Hepper]. *International Journal of Current Microbiology and Applied Sciences*,2020a;9(09):1441-1447.
- Patel RN, Madhu Bala. Genetic variability study for yield and its components in black gram [*Vigna mungo* (L.) Hepper]. *Journal of Pharmacognosy and Phytochemistry*,2020b;9(4):2061-2064.
- Rana VB, Makati JP, Chaudhary N, Joshi RJ, Vanapariya S, Shah ZS, Malaviya AV, Patel S. Evaluation of genetic variability parameters in 185 F<sub>3</sub> progenies of black gram [*Vigna*

- mungo* (L.) Hepper] for Seed Yield and Related Components. Biological Forum-An International Journal,2022;14(4):1056- 1060.
- Rao CM, Rao YK, Reddy M. Evaluation of mung bean germplasm for yield and yield components. Legume Research-An International Journal, 2006;29(1):73-75.
- Sathees N, Shoba D, Saravanan S, Merina PKS, Pillai MAR. Studies on genetic variability, association and path coefficient analysis in black gram [*Vigna mungo* (L.) Hepper]. International Journal of Current Microbiology and Applied Sciences, 2019;8(6):1892-1899.
- Souframanien J,Gopalakrishna T. A comparative analysis of genetic diversity in black gram genotypes using RAPD and ISSR markers. Theoretical and Applied Genetics, 2004; 109(8):1687-93.
- Sushmitharaj DV, Shoba D, Pillai MA. Genetic variability and correlation studies in black gram [*Vigna mungo* (L.) Hepper] with reference to YMV resistance.International Journal of Current Microbiology and Applied Sciences, 2018;6:2849-2856.
- Swamy AA, Reddy G. Genetic divergence and heterosis studies in mung bean (*Vigna radiata* (L.) Wilczek). Legume Research-An International Journal,2004; 27(2):115-18.
- Vyas D, Joshi A, Rajamani G, Jain D. Assessment of genetic diversity in black gram (*Vigna mungo* 23456(L.) Hepper) genotypes based on ISSR. Legume Research-An International Journal, 2018;41(2):175-81.

**Table 1: Measures of variability parameters for different characters in F<sub>4</sub> progenies of black gram**

Character	Range		$\sigma_g^2$	$\sigma_p^2$	GCV (%)	PCV (%)	$h_{bs}^2$ (%)	GA	GAM as % of mean
	Min.	Max.							
Days to 50% flowering	58.00	66.00	5.87	6.32	3.94	4.09	92.88	4.82	7.83
Plant height (cm)	32.10	58.01	20.10	48.46	9.48	14.72	41.48	5.96	12.59
Branches per plant	1.30	9.20	3.79	3.87	32.29	32.64	97.83	3.97	65.88
Clusters per plant	6.80	29.00	35.17	37.51	36.08	37.26	93.75	11.85	72.07
Pods per plant	13.80	88.40	460.8	470.1	37.79	38.17	98.03	43.85	77.20
Pod length (cm)	3.56	5.80	0.19	0.21	10.31	10.83	90.62	0.85	20.20
Seeds per pod	4.40	8.00	0.23	0.36	7.97	9.86	65.32	0.81	13.28
100 Seed weight (g)	1.80	6.50	0.75	0.82	19.33	20.25	91.07	1.70	38.05
Days to maturity	87.00	123.0	101.6	102.3	9.97	10.01	99.29	20.73	20.50
Harvest index (%)	19.84	63.96	60.86	99.20	15.93	20.34	61.35	12.61	25.74
Seed yield per plant (g)	2.66	19.30	22.13	28.03	34.70	39.05	78.95	8.62	63.60

$\sigma_g^2$  = Genotypic variance

$\sigma_p^2$  = Phenotypic variance

GCV = Genotypic coefficient of variation

PCV = Phenotypic coefficient of variation

$h_{bs}^2$  (%) = Heritability (Broad sense)

GA = Genetic advance

GAM = Genetic advance as per cent of mean (%)

**Table 2: Phenotypic correlation coefficients of seed yield per plant with other characters in F<sub>4</sub> progenies of black gram**

Traits	DTF	PH	BPP	CPP	PPP	PL	SPP	SW	DTM	HI	SY
<b>DTF</b>	<b>1.000</b>										
<b>PH</b>	-0.012 <sup>NS</sup>	<b>1.000</b>									
<b>BPP</b>	0.035 <sup>NS</sup>	0.841 <sup>**</sup>	<b>1.000</b>								
<b>CPP</b>	0.104 <sup>NS</sup>	0.790 <sup>**</sup>	0.771 <sup>**</sup>	<b>1.000</b>							
<b>PPP</b>	0.004 <sup>NS</sup>	0.515 <sup>**</sup>	0.448 <sup>**</sup>	0.450 <sup>**</sup>	<b>1.000</b>						
<b>PL</b>	0.082 <sup>NS</sup>	0.028 <sup>NS</sup>	-0.033 <sup>NS</sup>	-0.011 <sup>NS</sup>	0.166 <sup>NS</sup>	<b>1.000</b>					
<b>SPP</b>	0.032 <sup>NS</sup>	-0.134 <sup>NS</sup>	-0.160 <sup>NS</sup>	-0.137 <sup>NS</sup>	-0.056 <sup>NS</sup>	0.212 <sup>*</sup>	<b>1.000</b>				
<b>SW</b>	0.107 <sup>NS</sup>	-0.099 <sup>NS</sup>	-0.093 <sup>NS</sup>	-0.079 <sup>NS</sup>	-0.022 <sup>NS</sup>	0.120 <sup>NS</sup>	0.060 <sup>NS</sup>	<b>1.000</b>			
<b>DTM</b>	-0.098 <sup>NS</sup>	-0.081 <sup>NS</sup>	-0.096 <sup>NS</sup>	-0.051 <sup>NS</sup>	-0.005 <sup>NS</sup>	0.079 <sup>NS</sup>	0.008 <sup>NS</sup>	-0.175 <sup>NS</sup>	<b>1.000</b>		
<b>HI</b>	0.077 <sup>NS</sup>	0.746 <sup>**</sup>	0.857 <sup>**</sup>	0.749 <sup>**</sup>	0.436 <sup>**</sup>	0.008 <sup>NS</sup>	-0.054 <sup>NS</sup>	-0.099 <sup>NS</sup>	-0.017 <sup>NS</sup>	<b>1.000</b>	
<b>SY</b>	-0.078 <sup>NS</sup>	0.281 <sup>**</sup>	0.179 <sup>NS</sup>	0.273 <sup>**</sup>	0.526 <sup>**</sup>	0.252 <sup>*</sup>	0.220 <sup>*</sup>	0.094 <sup>NS</sup>	-0.164 <sup>NS</sup>	0.208 <sup>*</sup>	<b>1.000</b>

\*\* Significant at 1% level of probability, \* Significant at 5.0 % level of probability

DTF = Days to 50% flowering

PH = Plant height (cm)

BPP = Branches per plant

CPP = Clusters per plant

PPP = Pods per plant

PL = Pod length (cm)

SPP = Seeds per pod

SW = 100 Seed weight (g)

DTM = Days to maturity

HI = Harvest index

SY = Seed yield per plant (g)

**Table 3: Phenotypic path coefficient analysis of component characters towards seed yield per plant in F<sub>4</sub> progenies of black gram**

Traits	DTF	PH	BPP	CPP	PPP	PL	SPP	SW	DTM	HI	Correlation with seed yield per plant
<b>DTF</b>	<b>-0.1376</b>	-0.0007	-0.0087	0.0234	0.0017	0.0111	0.0070	0.0064	0.0180	0.0013	<b>-0.078<sup>NS</sup></b>
<b>PH</b>	0.0015	<b>0.0655</b>	-0.2099	0.1763	0.2505	0.0037	-0.0295	-0.0059	0.0149	0.0132	<b>0.281<sup>**</sup></b>
<b>BPP</b>	-0.0048	0.0551	<b>-0.2496</b>	0.1721	0.2183	-0.0044	-0.0353	-0.0055	0.0177	0.0152	<b>0.179<sup>NS</sup></b>
<b>CPP</b>	-0.0143	0.0518	-0.1925	<b>0.2231</b>	0.2191	-0.0015	-0.0302	-0.0046	0.0094	0.0133	<b>0.273<sup>**</sup></b>
<b>PPP</b>	-0.0004	0.0337	-0.1119	0.1004	<b>0.4869</b>	0.0224	-0.0124	-0.0013	0.0008	0.0077	<b>0.526<sup>**</sup></b>
<b>PL</b>	-0.0113	0.0018	0.0081	-0.0025	0.0807	<b>0.1352</b>	0.0468	0.0071	-0.0146	0.0001	<b>0.252<sup>*</sup></b>
<b>SPP</b>	-0.0043	-0.0087	0.0398	-0.0305	-0.0274	0.0286	<b>0.2211</b>	0.0035	-0.0014	-0.0009	<b>0.220<sup>*</sup></b>
<b>SW</b>	-0.0148	-0.0064	0.0232	-0.0175	-0.0105	0.0161	0.0131	<b>0.0597</b>	0.0323	-0.0017	<b>0.094<sup>NS</sup></b>
<b>DTM</b>	0.0134	-0.0053	0.0240	-0.0114	-0.0022	0.0107	0.0017	-0.0104	<b>-0.1844</b>	-0.0003	<b>-0.164<sup>NS</sup></b>
<b>HI</b>	-0.0105	0.0489	-0.2139	0.1670	0.2124	0.0010	-0.0119	-0.0059	0.0031	<b>0.0177</b>	<b>0.208<sup>*</sup></b>

\*\* Significant at 1.0 per cent level of probability, \* Significant at 5.0 per cent level of probability, Residual =0.57, Bold diagonal figures are the direct effects

DTF = Days to 50% flowering

PH = Plant height (cm)

BPP = Branches per plant

CPP = Clusters per plant

PPP = Pods per plant

PL = Pod length (cm)

SPP = Seeds per pod

SW = 100 Seed weight (g)

DTM = Days to maturity

HI = Harvest index (%)




Fig 1: GCV (%) and PCV (%) for eleven quantitative characters in F<sub>4</sub> progenies of black gram

GCV (%) = Genotypic coefficient of variation

PCV (%) = Phenotypic coefficient of variation





Fig 2: GAM and h<sup>2</sup><sub>(bs)</sub> for eleven quantitative characters in F<sub>4</sub> progenies of black gram

GAM = Genetic advance as per cent of mean (%)

h<sup>2</sup><sub>(bs)</sub> (%) = Heritability (Broad sense) (%)



**Fig 3. Phenotypic path diagram for seed yield per plant (g) in F<sub>4</sub> progenies of black gram**

DTF = Days to 50% flowering  
 PH = Plant height (cm)  
 PL = Pod length (cm)  
 BPP = Branches per plant  
 SPP = Seeds per pod  
 SW = 100 Seed weight (g)  
 DTM = Days to maturity  
 HI = Harvest index (%)  
 CPP = Clusters per plant  
 PPP = Pods per plant

UNDER PEER REVIEW