

EVIDENCE AND IMPACTS OF CLIMATE CHANGE ON SMALLHOLDER PADDY FARMERS IN PEMBA ISLAND

Abstract

This study presents evidence of climate change and its impacts on paddy farmers in Pemba Island. The phenomenon of ocean warming and accelerated ice mass loss has led to a significant rise in global sea levels, averaging 4.62 mm per year between 2013 and 2022, reaching an unprecedented peak in 2022. As a result, Pemba Island is experiencing escalating challenges due to rising sea levels, shifting precipitation patterns, and increasing temperatures. The primary objective of this study is to examine the evidence of climate change in the study area and assess the specific ways in which paddy farmers are affected by these changes on Pemba Island. Conducted in the Wete District, this research utilizes a cross-sectional design, involving 71 respondents selected through a combination of multistage cluster, purposive, and random sampling techniques. Data collection was carried out through surveys, focus groups, and interviews, and subsequently analyzed using SPSS and content analysis methodologies. The study also used climate data from Tanzania Meteorological Authority (temperature and rainfall data) over the past 31 years (1992-2023). The findings indicate that paddy farmers are grappling with higher temperatures, altered rainfall patterns, prolonged dry spells, intrusion of seawater, and an increase in pest infestations. As a result, their agricultural production is significantly reduced, exacerbating food insecurity and negatively impacting their livelihoods. In light of these findings, it is evident that robust adaptation measures must be implemented to address the challenges faced by paddy farmers. The study recommends the construction of sea walls as a means of mitigating the adverse effects of rising sea levels. Furthermore, developing drought-resistant paddy seeds varieties is proposed as a strategy to improve productivity and enhance resilience in the face of changing climatic conditions.

Keywords: evidence; impacts; climate change; paddy farmers; Wete district; Pemba Island

1.0 Introduction

Climate change is an urgent global challenge that has significant implications for human social, economic, and environmental well-being (Amjath-Babu *et al.*, 2016). The Earth's average surface temperature has been rising, with an increasing rate of ocean warming over the past two decades. In fact, the ocean heat content in 2022 reached the highest recorded levels (IPCC, 2007a; WMO, 2023). This warming, along with the accelerated loss of ice mass from ice sheets, has contributed to a rise in global mean sea level. Between 2013 and 2022, the global mean sea level increased by 4.62 mm per year, reaching a new record high in 2022 (WMO, 2023). Projections indicate that the rise in global sea level is likely to continue throughout the 21st century, with estimates ranging from 18-59 cm above 1990 levels by the end of the century (Meehl *et al.*, 2007).

The consequences of these changes are already becoming evident as climates and ecosystems undergo significant shifts (IPCC, 2007b). Island communities, such as Zanzibar, are particularly vulnerable to the impacts of climate change. Rising sea levels, changing precipitation patterns, and increasing air and sea surface temperatures are placing additional

stress on already limited island resources. At the same time, climate change policies often constrain local decision-making (Lazrus, 2012).

Zanzibar, due to its geographic position and limited land area as islands, is highly exposed to climate change and its associated impacts. The region is already experiencing sea level changes, and saltwater intrusion into underground aquifers, all of which threaten the availability and quality of water resources. Consequently, water stress has increased, and crop productivity has been affected (IPC, 2017). Furthermore, the projected rise in sea levels will exacerbate the problem by leading to coastal inundation and erosion, further reducing the already scarce land area. The degradation of coastal resources like mangroves and coral reefs, which support important sectors like fisheries and tourism, is also expected (RGoZ, 2010; Mwangora *et al.*, 2015).

Zanzibar faces additional challenges in adapting to climate change due to its limited economic capacity to respond effectively. The high cost of implementing adaptation measures further compounds vulnerability. Climate change not only poses a threat to achieving sustainable development and poverty reduction but also has the potential to undermine the progress made towards the Millennium Development Goals (MDGs) and Sustainable Development Goals (SDGs) through Zanzibar's Strategy for Growth and Reduction of Poverty (MKUZA) (RGoZ, 2010; Mwangora *et al.*, 2015). Small island developing states (SIDSs), including Pemba Island, are expected to be particularly affected by increasing temperatures, changes in precipitation patterns, and rising sea levels (Lazrus 2012). Their vulnerability stems from their sensitive ecological and economic systems, as well as other interacting stressors (Mangora *et al.*, 2015; Bakari, 2015; Makame, 2013). Changes in the timing and amount of rainfall have an impact on freshwater resources, which are crucial for island living and activities such as agriculture (Lazrus 2012). As a result, this can have potential adverse effects on island economies and food security.

Paddy farming was introduced to Pemba Island from Asia in the second century AD as a response to a wet climate and population growth on the island. This agricultural innovation was adopted from Asia due to monsoon exchanges (Walshaw, 2010). The wet climate, combined with a growing population, necessitated the intensification of paddy production to meet the food demands of the island's inhabitants, ultimately leading to rice (paddy) becoming the staple food on the island (Busungu, 2023). Despite the increasing challenges posed by climate change and variability, the tradition of paddy farming during wet seasons has persisted to this day.

However, like any other agricultural crops in Pemba and other developing countries, paddy farming is being affected by the impacts of climate change and variability (RGoZ, 2012; IPC, 2017). Crop production in Pemba, including paddy farming, is heavily dependent on weather and climate conditions. The island is also influenced by regional patterns of extreme weather events and rising sea levels, which result in floods, droughts, and storms. Despite the existing research on climate change and adaptation in Pemba Island and Zanzibar as a whole, empirical evidence regarding how paddy farmers in Pemba Island are affected by climate change and variability is relatively limited. Therefore, there was a need to understand how these farmers were impacted in order to develop adaptation strategies for future climate change. This study aimed to address this knowledge gap by identifying indicators of climate change and its impacts on paddy farming production in Pemba Island.

This study is aligned with the Zanzibar Climate Change Strategy (RGoZ, 2014), which emphasizes the need for climate-smart agriculture, improved natural resource management, and the development of climate information, capacity, and disaster risk management. Additionally, this study is in line with Sustainable Development Goal 2, which relates to food security. Therefore, this study contributes to identifying the impacts of climate change on paddy production, ultimately jeopardizing food security in the changing climate of Pemba Island.

2.0 Methodology

The study was conducted in the Wete District, North Pemba Region, Pemba Island, Zanzibar, Tanzania. Pemba Island is one of the two main islands that form part of the Zanzibar Archipelago. The other main island is Unguja, and together with several small islands, they make up Zanzibar. Wete District is one of two administrative districts of Pemba North Region in Tanzania. The district covers an area of 295 km². The district has a water border to the east and west by the Indian Ocean and is bordered to the north by Micheweni District. The district is latitudes 5° 6' 57.6" S and longitudes 39° 45' 18" E. The main activities carried out in Wete District are crop farming, livestock keeping, fishing, and trade. Like the remaining part of Pemba Island, all of the activities are climate-sensitive. Pemba Island acquired the nickname "Green Island" from an Arab writer due to its history of supplying agricultural produce, including rice, grains, and cereals, to Malindi and Mombasa (Pollard, 2009).



Figure 1: Wete District – Pemba Island

Pemba Island has a tropical climate, which is milder than both mainland Tanzania and Unguja Island. The average temperature is 25.5°C (78°F), and the average annual rainfall is 1,364 mm. Monthly average temperatures range between 24 and 27.4°C (75 and 81°F). There are two rainy seasons, with the majority of rainfall occurring between April and May, and a smaller rainy season between November and December. The drier months are January and February, with a longer dry season from June to October (Kai *et al.*, 2020). The decision to select Pemba Island for this study is twofold. First, historically, Pemba Island has been a supplier of agricultural produce, including rice, grains, and cereals, to Malindi and Mombasa. However, due to rising sea levels, changes in precipitation patterns, increasing temperatures, and floods (WMO, 2023; URT, 2021), Pemba Island is more vulnerable to the impacts of climate change compared to other areas in Zanzibar. Secondly, like other smallholder farmers in Tanzania, paddy farmers in Pemba Island rely on rainfed agriculture. This practice makes them particularly vulnerable to the impacts of climate change.

A cross-sectional research design was utilized for this study, which allowed for the collection of multiple cases at a single point in time (Creswell and Creswell, 2018). The selection of districts, wards, “shehias”, and households was done through a multistage cluster sampling procedure. This approach enabled the use of multiple sampling methods. Wete District was purposively selected based on its geographical location and the prevalence of paddy farming.

Wards, shehias, and households were then randomly selected using simple random sampling procedure. The study used a sample of 71 respondents.

Both quantitative and qualitative data were collected for this study. Quantitative data was obtained through a household questionnaire survey, while qualitative data was gathered through focus group discussions and key informant interviews. The questionnaire was pre-tested and revised before being administered to the heads of household paddy farmers in Wete District, North Pemba Region. The aim was to gather information on the impacts of climate change on paddy farmers. Three focus group discussions were conducted, each involving seven to nine participants. Key informant interviews were also conducted with experienced paddy farmers, taking into account their years of experience working in the field being 15 years and above. Meteorological data of rainfall and temperature covering a period of 31 years between 1992 and 2023 were collected from Tanzania Meteorological Authority (TMA).

A descriptive analysis was conducted using the Statistical Package for Social Sciences (IBM SPSS) version 20 software to analyse the quantitative data. Qualitative data were analyzed basing on their content. Climatic data from Tanzania Meteorological Authority (TMA) were analyzed using Microsoft Office Excel.

3.0 Results and Discussion

3.1 Evidence of Climate Change in Pemba Island

3.1.1 Increase in temperature

The findings revealed that respondents were aware of the changing climate. More than 79% of the respondents agreed that the temperature was increasing, while 20% of respondents were neutral and did not know whether the temperature was increasing or not. However, one respondent disagreed that the temperature was increasing. Rising temperatures associated with climate change will likely have a detrimental impact on crop production, livestock, fisheries, and allied sectors (Kumar *et al.*, 2017). This finding was complemented by a key informant interview, in which one key informant in Gando Ward, Wete District, stated:

"There is a noticeable increase in temperature, causing plants to dry up, paddy fields to become yellowish, and the soil to become dry and cracked due to high temperatures."




Figure 2: Minimum and maximum temperature of 31 years in the study area from 1992-2023

The trend analysis results for the period 1992 – 2023 show a significant increasing trend in mean annual maximum and minimum temperature ($P= 0.0905$, $R^2=0.5122$; $P=0.0112$, $R^2=0.0379$) respectively as shown in Fig. 2. The increase in mean annual minimum temperature was found to occur much faster than for mean annual maximum temperature. These findings show strong agreement on continued future warming in the range of 0.8°C to 1.8°C by the 2040s (URT 2021). According to climate science, further warming is inevitable over the next decade and likely beyond (Woetzel *et al.*, 2020). With increasing global average temperatures, climate models indicate a rise in climate hazards worldwide. These models also suggest that further warming will continue to increase the frequency and/or severity of acute climate hazards and intensify chronic hazards (Woetzel *et al.*, 2020). The projected seasonal temperature increases by 2°C in Tanzania 2050 will reduce average rice yields by 7.6 percent (Mafie 2021).

Table 1: The temperature is increasing

Temperature	Frequency	Percent
Agree	56	79
Neutral	14	20
Disagree	1	1
Total	71	100

3.1.2 Changes in rainfall pattern

The findings indicate that 49% of respondents agreed that there have been noticeable changes in the rainfall pattern of Pemba Island. Meanwhile, 47% of respondents remained neutral on the matter, and 4% disagreed that any changes had occurred. However, the changes in rainfall are complex, and do not appear to be simple precipitation trends across the Zanzibar islands (RGoZ, 2012). During focus group discussions, it was revealed that the rainfall on the island is unreliable. It occurs unpredictably, sometimes resulting in insufficient rain for crop production, while at other times, heavy rainfall leads to flooding in farmlands and low-lying

areas. Regardless of the amount of rainfall, whether it is minimal or excessive, it has significant impacts on crop production. The rainfall meteorological data confirms farmers observations on the anomalies of rainfall intensity and distribution in the study area. For example, in 1997 the annual rainfall was 2194.2mm, 1998 was 2082.4, 2001 was 869.6mm, 2003 was 977.9mm, 2004 was 967.8mm, 2017 was 2222.0mm, 2018 was 2047.2mm, 2019 was 2312.8mm, 2021 was 957.7 and 2022 was 958.7mm, show higher intensity and minimal rainfall events. Figure 3,4,5,&6 shows indications of changes in rainfall pattern.




Figure 3: Annual rainfall of 31 years (1992 – 2023) in the study area




Figure 4: Monthly(October, November and December) rainfall of 31 years in the study area




Figure 5: Monthly(January and February) rainfall of 31 years in the study area




Figure 6: Monthly (March, April and May) rainfall of 31 years in the study area

Table 2:Changes in Rainfall Pattern

Rainfall pattern change	Frequency	Percent
Agree	35	49
Neutral	33	47
Disagree	3	4
Total	71	100

3.1.3 Rise of sea level

The rise of sea levels is one of the evidences revealed on climate change in Wete District, Pemba Island. Approximately 66 percent of respondents agreed that the sea level is rising, while 25 percent remained neutral, and nine percent disagreed with the idea of sea level increase. However, these results were supported by the findings of focus group discussions which also indicated that the sea level was indeed rising. State of the climate in Africa 2022 report by the World Meteorological Organization (WHO, 2023) stated that the highest rate of 3.7mm/year sea level rise around Africa has been observed along the coastal areas of the Red Sea followed by the coastal areas of the western Indian Ocean (including Wete in Pemba Island) where the rate rarely exceeds 3.6mm/year. This increase in seawater has led to the intrusion of saltwater into paddy farms, resulting in crop destruction. As stated by Mwangora, *et al* (2015), the intrusion of saltwater into natural wells and agricultural lands has resulted in increased water stress and reduced crop productivity.

Table 3: Rise of Sea Level

There is rise of sea level	Frequency	Percent
Agree	47	66
Neutral	18	25
Disagree	6	9
Total	71	100

3.1.4 Increased dry spells

The findings revealed an increase in dry spells during the rainy season. 51 percent of respondents agreed that there were prolonged dry spells during the rainy season, 34 percent of respondents were neutral, and 15 percent of respondents did not agree that there was an increase in dry spells. These findings were complemented by a focus group discussion, which revealed that there was an increase in dry spells during the rainy season to the extent that the soil cracks. Increased dry spells adversely affect crop production by limiting the water supply to the crops, inhibiting plant growth, and reducing overall production (Woetzel, 2020). Consequently, this could have adverse impacts on the livelihoods of farmers.

Table 4: Increase in dry spells

There is Increase in dry spells	Frequency	Percent
Agree	36	51
Neutral	24	34
Disagree	11	15
Total	71	100

3.2 Impacts of climate change in paddy production

3.2.1 Intrusion of sea water into Paddy farms

Climate change has had a significant impact on paddy production in Wete District, North Pemba Region. One of the consequences of climate change is the intrusion of seawater into paddy farms. The research findings indicate that 75 percent of the respondents acknowledged that their paddy production was negatively affected by seawater intrusion. In contrast, 25 percent of the respondents claimed that their paddy farms were unaffected by sea water. The focus group discussions further supported these findings, revealing that when seawater enters the paddy farms, all crops in the affected areas perish. This intrusion of seawater not only jeopardizes food security but also affects the income of paddy farmers who rely on paddy production as their primary source of sustenance. The affected farms remain unproductive until rainfall washes away the salt brought by the sea water. According to the World Meteorological Organization (WMO) (2023), the global mean sea level has risen by 4.62 mm per year from 2013 to 2022, reaching a new record high in 2022. Considering this ongoing rise in sea levels, paddy production will continue to be impacted by seawater unless effective measures are taken to prevent its intrusion into paddy farms.

Table 5: Intrusion of Sea Water in Paddy Farms

Sea water enters paddy fields	Frequency	Percent
Yes	53	75
No	18	25
Total	71	100

3.2.2 Increased pests affecting paddy production

Climate change has also had an impact on paddy production by leading to a rise in pests in paddy farms. The economies of climate change in Zanzibar technical report, (2012) of the Revolutionary Government of Zanzibar, stated that climate change would lead to changes in pests and diseases. Approximately 61 percent of the respondents acknowledged an increase in pests that adversely affected paddy production in Wete. Conversely, 39 percent did not report any noticeable rise in pests affecting paddy production (see Table 6). These findings were further supported by the focus group discussions conducted in the study area. The discussions revealed that the increase in pests was primarily due to higher temperatures. This surge in pests not only affects paddy production but also increases the costs of production due to the need for purchasing pesticides. Consequently, the presence of pests in paddy farms results in reduced paddy production, causing farmers to experience food insecurity and a decline in income.

Table 6: Increase of pests affecting paddy production

Increase of pests	Frequency	Percent
Yes	43	61

No	28	39
Total	71	100

3.3 Impacts of climate change on food security among paddy farmers in the study area

Climate change has a direct and significant impact on food insecurity. The changing climate affects all four dimensions of food security: availability, accessibility, utilization, and stability of food systems (Rao, *et al.*, 2017; FAO, 2018). Many countries are facing the negative impacts of climate change, particularly in terms of decreased agricultural productivity, which leads to lower levels of national and household food security (Nyang'au, 2021). As global temperatures rise due to climate change, the production of food becomes more difficult and uncertain due to changes in weather patterns, extreme weather events, and other environmental disruptions (Rao *et al.*, 2017; WEF, 2023). Various studies show that paddy production is affected negatively by temperature and precipitation changes, that lead to decline in its yield (Davkota and Pajja, 2020; Liu *et al.*, 2020; Herath *et al.*, 2020; Sandet *al.*, 2022; Chandrasiri *et al.*, 2023; Zella *et al.*, 2024). In the study area, climate change played a major role in food insecurity among paddy farmers. The findings showed that 94 percent of respondents experienced food insecurity, while six percent did not (Table 7).

Table 7: Impacts of climate change on food security

Food insecurity	Frequency	Percent
Yes	67	94
No	4	6
Total	71	100

Paddy farmers in Wete District, Pemba Island have been experiencing various challenges due to climate change. These include increased temperatures, changes in rainfall patterns, longer dry spells during the rainy season, intrusion of seawater into paddy farms, and increased pest infestations that negatively affect paddy production (refer to table 1, 2, 4, 5, and 6). These changes have had a significant impact on the farmers' ability to produce an adequate amount of paddy crops, thus affecting food availability component, especially for poor household farmers. Low household income that led to low purchasing power affect access to food in the study area. Generally, availability, access, and stability of food security components, have all been compromised, as farmers' incomes also been affected. These findings were further supported by three informant interviews conducted in the study area:

The first informant stated:

"The increasing temperature has a detrimental effect on paddy production. The soil becomes dry and cracks, leading to the death of paddy plants. As the head of the household, this had a

significant impact on my ability to provide for my family. Paddy farming is our primary source of food and income, so the decrease in production has caused great hardship."

The second informant shared:

"We have been harvesting fewer paddy crops due to adverse weather conditions such as higher temperatures, increased pest attacks, and the intrusion of sea water into our farms. As the head of the household, I am highly affected by this situation. I struggle to provide enough food for my children, as well as clothing and healthcare in case of illness. Drought and famine have become significant challenges for us."

The third informant explained:

"As women, we bear the brunt of these challenges. Sometimes, our husbands are forced to seek casual labor elsewhere due to the low paddy harvest. In their absence, we are left to shoulder the entire responsibility of taking care of our families, including providing food, clothing, and healthcare for our children when they fall ill. This situation adds an immense burden to us women."

4.0 Conclusion

The findings indicate that paddy farmers in Wete District, Pemba Island are facing several challenges related to climate change. These include increased temperatures, changes in rainfall patterns, longer dry spells during the rainy season, intrusion of sea water into paddy farms, and increased pest infestations that negatively impact paddy production. These climate-related changes have led to a reduction in paddy production, which in turn has affected food security and the livelihoods of paddy farmers in Wete District, Pemba Island. If no strong adaptation measures are implemented to enhance the resilience of paddy farmers against the impacts of climate change, paddy production will continue to be adversely affected. The study recommends that the Ministry of Agriculture of the Revolutionary Government of Zanzibar and other stakeholders take action to protect paddy farmers from further impacts of climate change. This can be achieved by implementing adaptation strategies that build the resilience of paddy farmers. For example, constructing a concrete wall along the Wete District beach can help reduce or eliminate the intrusion of sea water into paddy farms. Additionally, the development of drought-resistant paddy seed varieties will enable farmers to increase paddy production regardless of the prevailing weather conditions in the study area. By prioritizing the resilience of paddy farmers against the impacts of climate change, there will be improvements in paddy productivity, self-sufficiency in food security, and the overall livelihoods of paddy farmers.

Disclaimer (Artificial intelligence)

Option 1:

Mary Kihupi Author hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

8.0 References

- Amjath-Babu, T.S., Krupnik, T.J., Aravindakshan, S., Arshad, M., Kaechele, H., (2016). Climate Change and Indicators of Probable Shifts in the Consumption Portfolio of Dryland Farmers in Sub-Saharan Africa: Implications for Policy, *Ecol. Indicat.* 67, 830-838
- Bakari, S.M. (2015). Climate Change Impacts and Adaptation Strategies of Small-Scale Agriculture in Micheweni District, Pemba, Tanzania. MA thesis, Sokoine University of Agriculture.
- Busungu, C. (2023) Past, Present and Future Perspectives of Rice Production in Tanzania. *Agricultural Sciences*, 2023, 14, 987-1006 <https://www.scirp.org/journal/as> ISSN Online: 2156-8561 ISSN Print: 2156-8553
- Chandrasiri, C. K., Tsusaka, T.W., Ho, T. D. N., Zulfiqar, F., and Datta, A. (2023). Impacts of climate change on paddy yields in different c zones of Sri Lanka: a panel data approach. *Asia-PacJRegSci* 7455-487. <https://doi.org/10.1007/s41685-022-00264-5>
- Creswell, J. W., & Creswell, J. D. (2018). *Research design* (5th ed.). SAGE Publications
- Devkota, N., and Pajja, N. (2020). Impacts of climate change on paddy production: Evidence from Nepal. *Asian Journal of Agriculture and Development* 17(2) 63 - 77
- FAO (2018). *The Future of Food and Agriculture: Alternative Pathways to 2050*. Food and Agriculture Organization of the United Nations, Rome, Italy 228pp.
- Herath, g., Hasanov, A., and Park, J. (2020). Impacts of climate change on paddy production in Malaysia: empirical analysis at the national and state level experience. In J. Xu, S. E. Ahamed, F. L. Cooke, and G. Duca (Eds), *Proceedings of the thirteenth International Conference on Management Science and Engineering Mangement* (1st ed., 1 656-664. *Springer*. https://doi.org/10.1007/978-3-030-21248-3_48
- IPC (2017). *Zanzibar Food Security and Nutrition Analysis Report*. Zanzibar Food Security and Nutrition Monitoring – IPC Technical Working Group 25pp.
- IPCC (2007a). *Summary for Policy Makers*. In: *Climate Change 2007: The physical science basis*. Contribution of working group I to the fourth assessment report International

- Panel on Climate Change. (edited by Solomon, S., Qin, D., Manning, M., Chen, Z. Marquis, M., Averyt, K.B., Tignor, M. and Miller, H. L.). Cambridge University Press, Cambridge, UK and New York, USA. 1-18.
- IPCC (2007b). Summary for Policy Makers. In: Climate change 2007: impacts, adaptation and vulnerability. Contribution of working group II to the fourth assessment report International Panel on Climate Change. (Edited by Parry, M. I, Canziani, O. F., Palutikof, G. P van der Linden, P.J., and Hanson, C.E.), Cambridge University Press, Cambridge, UK. 7-22.
- Kai, K.H., Kijazi, A.L. and Osima, S.E. (2020) An Assessment of the Seasonal Rainfall and 16 Its Societal Implications in Zanzibar Islands during the Season of October to December, 2019. *Atmospheric and Climate Sciences*, 10, 509-529. <https://doi.org/10.4236/acs.2020.104026>
- Lazrus, H. (2012). Sea change: Island Communities and Climate Change. *Annual Review of Anthropology*. DOI: 10.1146/annurev-anthro-092611-145730
- Liu, Y., Tang, L., Qiu, X., Liu, B., Chang, X., Liu, L., Zhang, X., Cao, W., and Zhu, Y. (2020). Impacts of 1.5 and 2.0°C global warming on rice production across China. *Agricultural and Forest Meteorology*. 284 ISSN 0168 – 1923 <https://doi.org/10.1016/J.agrformet.2020.107900>
- Mafie, G. K. (2021). The Impact of Climate Change on Agricultural Productivity in Tanzania. *International Economic Journal*, 36(1), 129–145. <https://doi.org/10.1080/10168737.2021.2010229>
- Meehl, G. A., Stocker, T. F, Collins W, Friedlingstein W, Gaye A.T., (2007). Global climate projections. In *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, ed. S Solomon, D Qin, M Manning, Z Chen, et al., pp. 747–846. Cambridge, UK: Camb. Univ. Press
- Makame, M. O. (2013). Vulnerability and Adaptation of Zanzibar East Coast Communities to Climate Variability and Change and Other Interacting Stressors Doctor of Philosophy dissertation, Rhodes University.
- Mangora, M., Shalli, M. S., Kaur, N., Jumah S.M., Mwita F. & Bakar S. (2015). Local Adaptation Investment Planning in Zanzibar. A Baseline Review of Policy and Institutional Framework. Retrieved from <https://www.researchgate.net/publication/27628835>.
- Nyang'au, J. O., Mohamed, J. H., Mango, N., Makate, C., and Wangea, A. N. (2021). Smallholder Farmers' Perception of Climate Change and Adoption of Climate Smart Agriculture Practices in Masaba South Sub-County, Kisii, Kenya. *Heligon* e06789
- Pollard, E. (2009). Settlement Adaptation to a Changing Coastline: Archaeological Evidence from Tanzania, During the First and Second Millennia AD. *Journal of Island and Coastal Archaeology*, 4(1): 82–107). Doi: 10.1080/15564890902779677.
- Rao, R. K., Sharma, P.K., and Raghuraman, M. (2017). Impact of Climate Change on Food Security. *Indian Journal of Agriculture and Allied Sciences* 3 (2) ISSN 2395-1109

- RGoZ (2010). Zanzibar Strategy for Growth and Reduction Poverty 2010-2015 (MKUZA II). Zanzibar: Revolutionary Government of Zanzibar.
- RGoZ (2012). The Economies of Climate Change in Zanzibar. Vulnerability, Impacts and Adaptation. Technical Report. Revolutionary Government of Zanzibar. 102pp.
- RGoZ (2014) Revolutionary Government of Zanzibar, Zanzibar Climate Change Strategy, May 2014.
- Sand, S., Wang, D., Fahad, S., Alharby, H. F., Bamagoos, A. A., Mjrashi, A., Alabdallah, N. M., ALZahrani, S.S., AbdELgaward, H., Adnan, M., Sayyed, R. Z., Ali., S., and Hassan, S. (2022). Comprehensive impacts of climate change on rice production and adaptive strategies in China. *Front microbiol.* Doi:10.3389/fmicb.2022.926059
- URT (2021). Nationally Determined Contribution. The United Republic of Tanzania Vice President's Office. 35pp
- WEF (2023). Food Security. How to Mitigate the Effects of Climate Change on Global Food Security. <https://www.weforum.org/agenda/2023/mitigate-climate-change-food-security>
- WMO (2023). State of the Climate in Africa 2022. *World Meteorological Organization*. Geneva, Switzerland. 34pp.
- Woetzel, J., Pinner, D., Samandari, H., Engel, H., Krishnan, M., McCullough, R., Melzer, T. and Boettiger, S. (2020). Climate Risks and Response Physical Hazards and Socioeconomic Impacts: How Will African Farmers Adjust to Changing Patterns of Precipitation. Case Study. McKinsey Global Institute. 20pp
- Walshaw, S. C. (2010). Converting to Rice: Urbanization, Islamization and Crops on Pemba Island, Tanzania, Ad 700–1500). *World Archaeology*, 42(1): 137–154). Doi:10.1080/00438240903430399.
- Zella, A. Y., Kitali, L. J., Lunyelele, S. S., Asenga, S. S., and Nturo, E. M. (2024) Effects of climate change among paddy farmers in Kahama District, Tanzania. *International Journal of Geography, Geology and Environment* 5(2)122-132 Doi: <https://doi.org/10.22271/27067483.2023.v5.i2b.180>