

Review Article

Black Soldier Fly *Hermetia illucens* (Diptera: Stratiomyidae) For Achieving Sustainable Development Through Circular Economy

ABSTRACT

A circular economy is a strategy aimed at optimizing resources so that sustainability can be achieved. The world needs to shift from linear economy to circular economy. Black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae) is gaining popularity in recent years for its tremendous capacity for organic waste management and bioconversion of organic waste into nutrient rich compost. Larvae and prepupae of black soldier fly have been used as feed for poultry birds, fishes, other animals and it can be used as nutrient rich food for humans. It is a promising source of protein and can be utilized for chitin, protein and grease extraction and biodiesel production. Due to its potential applicability at various industries and thereby recycling the organic waste and close the loop in the production system, it has gained attention in recent years for achieving some of the sustainable development goals adopted by united nations by transition from linear to circular economy. India has vast potential for BSF-based industries.

Keywords: Black soldier fly, *Hermetia illucens*, waste management; animal feed, biodiesel, sustainable development; circular economy.

1. INTRODUCTION

In 2015, the United Nations general assembly adopted the 17 sustainable development goals (SDGs), aiming to transition the world to a sustainable and resilient trajectory by 2030. [1]. By 2050, the global population is projected to reach 9.7 billion [2], presenting a significant challenge to food security. To meet the nutritional needs of this growing population, food production will need to increase by 70%, necessitating sustainable agricultural practices and innovative solutions to ensure global food security [3]. To address the pressing global issues of hunger, poverty, nutrition, and food security, a significant increase in food productivity is crucial. by 2050, the demand for cereals is expected to rise by approximately 50% for both food and feed purposes. Moreover, the demand for other essential food products like meat, dairy, fish, vegetable oils, and more is projected to grow at an even faster rate. [4, 5, 6]. To achieve a more sustainable food system, a gradual transition from a linear economy (characterized by a "make-take-waste" approach) to a circular economy (based on "reduce-reuse-recycle and recover" principles) is essential. Over the past decade, the concept of a circular economy has gained global significance, emerging as a vital theme worldwide. By embracing circular economy principles, we can foster sustainable and resource-efficient policies, leading to long-term benefits for both the environment and society, [8]. "Further, the term circular bioeconomy was introduced by the European commission, which defines it as: "The production of renewable biological resources and the conversion of these resources and waste streams into value-added products such as food, feed, bio-based products, and bioenergy. Sustainability and circularity must be at the center of the bioeconomy if it is to be successful. These objectives will promote the renewal of our

industry, the modernization of our primary production systems and the protection of the environment and will help enhance biodiversity” [9]

“Many species of insects are capable of converting biological wastes into valuable products like compost and also are the rich source of nutrients. Insects reduce the above-mentioned societal challenges, create healthier and more sustainable food, and reduce animal feed production and consumption. Insects are rich in proteins (37–63%) and fats (20–40%), with well-balanced amino acid and fatty acid profiles and they are good sources of minerals and vitamins” [10]. “Insects have a high feed conversion rate, requiring much less feed to produce the same amount of animal proteins. Most of the insects can be cultured on locally available industrial and agricultural waste streams, recycling a loss into a valuable protein source. Households across all continents wasted over 1 billion meals a day in 2022, while 783 million people were affected by hunger and a third of humanity faced food insecurity” [11]. “Insects have great potential to convert these food waste into energy. Insect biorefinery is a concept of using insect as a tool to convert biomass waste into energy and other beneficial products with concomitant remediation of the organic components. The exploitation of insects and their bioproducts have become more popular in recent years” [12]. “In near future, insect sector can become an important component of sustainable circular agriculture by closing nutrient and energy cycles, fostering food security, and minimising climate change and biodiversity loss, thereby contributing to sustainable development” [13, 14]

“Black soldier fly *hermetia illucens* L. (Diptera: Stratiomyidae) is a nutritious feed component for livestock with high protein levels. Larvae and pupae rich in nutrients. They are known to convert organic waste into compost which is also rich in nutrients” [12,13,14]. Black soldier fly (BSF) can be reared on a wide variety of organic residues. Therefore, the production of BSF within a circular agriculture is feasible and that can help in reducing the import of expensive feed components, such as fishmeal or soymeal for livestock production.

2. Biology and distribution of BSF

“BSF is wasp-like fly found in rural, urban and forest environments. It is native to the Neotropics, but now found in every zoogeographic region following decades of spread throughout the warmer parts of the world” [15]. “It is found on every continent except Antarctica. They are endemic to the tropical and warm temperate regions of the Western Hemisphere, dispersing to the continents on the Eastern Hemisphere by human interactions”. [16,17]. “It is eurythermal species which can tolerate wide extremes of temperature. It consists of four life stages viz. egg, maggot, pupa and adult (Fig.1). Adult female lay eggs in cracks and crevices near food in a dry place. Eggs hatch in about more than 4 days at a temperature of 27-29°C to about 3.5 days at 30°C” [18]. “Maggot is white, translucent with reddish brown head. Larva passes through six instars. The larval body comprises 11 segments covered by hairs and bristles. The larvae consistently feed on organic waste and reduces it to compost. The mouthparts of the larval stage are very strong and are well adapted for feeding on waste matter. The last instar larva become greyish and does not feed. Larva moults into pupa in about 18 days. Pupa migrates towards the dry place away from the food substrate” [16]. “The life cycle last approximate 45 days [Egg (4 days), Larva(18 days), Pupa (14 days), and adult (9 days)]. The total

life cycle of BSF completes about 45 days but may extend up to four months” [19] depending upon composition of food substrate, temperature, humidity etc.




Figure 1. Biology of black soldier fly[20]

3. BSF for organic waste management and compost production

“Waste management is one of the most important challenges of the twenty-first century. Many aspects of societies, economies and the environment are influenced by this issue. Addressing this challenge also contributes towards the achievement of more than half of the SDGs especially those related to health, climate change, food security, poverty alleviation, responsible consumption and production” [21]. “This can be achieved through waste valorization” [22]. “BSF larvae feed voraciously on different organic wastes like animal manure *viz.* poultry, swine and dairy manure” [23-29] . Kitchen waste [30-31] Human faeces and fecal sludge [29, 31, 32, 33, 34, 35], municipal organic waste [37], fruit and vegetable waste [38, 39, 40, 41], spent grains [41], slaughterhouse waste, food processing waste [41, 42], brewery waste and local beer waste [43], bakery waste [44]), and fish waste [28, 37, 45, 46]. “Breakthrough research reveals that BSF larvae do not produce detectable methane emissions when fed various organic substrates” [47]. “This finding has significant implications for sustainable waste management and the transition to a circular bioeconomy. Bioconversion can be an effective and eco-friendly waste

management and resource recovery technique to eliminate organic waste leading to a circular economy model. However, further research is crucial to optimize insect-based bioconversion processes for permanent organic waste elimination. By advancing this innovative approach, we can harness the potential of insect-based bioconversion to create a more sustainable and regenerative food system” [48]. “The conversion of food waste by BSF larvae is heavily influenced by presence of gut microbiota viz. *Enterococcus*, *Klebsiella*, *Morganella*, *Providencia*, and *Scrofimicrobium*, which play a vital role in substrate utilization” [49]. This highlights the importance of microbial communities in sustainable waste management and nutrient cycling.

“Composting using BSF requires a shorter time (12–15 days) than composting using microbes or earthworms” [50]. “BSF larvae can degrade organic waste up to 72% and the maturity of the compost is very precise” [51]. “ICAR-NBAIR has termed BSF compost as black gold due to its qualities. No significant changes in nutrient composition and compost-maintained shelf life under room temperature. The C:N ratio recorded from compost varied between 17.8- 22.0:1. The analysis of mature BSF compost revealed a humic substance content of 11.70% (w/w), moisture 26.05 % (w/w), organic matter 19.43% (w/w), pH of 7.19 and bulk density of 0.57 gmL⁻¹”[52]. “BSF larvae can also reduce pathogen bacteria, such as *Escherichia sp.*, *Salmonella sp.*, *Vibrio sp.*, and *Yersinia sp.*, allowing the compost end product to meet the requirements for use as fertilizer and/or soil improver” [53]. “Some challenges arise regarding the use of BSF larvae as a bioconversion agent, such as for heavy metal residues, pesticide residues, pathogens, and antimicrobial gene transmission and residues that require the best composting strategy for mitigation” [54]. “Toxic elements can give bioaccumulation problems in BSF larvae and frass but the risk depends on their level in the growing substrate” [55].

4. BSF as animal feed:

“In the field of animal nutrition, BSF larvae have emerged as a highly viable and sustainable source of protein. This species has attracted attention due to some special reasons which have the potency to convert waste food materials and manure into high quality insect proteins. The high protein content in BSF larvae, constituting up to 50% of their dry weight, makes them a potential sustainable alternative to traditional protein sources in animal feeds” [56]. “BSF gained special attention among the aquaculturists and it has been utilized for feed formulation in aquaculture. BSF larvae have an average content of 45.2% crude protein and 31.4% fat. However, fat and protein composition can vary as a function of growing substrate characteristics” [57]. “Maggots prefer to accumulate protein food residues rather than carbohydrate food residues” [58]. “They present an environmentally friendly alternative to conventional feed ingredients such as soybean meal and fishmeal” [59]. “BSF larvae meal showed a positive impact on improving health conditions in fishes by various workers” [60-61]. Linh *et al.* (2024) revealed that “fish meal can be effectively replaced by BSF larvae meal and that this not only has

a positive effect on immune-related gene expression and growth rate growth rate in fishes” [62].“Research has revealed the positive effect of BSF diet on broiler chickens [63-65]. Inclusion of live BSF larvae did not affect fear behavior in laying hens, but the relative abundance of certain gut microbiota was associated with fear-related behaviors” [66]. “BSF generates a sustainable protein source as well as the importance of its use as a substitute of protein-rich feedstuff in poultry feed manufacturing. Producing BSF on waste streams as protein component for feed can be done locally, generating a circular approach either to agriculture on-farm or in the local community thus generating an economy that is not dependent on external inputs. BSF has been identified as the most versatile because of the variety of biological wastes that can be used for its rearing, automation, scalability, nutritional value, and because of circular and environmental aspects” [67]. Studies by Perez-Velazquez *et al.* (2024) revealed that “the fish diets with the levels of replacement of 50% and 75%, with final weights of 23.04 and 23.11 g, respectively, elicited superior fish performance, as compared to the diet with 25% replacement (final weight of 20.09 g)” [68]. “BSF do not carry bacteria or diseases and larvae are capable of inactivating *Escherichia coli* and *Salmonella enterica*” [69]. “In the EU, insects have been allowed as protein sources for aquaculture since 2017” [70]. BSF is also used as ingredient in the food of pigs [71-72] turkey [73], rabbits [74-75], ducks [76]and quails [77].

5. BSF as Human food: Edible insects can be used to solve the problem of food scarcity [78-79]. Food safety reports indicating the microbial risks associated with edible insect species based their suggested limits on EU safety regulations for meat and seafood, due to the high moisture and nutrient content of the insects [80-81]. Insects are not listed in the Codex Alimentarius, a United Nations document on what is considered “food” that informs much global food regulatory policy, except as impurities that contaminate food [82]. BSF do not concentrate pesticides or mycotoxins. For commercial use in human foods, larvae could potentially be milled and converted into a textured protein with a strong flavor [83]. BSF larvae and pupae are safe for humans to eat, their eggs are not. Few cases of enteric/intestinal myiasis caused by BSF maggots owing to the consumption of unwashed fruit on which BSF eggs were present, has been reported [84]. A case of cutaneous myiasis was also recorded due to the same reason [85]. However, this problem will be eliminated in case of dried, cooked, and/or powdered BSF products. Currently the regulations in the EU do not support the use of BSF maggots in human food [86].

6. BSF for chitin extraction

“Chitin is a linear polymer of N-acetyl-d-glucosamine. Chitin and chitosan find application in the food industry, agriculture, wastewater treatment, tissue engineering, biomedical, biotechnological, sanitary and cosmetic sectors, and in the textile and paper industries” (87-88). “The chitin market is estimated to reach USD 2.941 billion in 2027” [89]. “Insect exoskeletons are rich in chitin. The exoskeleton of BSF contains up to 35% chitin” [90]. “Chitin can be extracted from each stage of

BSF [91]. Also, pupal exuviae and dead adults of BSF are chitin-rich waste materials” (Purkayastha and Sarkar, 2020). “Therefore, continuous production of BSF can yield this high-quality biopolymer. BSF has an ability to produce chitin polymers or polymer of glucosamine up to 7% of BSF biomass on dry matter basis” [92]. Xiong *et al.* (2023) designed “chitin extraction from pupae and puparium of BSF using microbial treatments” [93]. “Chitosan from BSF slows down the decay of strawberries [94]. Thus, by recycling wastes, ultimately leading to the production of chitin and chitosan polymers, BSF fits well with the principles of circular economy” [95-96].

7. BSF for biodiesel production

“Insect fat has been proposed as a promising resource for biodiesel production” [97]. “Biodiesel production from BSF can be achieved by lipid extraction or by transesterification” [98; 99; 100, 101, 102]. “This consumes less energy and increases sustainability” [103]. “All the parameters of biodiesel produced from BSFL satisfied the Korea fuel standard except oxidation stability” [104]. “Worldwide research on BSF is on the right track towards the avenues of production of green bioenergy” [105].

8. Case studies:

Chineme *et al.* (2023) investigated “how an open BSF biowaste system could apply circular economy principles to close a resource loop in a low-income community and highlights the benefits of adopting a modified circular economy in Kipunguni, Dar es Salam”. [106] Through community involvement, circular economy principles were effectively implemented to redirect a retail market’s fruit and vegetable waste from the landfill. Corn bran was substituted by BSF larvae as animal feed, and frass biofertilizer replaced commercial fertilizer. Thus, demonstrating circularity in the study location. The free-range open system produced 19.15 kg of BSF larvae, with 44.34% protein content, 20.6% crude fat and zero heavy metals.

Using experimental and secondary data, Abro *et al.* (2020) assessed potential socio-economic benefits of BSF larvae meal to the Kenyan poultry sector. [107]. They concluded that replacing 5-50% of the conventional feed sources by BSF can generate a potential economic benefit of 69-687 million USD (0.1-1% of the total GDP) and 16-159 million USD (0.02-0.24% of the GDP) if the entire commercial poultry sector adopts BSF larvae meal. These could translate to reducing poverty by 0.32-3.19 million (0.07-0.74 million) people, increasing employment by 25,000-252,000 (3300-33,000) people, and recycling of 2-18 million (0.24-2 million) tonnes of biowaste.

9. Aspects of BSF based business industry

“There is increasing demand for protein-rich feed in animal industry, aquaculture, meat industry. Also, there is increasing government support for the use of insect meal in animal feeds. The BSFmarket is expected to reach **\$3.96 billion by 2033**, at a CAGR of 31% from 2024–2033, while in terms of volume, the market is expected to reach 8.23 million tons by 2033, at a CAGR of 40.4% from 2024 to 2033. It is important to work on the legal framework and the harmonization of a differentiated legal status of insects as food and feed across the world to facilitate the global use of this sustainable source and promote investments on both household and industrial production” [108]. Research emphasis on establishing new supply chains to support entrepreneurs in finding solutions that can address logistical challenges to experiment with more efficient production scales in the BSF industry, which may increase in accessibility and competitiveness.

10. BSF for achieving sustainable development goals (SDGs)

“SDGs target five critical areas: people, planet, prosperity, peace, and partnership, addressing pressing global challenges to ensure a better future for all. These goals are No poverty (**SDG 1**), Zero hunger (**SDG 2**), Good health and well-being (**SDG 3**), Quality education (**SDG 4**), Gender equality (**SDG 5**), Clean water and sanitation (**SDG 6**), Affordable and clean energy (**SDG 7**), decent work and economic growth (**SDG 8**), industry, innovation and infrastructure (**SDG 9**), reduced inequalities (**SDG 10**), sustainable cities and communities (**SDG 11**), responsible consumption and production (**SDG 12**), climate action (**SDG 13**), life below water (**SDG 14**), life on land (**SDG 15**), peace, justice, and strong institutions (**SDG 16**) and partnerships for the goals (**SDG 17**)” [1]. “BSF farming is getting popularized nowadays as an innovative approach for waste to wealth, which is said to meet 12 out of 17 sustainable development goals (no poverty, zero hunger, good health and well-being, gender equality, clean water and sanitation, decent work and economic growth, industry, innovation and infrastructure, reduced inequalities, sustainable cities and communities, responsible consumption and production, climate action, life below water, life on land, peace, justice and strong institutions(fig.2)”. [14]




Fig.2: Developing a circular economy based on the production of black soldier flies (BSF)[14]

11. BSF research status in India and future prospects

In India, natural occurrence of BSF is reported previously by Ashuma *et al* (2007) Gujarathiand Pejaver. (2013) and Pathak *et. al* (2015) in manure and compost. [110-112.] Sable *et al* (2024) reported natural occurrence of BSF at 16 locations from 10 districts of Maharashtra and recorded various types of organic wastes utilized as feed by BSF in natural settings [113]. The biology of BSF is studied by Sharanbasappa *et al.* (2019) on muskmelon [114]; Khyade (2021) on kitchen waste [115] ; Shrikant and Sharanbasappa (2021) on organic wastes [116] , Sable and Tawale, (2024) on fruit and vegetable wastes [40] and Sable *et al* (2024) on kitchen waste [113] . Bioconversion efficiency of BSF was studied on various

organic waste streams by Singh *et al.* (2021) [117] Ganvir *et al.*, (2022) [44] and Karthkeyani *et al.* (2024) [32]. Naik and Katkar (2020) proposed the BSF technique for waste management [118].

India faces the problem of huge amount of organic waste residues in rural and urban areas and BSF can be employed for the management of these residues. Small and medium holding farmers, poultry farmers and aquaculturists, can adopt the production of BSF on household kitchen waste and agricultural waste. BSF is currently mass produced as feed on a large scale in China, USA, Canada, South Africa and several European countries [119]. Similarly, BSF will be helpful for small holder and medium holding farmers in India. It is highly essential to explore the potential of BSF in India for waste management, compost making and animal feed production.

12. Conclusions:

The global challenges of organic waste generation, pollution, and costly animal feed necessitate innovative solutions. BSF larvae provide a sustainable answer, transforming organic waste into valuable compost and animal feed. Implementing BSF-based systems can significantly mitigate waste management issues, environmental pollution, and greenhouse gas emissions. The insect's high reproductive rate, adaptability, and nutritional value make it an attractive option for large-scale production. Studies have demonstrated its potential for compost, chitin, biodiesel production and even human nutrition. Overall, BSF research highlights its vast potential for contributing to circular economy. Potential of BSF can be harnessed to achieve sustainable development goals. However, further research is needed to optimize breeding, rearing, and processing techniques. Standardization of protocols and regulations is also essential for industrial-scale implementation. There is a scarcity of BSF research in India and the potential of BSF as feed food and composting agent in India remains unexplored. There is great scope for utilization of BSF for wellbeing of small and medium holding farmers. India's tropical climate, abundant organic waste, and growing demand for sustainable solutions make it an ideal location for BSF-based industries to thrive.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

REFERENCES

1. United Nations General Assembly, Resolution 70/1: Transforming our world: The 2030 agenda for sustainable development. 2015. http://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E.
2. Food and agriculture Organization. High-Level Expert Forum: Global agriculture towards 2050.2009
3. Food and agriculture Organization. The state of world fisheries and aquaculture: opportunities and challenges. 2016. Food and Agriculture Organization of the United Nations, Rome, Italy
4. Pinotti L, Caprarulo V, Ottoboni M, Giromini C, Agazzi A, Rossi L, Tretola M, Baldi A, Savoini G, Đuragić O. FEEDNEEDS: Trends in R&D in the Italian and Serbian feed sectors. In: Battinelli P, Striber J, editors. Italian-Serbian Bilateral Cooperation on Science, Technology and Humanities. Museum of Yugoslav History; 2016. pp. 21-25.
5. Makkar HPS, Ankers P. Towards sustainable animal diets: a survey-based study. *Anim Feed Sci Technol*. 2014; 198:309-322.
6. Some S, Roy J, Chatterjee JS, Butt MH. Low demand mitigation options for achieving sustainable development goals: role of reduced food waste and sustainable dietary choice. *J Clean Prod*. 2022;369:133432.
7. Ekins P, Domenech T, Drummond P, Bleischwitz R, Hughes N, Lotti L. The Circular Economy: What, Why, How and Where. Background paper for an OECD/EC Workshop; 2019.
8. Milios L. Advancing to a Circular Economy: three essential ingredients for a comprehensive policy mix. *Sustain Sci*. 2018;13:861-878.
9. European Commission (2018) A sustainable bioeconomy for Europe: strengthening the connection between economy society and the environment. European Commission-2018
10. Van Huis A, van Itterbeeck J, Klunder H, Mertens E, Halloran A, Muir G, Vantomme P. Edible Insects: Future Prospects for Food and Feed Security. Food and Agriculture Organization of the United Nations (FAO); 2013. p. 171.
11. United Nations Environment Programme (UNEP). (2024). The world squanders over 1 billion meals a day <https://www.unep.org/news-and-stories/press-release/world-squanders-over-1-billion-meals-day-un-report>
12. Kee PE, Cheng YS, Yiim HS, Tan CY, Lam SS, Lan JCW, Ng HS, Khoo KS. Insect biorefinery: A circular economy concept for biowaste conversion to value-added products. *Environ Res*. 2023;221.
13. Moruzzo R, Mancini S, Guidi A. Edible Insects and Sustainable Development Goals. *Insects*. 2021;12(6):557. (link unavailable)

14. Barragan-Fonseca KY, Barragan-Fonseca KB, Verschoor G, van Loon JJ, Dicke M. Insects for peace. *Curr Opin Insect Sci.* 2020;40:85–93.
15. Marshall SA, Woodley NE, Hauser M. The historical spread of the black soldier fly, *Hermetia illucens* (L.) (Diptera, Stratiomyidae, Hermetiinae), and its establishment in Canada. *J Ent Soc Ont.* 2015;146:51–4.
16. May B. The occurrence in New Zealand and the life-history of the soldier fly '*Hermetia illucens*'. *New Zealand Journal of Science.* 1961;4:55–65.
17. Rozkosny R. A biosystematics study of the European Stratiomyidae (Diptera). Dr W. Junk Publishers; 1982. Series Entomologica.
18. Tomberlin JK, Sheppard DC. Factors influencing mating and oviposition of black soldier flies (Diptera: Stratiomyidae) in a colony. *Journal of Entomological Science.* 2002;37:345–352.
19. Furman DP, Young RD, Catts PE. *Hermetia illucens* (Linnaeus) as a factor in the natural control of *Musca domestica* Linnaeus. *J Econ Entomol.* 1959;52:917–921.
20. ChiaSY. (2021). Black soldier fly larvae as a sustainable animal feed ingredient in Kenya. 10.18174/502357.
21. Wilson DC, Rodic L, Modak P, Soos R, Carpintero Rogero A, Velis C, Iyer M, Simonett O. Global waste management outlook. International Solid Waste Association (ISWA) and United Nations Environmental Programme (UNEP); 2015.
22. Lohri CR, Diener S, Zabaleta I, Mertenat A, Zurbrügg C. Treatment technologies for urban solid biowaste to create value products: A review with focus on low- and middle-income settings. *Rev Environ Sci Biotechnol.* 2017;16:81–130.
23. Sheppard DC, Newton GL, Thompson SA, Savage S. A value added manure management system using the black soldier fly. *Bioresource Technology.* 1994;50:275–279.
24. Yu G, Cheng P, Chen Y, Li Y, Yang Z, Chen Y, Tomberlin JK. Inoculating poultry manure with companion bacteria influences growth and development of black soldier fly (Diptera: Stratiomyidae) larvae. *Environmental Entomology.* 2011;40:30–35.
25. Myers HM, Tomberlin JK, Lambert BD, Kattes D. Development of black soldier fly (Diptera: Stratiomyidae) larvae fed dairy manure. *Environ Entomol.* 2008;37(1):11–15.
26. Li Q, Zheng L, Qiu N, Cai H, Tomberlin JK, Yu Z. Bioconversion of dairy manure by black soldier fly (Diptera: Stratiomyidae) for biodiesel and sugar production. *Waste Manag.* 2011;31:1316–1320.
27. Newton GL, Sheppard DC, Watson DW, Burtle GJ, Dove CR, Tomberlin JK, Thelen EE. The black soldier fly, *Hermetia illucens*, as a manure management/resource recovery tool. In: Symposium on the state of the science of animal manure and waste management, (ed.), Nowak P. San Antonio, Texas; 2005.

28. Nguyen TT, Tomberlin JK, Vanlaerhoven S. Ability of black soldier fly (Diptera: Stratiomyidae) larvae to recycle food waste. *Environ Entomol.* 2015;44:406–410.
29. Lalander C, Diener S, Zurbrügg C, Vinnerås B. Effects of feedstock on larval development and process efficiency in waste treatment with black soldier fly (*Hermetia illucens*). *J Clean Prod.* 2019;208:211-219.
30. Miranda CD, Cammack JA, Tomberlin JK. Life-History Traits of the Black Soldier Fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae), Reared on Three Manure Types. *Animals.* 2019;9(5):281.
31. Srikanth BH, Sharanabasappa Deshmukh. Growth performance and bioconversion rate of black soldier fly, *Hermetia illucens* (L.) when reared on organic feed wastes. *Indian J Entomol.* 2021;83:155-158.
32. Karthikeyani T, Sivasubramanian K, Maheswari M, Chitra N, Saravanan S, Jothimani P, Karthika S. The Efficiency of Black Soldier Fly Larvae with Vegetable, Fruit and Food Waste as Biological Tool for Sustainable Management of Organic Waste. *Int J Environ Clim Change.* 2024;14(2):441–448.
33. Lalander C, Diener S, Magri ME, Zurbrügg C, Lindström A, Vinnerås B. Faecal sludge management with the larvae of the black soldier fly (*Hermetia illucens*) — from a hygiene aspect. *Sci Total Environ.* 2013;458-460:312-318.
34. Banks IJ, Gibson WT, Cameron MM. Growth rates of black soldier fly larvae fed on fresh human faeces and their implication for improving sanitation. *Trop Med Int Health.* 2014;19(1):14-22.
35. Joly G. Valorising organic waste using the black soldier fly (*Hermetia illucens*), in Ghana. MSc thesis. KTH Royal Institute of Technology; 2018.
36. Diener S, Studt S, Nandayure M, Roa Gutiérrez F, Zurbrügg C, Tockner K. Biological treatment of municipal organic waste using black soldier fly larvae. *Waste Biomass Valoris.* 2011; 2:357-363.
37. Saragi ES, Bagastyo AY. Reduction of organic solid waste by black soldier fly (*Hermetia illucens*) larvae. The 5th Environmental Technology and Management Conference “Green Technology towards Sustainable Environment” November 23-24, 2015, Bandung, Indonesia. Cheng, J.Y.K.; Lo, I.M.C. (2016). Investigation of the available technologies and their feasibility for the conversion of food waste into fish feed in Hong Kong. *Environ Sci Pollut Res.* 23: 7169-7177.
38. Cheng JYK, Lo IMC. Investigation of the available technologies and their feasibility for the conversion of food waste into fish feed in Hong Kong. *Environ Sci Pollut Res.* 2016; 23:7169-7177.
39. Rampure SM, Velayuthannair K. Influence of agricultural wastes on larval growth phases of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae): An integrated approach. *J Appl Nat Sci.* 2023;15(2):860-869.
40. Sable A, Tawale J. Biology of black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae) on fruit and vegetable wastes. In: Proceedings of the National

- Conference on Recent Trends and Advances in Entomology; 2024 Mar 26-27; Entomology Research Institute, Loyola College, Chennai. p. 36.
41. Dortmans B, Diener S, Verstappen BM, Zurbrügg C. Black soldier fly biowaste processing - a step-by-step guide. Eawag: Swiss Federal Institute of Aquatic Science and Technology; 2017.
 42. Mohd-Noor SN, Wong CY, Lim JW, Mah-Hussin MIA, Uemura Y, Lam MK, Ramli A, Bashir MJK, Tham L. Optimization of self-fermented period of waste coconut endosperm destined to feed black soldier fly larvae in enhancing the lipid and protein yields. *Renew Energy*. 2017; 111:646-654.
 43. Sankara F, Pousga S, Coulibaly K, Nacoulma JP, Ilboudo Z, Ouédraogo I, Somda I, Kenis M. Optimization of Production Methods for Black Soldier Fly Larvae (*Hermetia illucens* L.) in Burkina Faso. *Insects*. 2023;14(9):776.
 44. Ganvir KP, Darvekar AN, Raut VD, Thorat RK. Effect of locally generated food waste on bioconversion and nutrient parameters of black soldier fly larvae, *Hermetia illucens* L. *J Entomol Zool Stud*. 2022;10(5):108-116.
 45. Liland NS, Sørensen M, Belghit I, Willora FP, Torrissen A, Torrissen O. Closing the gap – producing black soldier fly larvae on aquaculture side streams. *J Insects Food Feed*. 2023;9(7):885-892.
 46. St-Hilaire S, Cranfill K, McGuire MA, Mosley EE, Tomberlin JK, Newton GL, Sealey W, Sheppard DC, Irving S. Fish offal recycling by the black soldier fly produces a foodstuff high in omega-3 fatty acids. *J World Aquacult Soc*. 2007;38(2):309-313.
 47. Bava L, Jucker C, Gislou G, Lupi D, Savoldelli S, Zucali M, Colombini S. Rearing of *Hermetia illucens* on different organic by-products: influence on growth, waste reduction, and environmental impact. *Animals*. 2019; 9:289.
 48. Hamam M, D'Amico M, Di Vita G. Advances in the insect industry within a circular bioeconomy context: a research agenda. *Environ Sci Eur*. 2024; 36:29.
 49. Ganesan AR, Mohan K, Kandasamy S, Surendran RP, Kumar R, Rajan DK, Rajarajeswaran J. Food waste-derived black soldier fly (*Hermetia illucens*) larval resource recovery: A circular bioeconomy approach. *Process Saf Environ Prot*. 2024; 184:170-179.
 50. Purnamasari, L. and Himmatul K. Black Soldier Fly (*Hermetia illucens*) as a Potential Agent of Organic Waste Bioconversion. *ASEAN Journal on Science & Technology for Development*. 2022;39(2)69–83. DOI 10.29037/ajstd.780.
 51. Triadi RMA, Mirwan M, Farahdiba AU, Rositasari PR, Prawardani S. Application of composting technology to reduce organic waste by utilizing black soldier fly (*Hermetia illucens*) larvae. In: 3rd International Conference Eco-Innovation in Science, Engineering, and Technology; 2022. NST Proceedings. p. 91-94. doi: 10.11594/nstp.2022.2715
 52. Indian Council of Agricultural Research - National Bureau of Agricultural Insect Resources (ICAR-NBAIR). Shelf-life studies on black soldier fly compost. Annual Report 2019. 2019;33p.

53. Awasthi MK, Liu T, Awasthi SK, Duan Y, Pandey A, Zhang Z. Manure pretreatments with black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae): A study to reduce pathogen content. *Sci Total Environ.* 2020; 737:139842. doi: 10.1016/j.scitotenv.2020.139842.
54. Rajasekar B, Swetha TN, Harshita DN, Mishra P, Hudge B. A review on black soldier fly *Hermetia illucens* as a potential source for organic waste management. *Pharma Innovation.* 2023;12(6):369-3474.
55. Addeo NF, Scivicco M, Vozzo S, Bovera F, Asiry KA, Alqurashi S, Cacciola NA, Severino L. Mineral profile and heavy metals bioaccumulation in black soldier fly (*Hermetia illucens*, L.) larvae and frass across diverse organic substrates. *Italian J Anim Sci.* 2024;23(1):179-188. DOI: 10.1080/1828051X.2024.2302845
56. Shumo M, Osuga IM, Khamis FM, Tanga CM, Fiaboe KK, Subramanian S, Ekesi S, van Huis A, Borgemeister C. The nutritive value of black soldier fly larvae reared on common organic waste streams in Kenya. *Sci Rep.* 2019;9:10110.
57. Makkar HP, Tran G, Heuzé V, Ankers P. State-of-the-art on use of insects as animal feed. *Anim Feed Sci Technol.* 2014;197:1–33.
58. Jalil NAA, Abdullah SH, Ahmad IK, Basri NEA, Mohamed ZS. Decomposition of food waste from protein and carbohydrate sources by black soldier fly larvae, *Hermetia illucens* L. *J Environ Biol.* 2021;42:756-761.
59. Salahuddin M, Abdel-Wareth AAA, Hiramatsu K, Tomberlin JK, Luza D, Lohakare J. Flight toward Sustainability in Poultry Nutrition with Black Soldier Fly Larvae. *Animals.* 2024;14(3):510.
60. Yandigeri M, Amala U. Black Soldier Fly, a detritivorous insect as a waste decomposer and raw material for aquacultural feed. *EnviroWorld.* 2019:21–22.
61. Amala U, Yandigeri MS, Mohan M, Panikkar P, Jesna PK, Khan FM, Das BK, Ballal CR, Selvaraj K, Vijaykumar ME. Black soldier fly – a promising protein supplement in aquacultural feed. *ICAR News.* 2019;25(3):27.
62. Linh NV, Wannavijit S, Tayyamath K, Dinh-Hung N, Nititanarapee T, Sumon MAA, Srinual O, Permpoonpattana P, Doan HV, Brown CL. Black Soldier Fly (*Hermetia illucens*) Larvae Meal: A Sustainable Alternative to Fish Meal Proven to Promote Growth and Immunity in Koi Carp (*Cyprinus carpio* var. koi). *Fishes.* 2024;9(2):53.
63. Pieterse E, Erasmus SW, Uushona T, Hoffman LC. Black soldier fly (*Hermetia illucens*) pre-pupae meal as a dietary protein source for broiler production ensures a tasty chicken with standard meat quality for every pot. *J Sci Food Agric.* 2019;99:893–903. doi: 10.1002/jsfa.9261.
64. Schiavone A, Dabbou S, De Marco M, Cullere M, Biasato I, Biasibetti E, Capucchio M, Bergagna S, Dezzutto D, Meneguz M. Black soldier fly larva fat inclusion in finisher broiler chicken diet as an alternative fat source. *Animal.* 2018;12:2032–2039. doi: 10.1017/S1751731117003743.

65. Schiavone A, Dabbou S, Petracci M, Zampiga M, Sirri F, Biasato I, Gai F, Gasco L. Black soldier fly defatted meal as a dietary protein source for broiler chickens: Effects on carcass traits, breast meat quality and safety. *Animal*. 2019;13:2397–2405. doi: 10.1017/S1751731119000685.
66. Huang C, Hernandez CE, Wall H, Tahamtani FM, Ivarsson E, Sun L. Live black soldier fly (*Hermetia illucens*) larvae in feed for laying hens: effects on hen gut microbiota and behavior. *Poult Sci*. 2024;103(3).
67. Maroušek J, Strunecký O, Maroušková A. Insect rearing on biowaste represents a competitive advantage for fish farming. *Rev Aquac*. 2023;15:965–975.
68. Perez-Velazquez M, Millanes-Mora MA, González-Félix ML. Assessment of hydrolysis of partially defatted black soldier fly larvae meal in diets for Nile tilapia *Oreochromis niloticus*. *Anim Feed Sci Technol*. 2024;307:115831.
69. Erickson CM, Islam M, Sheppard C, Liao J, Doyle P. Reduction of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar enteritidis in chicken manure by larvae of the black soldier fly. *J Food Prot*. 2004;67:685–690.
70. European Commission. Regulation (EU) 893/2017 of 24 May 2017 amending Annexes I and IV to Regulation (EC) No 999/2001 of the European Parliament and of the Council and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011 as regards the provisions on processed animal protein. *Off J Eur Union*. 2017;138:92–116.
71. Biasato I, Renna M, Gai F, Dabbou S, Meneguz M, Perona G, Martinez S, Lajusticia ACB, Bergagna S, Sardi L. Partially defatted black soldier fly larva meal inclusion in piglet diets: Effects on the growth performance, nutrient digestibility, blood profile, gut morphology and histological features. *J Anim Sci Biotechnol*. 2019;10:12. doi: 10.1186/s40104-019-0325-x.
72. Yu M, Li Z, Chen W, Rong T, Wang G, Wang F, Ma X. Evaluation of full-fat *Hermetia illucens* larvae meal as a fishmeal replacement for weanling piglets: Effects on the growth performance, apparent nutrient digestibility, blood parameters and gut morphology. *Anim Feed Sci Technol*. 2020;264:114431. doi: 10.1016/j.anifeedsci.2020.114431.
73. Sypniewski J, Kierończyk B, Benzertiha A, Mikołajczak Z, Pruszyńska-Oszmałek E, Kołodziejcki P, Sassek M, Rawski M, Czekala W, Józefiak D. Replacement of soybean oil by *Hermetia illucens* fat in turkey nutrition: Effect on performance, digestibility, microbial community, immune and physiological status and final product quality. *Br Poult Sci*. 2020;61:294–302. doi: 10.1080/00071668.2020.1716302.
74. Dalle Zotte A, Cullere M, Martins C, Alves SP, Freire JP, Falcão-e-Cunha L, Bessa RJ. Incorporation of Black Soldier Fly (*Hermetia illucens* L.) larvae fat or extruded linseed in diets of growing rabbits and their effects on meat quality traits including detailed fatty acid composition. *Meat Sci*. 2018;146:50–58. doi: 10.1016/j.meatsci.2018.08.002.

75. Martins C, Cullere M, Dalle Zotte A, Cardoso C, Alves SP, Bessa R, Freire JPB, Falcao-e-Cunha L. Incorporation of two levels of black soldier fly (*Hermetia illucens* L.) larvae fat or extruded linseed in diets of growing rabbits: Effects on growth performance and diet digestibility. *Czech J Anim Sci.* 2018;63:356–362. doi: 10.17221/22/2018-CJAS.
76. Gariglio M, Dabbou S, Crispo M, Biasato I, Gai F, Gasco L, Piacente F, Odetti P, Bergagna S, Plachà I. Effects of the dietary inclusion of partially defatted black soldier fly (*Hermetia illucens*) meal on the blood chemistry and tissue (Spleen, Liver, Thymus, and Bursa of Fabricius) histology of muscovy ducks (*Cairina moschata domestica*). *Animals.* 2019;9:307. doi: 10.3390/ani9060307.
77. Cullere M, Tasoniero G, Giaccone V, Miotti-Scapin R, Claeys E, De Smet S, Dalle Zotte A. Black soldier fly as dietary protein source for broiler quails: Apparent digestibility, excreta microbial load, feed choice, performance, carcass and meat traits. *Animal.* 2016;10:1923–1930. doi: 10.1017/S1751731116001270.
78. Martin D. *Edible: An Adventure into the World of Eating Insects and the Last Great Hope to Save the Planet.* Houghton Mifflin Harcourt: Boston, MA, USA, 2014.
79. Nadeau L, Nadeau I, Franklin F, Dunkel F. The potential for entomophagy to address undernutrition. *Ecol Food Nutr.* 2015;55:200–208.
80. European Commission. Regulation (EC) No. 2073/2005 on microbiological criteria for foodstuffs. *Off J Eur Union.* 2005;338:1–26.
81. NVWA (Netherlands Food and Consumer Product Safety Authority). *Advisory Report on the Risks Associated with the Consumption of Mass-Reared Insects.* 2014.
82. Van Huis A, Van Itterbeeck J, Klunder H, Mertens E, Halloran A, Muir G, Vantomme P. *Edible Insects. Future Prospects for Food and Feed Security.* FAO: Rome, Italy, 2013; Volume 171, ISBN 978-92-5-107595-1.
83. Wang Y-S, Shelomi M. Review of Black Soldier Fly (*Hermetia illucens*) as Animal Feed and Human Food. *Foods.* 2017;6(10):91.
84. Meleney HE, Harwood PD. Human intestinal myiasis due to the larvae of the soldier fly, *Hermetia illucens* L. (diptera, stratiomyidae). *Am J Trop Med Hyg.* 1935;1:45–49.
85. Lee HL, Chandrawathani P, Wong WY, Tharam S, Lim WY. A case of human enteric myiasis due to larvae of *Hermetia*.
86. Adler AI, Brancato FP. Human Furuncular Myiasis Caused by *Hermetia illucens* (Diptera: Stratiomyidae). *J Med Entomol.* 1995;32(5):745–746. doi: 10.1093/jmedent/32.5.745.
87. Besler M, Steinhart H, Paschke A. Stability of food allergens and allergenicity of processed foods. *J Chromatogr B Biomed Sci Appl.* 2001;756:207–228. doi: 10.1016/S0378-4347(01)00110-4.

88. Bakshi PS, Selvakumar D, Kadirvelu K, Kumar NS. Chitosan as an environment friendly biomaterial – A review on recent modifications and applications. *Int J Biol Macromol*. 2020;150:1072–1083.
89. Taranathan RN, Kittur FS. Chitin—The undisputed biomolecule of great potential. *Crit Rev Food Sci Nutr*. 2003;43:61–87.
90. Shamshina JL, Berton P, Rogers RD. Advances in Functional Chitin Materials: A Review. *ACS Sustain Chem Eng*. 2019;7:6444–6457.
91. Thomas H, Elena T, Aman P, Rosanna S, Patrizia F, Susanne Z. Current state of chitin purification and chitosan production from insects. *J Chem Technol Biotechnol*. 2020;95:2775–2795.
92. Wu CS, Wang SS. Bio-based electrospun nanofiber of polyhydroxyalkanoate modified with Black Soldier Fly's pupa shell with antibacterial and cytocompatibility properties. *ACS Appl Mater Interfaces*. 2018;10:42127–42135.
93. Purkayastha D, Sarkar S. Physicochemical structure analysis of chitin extracted from pupa exuviae and dead imago of Wild Black Soldier Fly (*Hermetia illucens*). *J Polym Environ*. 2020;28:445–457.
94. Surendra K, Olivier R, Tomberlin JK, Jha R, Khanal SK. Bioconversion of organic wastes into biodiesel and animal feed via insect farming. *Renewable Energy*. 2016;98:197–202.
95. Xiong A, Ruan L, Ye K, Huang Z, Yu C. Extraction of Chitin from Black Soldier Fly (*Hermetia illucens*) and Its Puparium by Using Biological Treatment. *Life (Basel)*. 2023;13(7):1424.
96. Triunfo M, Guarnieri A, Ianniciello D, Coviello L, Vitti A, Nuzzaci M, Salvia R, Scieuzo C, Falabella P. *Hermetia illucens*, an innovative and sustainable source of chitosan-based coating for postharvest preservation of strawberries. *iScience*. 2023;26(12):108576.
97. Mannucci A, Panariello L, Abenaim L, Coltelli MB, Ranieri A, Conti B, Santin M, Castagna A. From Food Waste to Functional Biopolymers: Characterization of Chitin and Chitosan Produced from Prepupae of Black Soldier Fly Reared with Different Food Waste-Based Diets. *Foods*. 2024;13:278.
98. Lai EPC. Biodiesel: environmental friendly alternative to petrodiesel. *J Pet Environ Biotechnol*. 2015;5:122.
99. Li Q, Zheng L, Cai H, Garza E, Yu Z, Zhou S. From organic waste to biodiesel: black soldier fly, *Hermetia illucens*, makes it feasible. *Fuel*. 2011;90:1545–1548.
100. Zheng L, Hou Y, Li W, Yang S, Li Q, Yu Z. Biodiesel production from rice straw and restaurant waste employing black soldier fly assisted by microbes. *Energy*. 2012;47:225–229.
101. Elsayed M, Ran Y, Ai P, Azab M, Mansour A, Jin K, Zhang Y, Abomohra AEF. Innovative integrated approach of biofuel production from agricultural wastes by anaerobic digestion and black soldier fly larvae. *Journal of Cleaner Production*. 2020;263:121495.

102. Jung S, Jung JM, Yiu Fai Tsang YF, Bhatnagar A, Chen W-H, Lin K-YA, Kwon EE. Biodiesel production from black soldier fly larvae derived from food waste by non-catalytic transesterification. *Energy*. 2022;238:121700.
103. Mohan K, Sathishkumar P, Rajan DK, Rajarajeswaran J, Ganesan AR. Black soldier fly (*Hermetia illucens*) larvae as potential feedstock for the biodiesel production: Recent advances and challenges. *Sci Total Environ*. 2023;859(Pt 1):160235.
104. Park JY, Jung S, Na YG, Jeon CH, Cheon HY, Yoon EY, Lee SH, Kwon EE, Kim JK. Biodiesel production from the black soldier fly larvae grown on food waste and its fuel property characterization as a potential transportation fuel. *Environmental Engineering Research*. 2022;27(3):200704.
105. Mangindaan D, Kaburuan ER, Meindrawan B. Black Soldier Fly Larvae (*Hermetia illucens*) for Biodiesel and/or Animal Feed as a Solution for Waste-Food-Energy Nexus: Bibliometric Analysis. *Sustainability*. 2022;14:13993.
106. Chineme A, Assefa G. Open and Closed Black Soldier Fly Systems Tradeoff Analysis. *Sustainability*. 2023;15(24):16677.
107. Abro Z, Kassie M, Tanga C, Beesigamukama D, Diro G. Socio-economic and environmental implications of replacing conventional poultry feed with insect-based feed in Kenya. *J Clean Prod*. 2020;265.
108. Black Soldier Fly Market by Product, by Application, and Geography - Global Forecast to 2033. [researchandmarkets.com/reports/4986109/black-soldier-fly-market-by-product-by#:~:text=In%20terms%20of%20value%2C%20the,8%2C003.7%20thousand%20tons%20by%202033.](https://www.researchandmarkets.com/reports/4986109/black-soldier-fly-market-by-product-by#:~:text=In%20terms%20of%20value%2C%20the,8%2C003.7%20thousand%20tons%20by%202033.)
109. Moruzzo R, Mancini S, Guidi A. Edible Insects and Sustainable Development Goals. *Insects*. 2021; 12:557.
110. Ashuma BM, Kaur P, Rath S, Juyal P. First report of *Hermetia illucens* larvae in poultry houses of Punjab. *Journal of Parasitic Diseases*. 2007;31(2):145-146.
111. Gujarathi GR, Pejaver MK. Occurrence of Black Soldier Fly *Hermetia illucens* (Diptera: Stratiomyidae) in Biocompost. *Research Journal of Recent Sciences*. 2013;2(4):65-66.
112. Pathak R, Sharma S, Prasad R. Study on occurrence of black soldier fly larvae in composting of kitchen waste. *International Journal of Research in Biosciences*. 2015;4(4):38-45.
113. Sable A and Chavan K. Natural occurrence and biology of black soldier fly, *Hermetia illucens* L. (Diptera: Stratiomyidae) under local conditions of Maharashtra, India. *Uttar Pradesh Journal of Zoology*. 2024;45(18):10-25

114. Sharanabasappa D, Srikanth BH, Maruthi MS, Pavithra HB. Biology of black soldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae) on muskmelon fruit. Indian Journal of Entomology. 2019; 81:153-155.
115. Khyade VB. Rearing of black soldier fly *Hermetia illucens* (Linnaeus) (Diptera: Stratiomyidae) in local conditions of Baramati (India). Uttar Pradesh Journal of Zoology. 2021;42(5):64-72.
116. Srikanth BH, Sharanabasappa D. Growth performance and bioconversion rate of black soldier fly, *Hermetia illucens* (L.) when reared on organic feed wastes. Indian Journal of Entomology. 2021; 83:155-158.
117. Singh A, Srikanth BH, Kumari K. Determining the Black Soldier fly larvae performance for plant-based food waste reduction and the effect on Biomass yield. Waste Manag. 2021;130:147-154. doi: 10.1016/j.wasman.2021.05.028. Erratum in: Waste Manag. 2022;144:122. PMID: 34090238.
118. Naik SN, Katkar AA. Innovative Approach towards Solid Waste Management on Agonda Beach in Goa. International Research Journal of Engineering and Technology. 2020;7(6):2049-2055.
119. SY, Tanga CM, van Loon JJA, Dicke M. Insects for sustainable animal feed: inclusive business models involving smallholder farmers. Curr Opin Environ Sustain. 2019;41:23-30.

ABBREVIATIONS

BSF- Black soldier fly, SDG- Sustainable development goal