

Original Research Article

Effect of Pruning and Micronutrient Sprays on Physico-chemical Properties of Ber

ABSTRACT

An investigation was undertaken to study the effect of pruning height and micronutrients on the physicochemical properties of ber. The experiment was laid out in the farmers field at conducted in the farmers' field at the Manikpur area of Bongaigaon district, Assam (26°26'51" North latitude and 90°46'47" East longitude) for two consecutive years. Two types of Thailand ber were exposed to two different pruning heights and eight different types of micronutrient spray combinations. Results depicted that the Roundish ovate reddish type variety performed better. The highest total sugar content (14.91%), TSS/acid ratio (44.07) and ash content (2.66%) were observed in, and ash content (2.66%) were observed at a pruning height of 75 cm. Among micronutrients, borax at 0.5% and zinc sulphate sulphate (ZnSO₄) at 0.5% in combination proved to be better in terms of regarding total sugar (14.83%) and ash content (2.64%). So, the experiment proved that pruning height of 75 cm from the ground level and spraying of borax 0.5% and zinc sulphate 0.5% in combination were better compared to the other treatments.

Formatted: Subscript

Keywords: Ash content; Ber; Sugar; Titratable acidity; TSS

1. INTRODUCTION

Ziziphus mauritiana (Lamk.), the Indian ber of Rhamnaceae family, occupies an important place among minor fruits of India. India produces around 513 thousand metric tons of ber annually from an area of around 50 thousand hectares [1]. This fruit crop is in cultivation in India from has been in cultivation in India since very ancient times. The sage Vedavyasa was also popular as 'Badrayan' (the person residing in the ber forest) as he created his cottage in the forest where plenty of ber plants were there [2]. The nutritious fruit ber is superior to apple in terms of vitamin C, calcium, phosphorus, and protein [3].

Comment [VM1]: Values of nutritional contents in ber can be included in brief.

Annual pruning is mandatory in ber as it bears fruits on the current season growth [3]. Again, it is noticed that micronutrients play vital role in fruit set, fruit retention, fruit development and has been noticed that micronutrients play a vital role in fruit set, fruit retention, fruit development, and the quality of ber. Boron proved its efficiency in increasing the blooming and retention of flowers along with and developing development of seed seeds and fruit [4]. The stimulation of carbohydrates and proteins in plant cells as well as formation of DNA are regulated by the micronutrient zinc, besides its role in, as well as the formation of DNA, are regulated by the micronutrient zinc, besides its role in the biosynthesis of indole-3-butyric acid [5]. Therefore, an experiment was laid out to assess the role of pruning and micronutrients in the physico-chemical conducted to assess the role of pruning and

micronutrients in the physicochemical properties of two types of Thailand ber variety varieties under Assam condition conditions.

2. MATERIAL AND METHODS

The experiment was conducted for two years during 2020-21 and 2021-22, 2020-21 and 2021-22, in a farmers' field located at Manikpur area of Bongaigaon district, Assam, India (26°26'51" North latitude and 90°46'47" East longitude). The soil was strongly acidic (pH 5.1) with in the Manikpur area of Bongaigaon district, Assam, India (26°26'51" N) and 90°46'47" E). The soil was strongly acidic (p^H 5.1) with a high organic carbon content (1.03%).

Formatted: Superscript

Two types of Thailand Thailandbers were selected as study material, viz, V₁: Round green type, V₂: Roundish ovate reddish type. Heading back system of pruning was followed, and two different height heights of pruning were considered for the experiment (P₁: 50 cm from the ground level and P₂: 75 cm from the ground level). Pruning was performed just after harvest during March. Borax (11% B) and zinc sulphate ($ZnSO_4$) (33% Zn) were the two micronutrient sources taken for the study with various levels as N₀: No micronutrient, N₁: Borax 0.4%, N₂: Borax 0.5%, N₃: $ZnSO_4$ 0.4%, N₄: $ZnSO_4$ 0.5%, N₅: Borax 0.4% + $ZnSO_4$ 0.4%, N₆: Borax 0.4% + $ZnSO_4$ 0.5%, N₇: Borax 0.5% + $ZnSO_4$ 0.4%, N₈: Borax 0.5% + $ZnSO_4$ 0.5%. Spraying of micronutrients were done twice, first at fruit set and the next one is The spraying of micronutrients was done twice, first at the fruit set and the next one 30 days after the first spray.

Comment [VM2]: Include the name of the manufacturer and country of origin

Formatted: Subscript

Comment [VM3]: How the micronutrients were sprayed, which sprayer was used and during what time of day, it was sprayed?

2.1. Moisture content (%)

The moisture content was determined by following standard procedure with the help of the following formula –

$$\text{Moisture content (\%)} = \frac{\text{Fresh weight (g)} - \text{Dry weight (g)}}{\text{Fresh weight (g)}} \times 100$$

2.2. Total soluble solids (TSS) ($^{\circ}$ Brix) content

The fresh fruit was after removal of seed. To make the homogenous mixture, the cut pieces were ground. Then, juice was extracted and filtered with a muslin cloth. A drop of juice was placed in the digital hand refractometer to find measure the TSS ($^{\circ}$ Brix) at room temperature.

Comment [VM4]: Correct the sentence since it is difficult to comprehend.

Comment [VM5]: Name of the manufacturer and country of origin

2.3. Reducing sugar (%)

Lane-Eynon method was used for determining the reducing sugar.

Comment [VM6]: Missing reference and year

$$\text{Reducing sugar (\%)} = \frac{\text{mg of invert sugar} \times \text{dilution} \times 100}{\text{titre value} \times \text{weight of sample}}$$

Where, mg of invert sugar = Factor 0.05

2.4. Total sugar (%)

To determine the total sugar, the Lane-Eynon method was executed.

Comment [VM7]: Missing reference and year

$$\text{Total invert sugar (\%)} = \frac{\text{mg of invert sugar} \times \text{dilution} \times \text{second volume make up} \times 100}{\text{titre value} \times \text{weight of sample} \times \text{sample kept overnight}}$$

Where, mg of invert sugar = Factor 0.05

$$\text{Sucrose (\%)} = \{\text{Total invert sugar (\%)} - \text{Reducing sugar (\%)}\} \times 0.95$$

$$\text{Total sugar (\%)} = \text{Sucrose (\%)} + \text{Reducing sugar (\%)}$$

2.5. Non-reducing sugar (%)

The value of non-reducing sugar was directly calculated by using following formula –

$$\text{Non-reducing sugar (\%)} = \text{Total sugar (\%)} - \text{Reducing sugar (\%)}$$

Comment [VM8]: Missing reference and year

2.6. Titratable acidity (%)

~~Determination of titratable acidity in percentage was done~~The titratable acidity (%) was determined following the procedure prescribed by AOAC [6]. Titratable acidity was calculated in terms of percentage anhydrous citric acid.

$$\text{Titratable acidity (\%)} = \frac{\text{Titre value} \times N \times V \times \text{equivalent weight of citric acid} \times 100}{\text{weight of sample} \times \text{aliquot} \times 1000}$$

Where, N = Normality of alkali and V = Volume made up

2.7. TSS/acid ratio

The ratio of TSS and acid is an important aspect in relation to consumer preference. It was calculated out by dividing the value of TSS by the value of titratable acidity.

2.8. Ash content (%)

The ash content was determined by burning out all the organic material present in the sample in a temperature of 600°C in a muffle furnace [6]. The following calculation was used to find out the ash content –

$$\text{Ash (\%)} = \frac{W_1 - W}{W_2 - W} \times 100$$

Where, W = Weight of the crucible

W₁ = Weight of sample heated at 600°C

W₂ = Weight of crucible + sample

The relevant parameters of the study were subjected to standard statistical procedures [7].

Comment [VM9]: Please provide detailed information about the design of the experiment used in this study. Additionally, specify the significance test employed to analyze the data, such as univariate or multivariate ANOVA, and indicate the software utilized for data analysis.

3. RESULTS AND DISCUSSION

~~For different physico-chemical parameters record~~The recorded physico-chemical parameters were taken as per standard protocol as mentioned in the materials and methods and are presented and discussed hereunder.

3.1. Moisture content (%)

It is evident from Table 1 that only variety influenced the fruit moisture content significantly, whereas the pruning height as well as micronutrient sprays had no significant effect on fruit moisture. The maximum moisture of 79.36 per cent was recorded for variety V₁. The result Table 1 shows that only variety significantly influenced the fruit moisture content, whereas pruning height and micronutrient sprays had no significant effect on fruit moisture. Variety V₁ recorded maximum moisture of 79.36 percent, which is in conformity with the works of other research workers [8].

Table 1. Moisture and TSS content as influenced by variety, pruning height, and micronutrient sprays

| Treatments | Moisture (%) | | | TSS (°Brix) | | |
|----------------------------|--------------|-------------|-------------|-------------|-------------|-------------|
| | 2020 – 21 | 2021 – 22 | Pooled | 2020 – 21 | 2021 – 22 | Pooled |
| V ₁ | 79.73 | 78.98 | 79.36 | 13.53 | 13.46 | 13.49 |
| V ₂ | 78.29 | 78.76 | 78.52 | 14.91 | 15.09 | 15.00 |
| CD_{P=0.05} | 1.18 | 1.00 | 0.70 | 0.07 | 0.05 | 0.04 |
| P ₁ | 79.23 | 78.59 | 78.91 | 13.86 | 13.92 | 13.89 |
| P ₂ | 78.79 | 79.15 | 78.97 | 14.58 | 14.63 | 14.61 |
| CD_{P=0.05} | NS | NS | NS | 0.07 | 0.05 | 0.04 |
| N ₀ | 79.46 | 79.35 | 79.41 | 13.91 | 13.99 | 13.95 |
| N ₁ | 78.88 | 79.25 | 79.07 | 14.01 | 14.07 | 14.04 |
| N ₂ | 79.49 | 77.96 | 78.73 | 14.09 | 14.12 | 14.10 |
| N ₃ | 78.91 | 79.38 | 79.15 | 14.16 | 14.18 | 14.17 |
| N ₄ | 78.68 | 80.00 | 79.34 | 14.20 | 14.22 | 14.21 |
| N ₅ | 79.56 | 78.10 | 78.83 | 14.27 | 14.32 | 14.29 |
| N ₆ | 77.10 | 79.45 | 78.28 | 14.35 | 14.43 | 14.39 |
| N ₇ | 79.88 | 79.09 | 79.49 | 14.46 | 14.53 | 14.49 |
| N ₈ | 79.13 | 77.23 | 78.18 | 14.53 | 14.63 | 14.58 |
| CD_{P=0.05} | NS | NS | NS | 0.15 | 0.10 | 0.09 |

Comment [VM10]: Levels of significance indicating statistical significance (such as a, b, c, d) must be included in the table.

3.2. TSS (°Brix)

All the three factors viz., variety, pruning height and application of micronutrients three factors viz., variety, pruning height and application of micronutrients, significantly affected the total soluble solid content in fruits (Table 1). A TSS value of 15.00 °Brix in the variety V₂ was the highest and 4.61 °Brix was noticed in pruning height P₂. This resulted in higher accumulation of photosynthates and minerals in pruned trees leading to the highest content of TSS in medium pruned medium pruned trees [9]. Medium pruning (P₂) performed better in terms regarding the TSS content of peach cv. Flordaking [10]. Among the micronutrients, the highest TSS was found in N₈ (14.58 °Brix) and the lowest in N₀ (13.95 °Brix). Researchers also observed significant differences among various ber varieties [11]. Open An open plant canopy allows more sunlight penetration. Boron alone, or in association with Zn had positive influence on positively influenced fruit TSS content. Boron mediated transport of

Formatted: Not Superscript/ Subscript

Formatted: Font: Italic

~~higher assimilates might be the reason behind~~ Boron-mediated transport of higher assimilates might be the reason for this [12].

3.3. Total sugar (%)

As depicted by ~~the Table 2, amount of total sugar in fruit was significantly influenced by variety, pruning~~ Table 2, the amount of total sugar in fruit was significantly influenced by ~~variety, pruning~~ and micronutrient treatments. Among varieties, V₂ showed significantly higher ~~amount of total sugar content (15.14%) compared to total sugar content (15.14%) than~~ V₁ (13.54%). The significant deviation in total sugar content of fruits due to variety tallies the experimental finding of [11]. ~~Pruning height of 75 cm from the ground level (P₂) found to be significantly superior to A pruning height of 50 cm from the ground level (P₁) in terms of total sugar content with values of 14.91 and 13.77 per cent~~ pruning height of 75 cm from the ground level (P₂) was significantly superior to the pruning height of 50 cm from the ground level (P₁) in terms of total sugar content with values of 14.91 and 13.77 per cent, respectively. Similarly, higher amount of total sugar was achieved in 50 per cent pruning intensity due to a higher amount of total sugar was achieved at 50 percent pruning intensity due to the accumulation of more photosynthates and nutrients in pruned shoots [13]. Among micronutrient levels, N₈ showed significantly higher ~~content of total sugar (14.83%) compared to total sugar content (14.83%) than~~ all other levels, whereas N₀ exhibited the lowest quantity of total sugar (13.86%). The effect of boron can be attributed to its vital role in ~~translocation of sugar by formation of the translocation of sugar by the formation of a 'sugar-borate complex' with sugar in higher plants [14]. Zinc-mediated metabolism of starch and nucleic acid might be the cause for Zinc-mediated metabolism of starch and nucleic acid might be the cause for~~ the positive effect of this micronutrient in increasing the total sugar content [15].

Table 2. Total and reducing sugar content as influenced by variety, pruning height, and micronutrient sprays

| Treatments | Total sugar (%) | | | Reducing sugar (%) | | |
|----------------------------|-----------------|-------------|-------------|--------------------|-------------|-------------|
| | 2020 – 21 | 2021 – 22 | Pooled | 2020 – 21 | 2021 – 22 | Pooled |
| V ₁ | 13.58 | 13.49 | 13.54 | 7.41 | 7.32 | 7.37 |
| V ₂ | 15.26 | 15.03 | 15.14 | 8.13 | 8.03 | 8.08 |
| CD_{P=0.05} | 0.06 | 0.05 | 0.04 | 0.03 | 0.03 | 0.02 |
| P ₁ | 13.84 | 13.69 | 13.77 | 7.51 | 7.40 | 7.45 |
| P ₂ | 15.00 | 14.83 | 14.91 | 8.04 | 7.94 | 7.99 |
| CD_{P=0.05} | 0.06 | 0.05 | 0.04 | 0.03 | 0.03 | 0.02 |
| N ₀ | 13.92 | 13.80 | 13.86 | 7.54 | 7.47 | 7.50 |
| N ₁ | 14.05 | 13.92 | 13.99 | 7.60 | 7.52 | 7.56 |
| N ₂ | 14.17 | 14.03 | 14.10 | 7.65 | 7.57 | 7.61 |
| N ₃ | 14.31 | 14.17 | 14.24 | 7.72 | 7.63 | 7.67 |
| N ₄ | 14.44 | 14.26 | 14.35 | 7.78 | 7.67 | 7.72 |
| N ₅ | 14.57 | 14.35 | 14.46 | 7.83 | 7.70 | 7.77 |
| N ₆ | 14.67 | 14.47 | 14.57 | 7.89 | 7.76 | 7.83 |
| N ₇ | 14.76 | 14.58 | 14.67 | 7.95 | 7.81 | 7.88 |
| N ₈ | 14.89 | 14.77 | 14.83 | 8.00 | 7.90 | 7.95 |
| CD_{P=0.05} | 0.13 | 0.11 | 0.08 | 0.07 | 0.07 | 0.05 |

Comment [VM11]: Levels of significance indicating statistical significance (such as a, b, c, d) must be included in the table.

3.4. Reducing sugar (%)

The data pertaining to reducing sugar content in fruits also followed the same trend as total sugar content. Table 2 ~~illustrated that reducing sugar content was significantly influenced by variety, level of pruning~~ illustrates that reducing sugar content was significantly influenced by ~~variety, level of pruning~~ and micronutrient treatments. Among varieties, V₂ showed significantly higher level of reducing sugar content (8.08%) compared to V₁ (7.37%). Significant difference as a result of variety was also obtained by [8]. Pruning level P₂ (7.99%) performed better than P₁ (7.45%) in terms of reducing sugar content. More synthesis of photosynthates in moderately pruned apple trees as illustrated by [16] might be the probable cause of significant variation achieved due to pruning. As in case of total sugar content, N₈ exhibited significantly higher content of reducing sugar (7.95%) compared to all other levels. Again, N₀ exhibited the lowest content reducing sugar (7.50%). Besides aiding in sugar transport, source to sink movement of growth regulators is also regulated by boron [17]. Conversion of starch to sugar through hydrolysis is promoted by zinc. As a result, significantly higher amount of reducing sugar in zinc treated fruits were obtained [18].

3.5. Non-reducing sugar (%)

Table 3 depicted that all the factors ~~i.e. variety, pruning level and micronutrient sprays had significant effect on~~, ~~i.e., variety, pruning level, and micronutrient sprays, had a significant effect on the~~ non-reducing sugar content of Thailand ber fruit. The variety V₂ (7.07%) was significantly superior to V₁ (6.17%) in terms of non-reducing sugar content. This variation occurred because of significant diversity in total sugar and reducing sugar content among varieties. In case of pruning, P₂ (6.92%) performed significantly better than P₁ (6.31%) in regards to non-reducing sugar content of fruits. More assimilation of food material increased the non-reducing sugar content in medium pruned ber trees [9]. Similar observations in ber were also reported earlier [19]. Non-reducing sugar content of N₈ was significantly higher (6.88%) compared to all other micronutrient treatments, while, N₀ exhibited the lowest content non reducing sugar (6.36%). As a result of significant impact of boron and zinc on total sugar and reducing sugar, significant effect of these micronutrients were also noticed in case of non-reducing sugar content.

Table 3. Non reducing sugar and titratable acidity content as influenced by variety, pruning height and micronutrient sprays

| Treatments | Non reducing sugar (%) | | | Titratable acidity (%) | | |
|----------------------------|------------------------|--------------|--------------|------------------------|--------------|--------------|
| | 2020 – 21 | 2021 – 22 | Pooled | 2020 – 21 | 2021 – 22 | Pooled |
| V ₁ | 6.17 | 6.18 | 6.17 | 0.32 | 0.31 | 0.32 |
| V ₂ | 7.13 | 7.00 | 7.07 | 0.53 | 0.53 | 0.53 |
| CD_{P=0.05} | 0.040 | 0.031 | 0.023 | 0.008 | 0.008 | 0.006 |
| P ₁ | 6.33 | 6.30 | 6.31 | 0.49 | 0.48 | 0.48 |
| P ₂ | 6.96 | 6.88 | 6.92 | 0.37 | 0.36 | 0.36 |
| CD_{P=0.05} | 0.040 | 0.031 | 0.023 | 0.008 | 0.008 | 0.006 |
| N ₀ | 6.39 | 6.33 | 6.36 | 0.48 | 0.48 | 0.48 |
| N ₁ | 6.45 | 6.41 | 6.43 | 0.47 | 0.46 | 0.46 |
| N ₂ | 6.52 | 6.47 | 6.49 | 0.45 | 0.44 | 0.45 |

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Comment [VM12]: Reducing, non-reducing and total sugars can be clubbed into a single table whereas TSS and Titratable acidity in another table for the convenience.

Comment [VM13]: Levels of significance indicating statistical significance (such as a, b, c, d) must be included in the table.

| | | | | | | |
|----------------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| N ₃ | 6.59 | 6.55 | 6.57 | 0.44 | 0.42 | 0.43 |
| N ₄ | 6.67 | 6.59 | 6.63 | 0.43 | 0.42 | 0.42 |
| N ₅ | 6.74 | 6.64 | 6.69 | 0.41 | 0.40 | 0.41 |
| N ₆ | 6.77 | 6.71 | 6.74 | 0.40 | 0.40 | 0.40 |
| N ₇ | 6.81 | 6.77 | 6.79 | 0.38 | 0.37 | 0.37 |
| N ₈ | 6.89 | 6.86 | 6.88 | 0.38 | 0.38 | 0.38 |
| CD_{P=0.05} | 0.086 | 0.065 | 0.049 | 0.016 | 0.017 | 0.012 |

3.6. Titratable acidity (%)

Significant variations were noted in ~~case of titratable acidity of fruits in regards to variety, pruning the titratable acidity of fruits regarding variety, pruning,~~ and micronutrient level (Table 3). Variety V₁ recorded significantly lower titratable acidity of 0.32 per cent, ~~while~~ compared with V₂ (0.53%). As found in the current study ~~Ghosh and Mathew (2002) also saw significant effect of variety on,~~ Ghosh and Mathew (2002) also saw a significant effect of variety on the titratable acidity of fruits. In ~~case of~~ pruning, P₂ exhibited the lowest titratable acidity of 0.36 ~~per cent~~ compared to P₁ (0.48%). Reduction in pruning intensity significantly reduced the titratable acidity in the present investigation, which ~~is in conformity~~ conforms with the illustration of [20], which could be attributed to either ~~conversion of acid to sugar or the~~ conversion of acid to sugar or the utilization of acids in the ~~process of metabolism~~ metabolism process. Among different ~~level of micronutrients,~~ N₇ recorded the lowest level of titratable acidity with value of 0.37 per cent, which was at par with 0.38 per cent levels of micronutrients. N₇ recorded the lowest level of titratable acidity with the value of 0.37 percent, which was at par with 0.38 percent recorded in N₈. Further, the highest titratable acidity was found in N₀ (0.48%). The probable cause for the phenomenon might be ~~because of conversion of acids to sugar due to reversal of glycolytic pathway or acids had the conversion of acids to sugar due to the reversal of the glycolytic pathway, or acids been~~ utilized in respiration as a substrate or both [21].

Comment [VM14]: Missing reference? Also, this does not match with the style of the journal. Cite as per the journal guidelines.

3.7. TSS/acid-titratable acidity ratio

The ~~ratio of total soluble solid and acid~~ TSS and titratable acidity ratio differed significantly ~~w.r.t. variety, pruning level and,~~ and micronutrient levels (Table 4). ~~Significantly~~ A significantly higher value of 45.33 was observed in V₁, while V₂ recorded a value of 29.07. ~~In regards to~~ Regarding pruning height, P₂ exhibited a significantly higher TSS/acid ratio of 44.07 than P₁ (30.33). The ratio of TSS/acid was the maximum in N₇ (43.04) closely followed by N₈ (42.94). The lowest TSS/acid ratio was found in N₀ (31.24) ~~closely followed,~~ followed by N₁ (32.32). Interaction studies revealed significant difference of TSS/acid ratio among various interaction (Table 4.). More TSS and less titratable acidity tends to higher TSS/acid ratio and vice versa. The probable reasons already discussed which led to variation in TSS and titratable acidity. The significant variation in TSS and titratable acidity led to significant changes in TSS/acid ratios.

Comment [VM15]: Use full form

Table 4. TSS/acid ratio and ash content as influenced by variety, pruning height and micronutrient sprays

| Treatments | TSS : acid | | | Ash (%) | | |
|----------------|------------|-----------|--------|-----------|-----------|--------|
| | 2020 – 21 | 2021 – 22 | Pooled | 2020 – 21 | 2021 – 22 | Pooled |
| V ₁ | 44.08 | 46.57 | 45.33 | 2.37 | 2.38 | 2.37 |

Comment [VM16]: Levels of significance indicating statistical significance (such as a, b, c, d) must be included in the table.

| | | | | | | |
|----------------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| V ₂ | 28.90 | 29.25 | 29.07 | 2.65 | 2.66 | 2.65 |
| CD_{P=0.05} | 0.861 | 1.016 | 0.655 | 0.015 | 0.013 | 0.010 |
| P ₁ | 30.02 | 30.63 | 30.33 | 2.35 | 2.39 | 2.37 |
| P ₂ | 42.96 | 45.19 | 44.07 | 2.66 | 2.65 | 2.66 |
| CD_{P=0.05} | 0.861 | 1.016 | 0.655 | 0.015 | 0.013 | 0.010 |
| N ₀ | 31.14 | 31.34 | 31.24 | 2.39 | 2.41 | 2.40 |
| N ₁ | 31.40 | 33.25 | 32.32 | 2.43 | 2.44 | 2.44 |
| N ₂ | 33.74 | 34.89 | 34.32 | 2.45 | 2.46 | 2.46 |
| N ₃ | 34.02 | 36.96 | 35.49 | 2.48 | 2.48 | 2.48 |
| N ₄ | 35.30 | 38.32 | 36.81 | 2.51 | 2.52 | 2.51 |
| N ₅ | 38.34 | 39.26 | 38.80 | 2.53 | 2.54 | 2.53 |
| N ₆ | 39.06 | 40.61 | 39.83 | 2.56 | 2.57 | 2.57 |
| N ₇ | 43.09 | 42.98 | 43.04 | 2.60 | 2.62 | 2.61 |
| N ₈ | 42.33 | 43.56 | 42.94 | 2.63 | 2.64 | 2.64 |
| CD_{P=0.05} | 1.827 | 2.155 | 1.389 | 0.031 | 0.028 | 0.020 |

3.8. Ash content (%)

A view of Table 4 enlightened that all the factors ~~i.e. variety, pruning height and micronutrient levels had significant influence on ash content of Thailand ber, i.e., variety, pruning height, and micronutrient levels, had significant influence on the ash content of Thailand's~~ fruits under study. The ash content of V₂ was 2.65 ~~per cent~~ percent, which was significantly superior to V₁ (2.37%). As observed in the current investigation, significant varietal differences among Thai apple ber accessions were also observed [8]. Significantly higher ash content was witnessed in P₂ (2.66%) compared to P₁ (2.37%). Moderate pruning provides ~~more open canopy of ber plants, which aids in penetration of more sun rays leading to more production of photosynthates [2], which in turn triggered a more open canopy of ber plants, which aids in the penetration of more sun rays, leading to more production of photosynthates [2], which in turn triggers~~ changes in ash content. The micronutrient treatment N₈ was found to be the best in ~~terms of~~ ash content and the value of 2.64 per cent was significantly higher than all other micronutrient treatments. The lowest ash content was ~~detected in~~ N₀ with a value of 2.40 per cent. Application of micronutrient boron enhances the uptake and transport of other nutrients, viz., N, P, K, Cu, Zn and Fe from soil [22] which instigated increase in ash content of fruits. Zinc influenced ash content positively being a constituent of the enzyme carbonic anhydrase, which is a mandatory enzyme for photosynthesis [23].

Formatted: Font: Not Italic

4. CONCLUSION

In the present study, it was revealed that Roundish ovate reddish type variety (V₂) was superior to the Round green type variety (V₁) in terms of physico-chemical parameters. In regards to those parameters pruning height of 75 cm (P₂) was better and micronutrient

sprays with Borax 0.5% and Zinc Sulphate 0.5% (N₈) performed the best. On the basis of results, the best pruning height and micronutrient sprays can be advocated for adoption.

Comment [VM17]: The conclusion is very short and can be extended further for more precise recommendations by justifying the hypothesis of the experiment.

REFERENCES

1. Anonymous Horticultural Statistics at a glance 2018: Horticultural Statistics Department, Govt. of India, New Delhi; 2019.
2. Kumar H, Katiyar PN, Singh AK, Rajkumar BV. Effect of different pruning severity on growth and yield of ber (*Zizyphus mauritiana* Lamk), cv. Banarsi Karaka. Int J CurrMicrobiolAppld Sci. 2014;3(5):935-940.
3. Singh S, Pratap B, Yadav A, Shivam. Assess the effect of pruning and plant growth regulators on growth, flowering and fruiting of ber tree. J Pharmacognosy Phytochem. 2017;6(5):735-738.
4. Peres LA, Reyes RD. Effect of nitrogen, boron and lime on *Carica papaya*. J AgrilUniv Puerto Rico. 1983;67:181-187.
5. Saadati S, Moallemi N, Mortazavi SMH, Seyyednejad SM. Foliar applications of zinc and boron on fruit set and some fruit quality of olive. Vegetos. 2016;29(2):100-120.
6. AOAC. *Official methods of analysis* (10th edn.): Association of Official Agricultural Chemist, Washington, D. C; 1965.
7. Panse VG, Sukhatme PV. *Statistical Methods for Agriculture Workers*: Indian Council of Agricultural Research, New Delhi; 1985.
8. Khanikar H, Langthasa S, Hazarika DN, Goswami RK, Deori GD. Studies on Physico-chemical characters of Thai Apple ber (*Zizyphus mauritiana* Lamk.) grown in Assam. J Pharmacognosy Phytochem. 2021;10(3):187-192.
9. Kumar A, Pratap B, Vishwakarma G. Effect of pruning on growth, yield and fruit quality characteristics of ber (*Zizyphus mauritiana* Lamk.) cv. Gola. Progressive Hort. 2016;48(2):148-153.
10. Kumar M, Rawat V, Rawat JMS, Tomar YK. Effect of pruning intensity on peach yield and fruit quality. Sci Hortic. 2010;125(3):218-221.
11. Ghosh SN, Mathew B. Performance of nine ber cultivars (*Zizyphus mauritiana* Lamk) on top working in semi-arid region of West Bengal. J Appld Hort. 2002;4(1) :49-51.
12. Tadayon MS, Moafpourian G. Effects of exogenous epi-brassinolid, zinc and boron foliar nutrition on fruit development and ripening of grape (*Vitis vinifera* L. cv. 'Khalili'). Sci Hortic. 2019;244:94-101.
13. Singh S, Pratap B, Gupta S, Yadav D, Kumar A, Debapriya S, et al. Assess the effect of pruning and plant growth regulators on yield and quality of ber fruit. Int J CurrMicrobiolAppld Sci. 2019;8(1):539-547.

14. Gauch HG, Duggar WM. The role of boron in the translocation of sucrose. *Plant Physiol.*1953;28:457-466.
15. Alloway BJ. Fundamental aspects of zinc in soil and plants. In: *Zinc in Soils and crop nutrition*. 2ndEdn., IZA and IFA, Brussels, Belgium and Paris, France, pp. 30-52; 2008.
16. Ferre DC, Myers SC, Rom CR, Taylor BH. Physiological aspects of summer pruning. *Acta Hort.*1984;146:243-252.
17. Shireen F, Nawaz MA., Chen C, Zhang Q, Zheng Z, Sohail H, et al. Boron: Functions and approaches to enhance its availability in plants for sustainable agriculture. *Int J Mol Sci.* 2018;19:95-98.
18. Jat G, Kacha HL. Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *J AgriSearch.* 2014;1(2):86-91.
19. Yadav AK, Singh SP Singh A. Screening of ber (*Zizyphus mauritiana* Lamk.) cultivars based on external, physical and chemical attributes for cultivation in eastern Uttar Pradesh. *Indian Horti J.*2005;18(3):144-148.
20. Sahu CK, Patel MK, Panda CM. Effect of pruning and plant growth regulator on physico-chemical quality of sapota (*Manilkara zapota* L.) cv. Cricket Ball. *Pharma Innov J.*2018;7(8):335-338.
21. Kunda S, Mitra SK. Response of guava to foliar spray of copper, boron and zinc. *Indian Agri.*1999;43(1-2):49-54.
22. Ahmed N, Abidm M, Abmadm F, Ullahm MA, Javaidm Q, Ali MA. Impact of boron fertilization on dry matter production and mineral constitution of irrigated cotton. *Pakistan J Bot.* 2011;43:2903-2910.
23. Bisen B, Tiwari JP, Mishra KK, Lal S. Effect of zinc sulphate on vegetative growth, yield and leaf nutrient status of guava (*Psidium guajava* L.). *Int J CurrMicrobiolAppl Sci.*2020;9(7):1265-1273.