

Influence of organic nutrient sources on nutrient, quality and yield parameters of vegetable clusterbean

Abstract:

The investigation “organic nutrient source and their influence on nutrient dynamics, quality and yield parameters of vegetable clusterbean” during the year 2018, was conducted at irrigated wetland farms, TNAU, Coimbatore. The RBD (randomized complete block design) design was adopted and replicated thrice. The total number of treatments were 12 with various basal applications of organic manures namely farm yard manure, vermicompost, humic acid, ghanjeevamrutham and also foliar spray combinations namely panchagavya and banana pseudo stem sap. The standard recommended cultural practices of organic farming were followed for all treatments except fertilizer application and synthetic chemical use. The data revealed that soil chemical properties viz., pH, EC and organic carbon were not significantly influenced by various organic nutrient sources. Significantly, higher values of N, P and K availability in soil and their uptake by clusterbean crop and crude protein content, crude fiber, ascorbic acid, moisture percentage in cluster bean pods were recorded with 100 % recommended dose of fertilizer and 2% foliar spray of TNAU pulse wonder. However, organic treatments were on par with inorganic treatments for all the nutrient, quality and yield parameters. So, organic treatments like organic farmers practice and FYM along with Panchagavya were recommended to improve nutrient use efficiency and quality of yield, eventually to benefit farmers and complete agricultural ecosystem in terms of profit and sustainability, respectively.

Key words: Clusterbean, FYM, N uptake, Organic manures, Panchagavya,

Comment [JR1]: Enter the objective of the study.
Suggestion:
the experiment was taken up to study the influence of various basal organic manures and foliar spray on soil nutrient availability and their uptake and also on quality parameters of vegetable cluster bean.

Introduction:

The clusterbean (*Cyamopsis tetragoloba*(L). Taub.) is an annual nitrogen fixing, drought tolerant legumecrop with deep root system. It is also valued as potential industrial crops, because of Gaur (galactomanan) gum content in endosperm of seed. Inorganic fertilizers has resulted in reduced crop yield, as a result of soil acidity, nutrient imbalance, poor physical properties of the soil and nutrient retention(Ballal and Kadam, 2016). Organic farming has gained importance in recent years, as a result of the of intrinsic benefits, such as lowering the hazards of human, animal and environmental exposure to toxic chemicals. It also helps in recycling animal wastes back into the farm and therefore reducing the cost of cultivation. Therefore, the application of plant nutrients through organic sources like FYM, Vermicopmost, *Panchagavya*, bananapseudostem sap remains a preferable option for ensuring increased fruit shelf life, quality and sustainable crop production (Alexander et al., 2009). Considering the benefits of organic farming, the experiment was taken up to study the influence of various basal organic manures and foliar spray on soil nutrient availability and their uptake and also on quality parameters of vegetable clusterbean.

Material and Methods:

The field experiment was taken up at irrigated wetland farms of Tamil Nadu Agricultural University, Coimbatore during *Kharif* season from July to October, 2018. The experiment was carried out in randomized complete block design with twelve treatments and three replications. Treatment details are as follows ; T₁: Organic farmer's practice (farm yard manure (FYM) @10 t/ha + *Jeevamruth* @ 500 l/ha along with irrigation on 3rd days after sowing (DAS), 30 DAS and 60 DAS + *Panchagavya* @ 3% as foliar spray (FS) on 30, 45 and 60 days after sowing (DAS), T₂: FYM @ 25 t/ha + *Panchagavya* @ 3% as FS on 30, 45 and 60 DAS, T₃: FYM @ 12.5 t/ha + *Panchagavya* @ 3% as FS on 30, 45 and 60 DAS, T₄: FYM @ 12.5 t/ha + banana pseudostem sap @ 2% as FS on 30, 45 and 60 DAS, T₅:

Comment [JR2]:

In relation to the introduction:

1. Review all citations in the introduction.

Author Guidelines

All references must be numbered consecutively and citations of references in text should be identified using numbers in square brackets (e.g., "as discussed by Smith [9]"; "as discussed elsewhere [9, 10]"

2. Insert more and recent citations

Comment [JR3]: In relation to the Material and Methods

1. Review all citations in the according

Author Guidelines

2. Redraw all tables. in the according

Author Guidelines

Vermicompost @ 5 t/ha + Panchagavya @ 3% as FS on 30, 45 and 60 DAS, T₆: Vermicompost @ 5t/ha + banana pseudostem sap @ 2% as FS on 30, 45 and 60 DAS, T₇: *Ganajeevamruth* @ 500 kg/ha + *Panchagavya* @ 3% as FS on 30, 45 and 60 DAS, T₈: *Ganajeevamruth* @ 500 kg/ha + banana pseudostem sap @ 2% as FS on 30, 45 and 60 DAS, T₉: Humic acid @ 6 l/ha + *Panchagavya* @ 3% as FS on 30, 45 and 60 DAS, T₁₀: Humic acid @ 6 l/ha + banana pseudostem sap @ 2% as FS on 30, 45 and 60 DAS, T₁₁: RDF of NPK fertilizers (50:50:25) + TNAU pulse wonder @ 1% as FS, T₁₂: Absolute control.

A well decomposed farm yard manure, vermicompost, *ganajeevamruth* and humic acid were administered on to clusterbean crop as basal treatments, whereas *Panchagavya*, banana pseudostem sap were foliar sprayed on 30, 45, and 60 DAS and Jeevamruth was dissolved into irrigation water on 3rd day after sowing, 30 DAS and 60 DAS. The banana pseudostem sap was collected from 8 year old banana plant, by making sharp cut on one side of a stem. Two milliliter of the extracted sap was diluted in one liter of water and applied as foliar spray.

Field was ploughed twice using a cultivator and once using rotovalor to get fine tilth. A bullock-drawn ridger was used to create ridges 45 centimetres apart. The dimension of the gross plot was 6.75 m × 3.15m. The field was irrigated and basal organic manures were applied prior to seeding. The experiment employed the clusterbean MDU 1 variety released from TNAU. Rhizobium was applied to the seeds at a rate of 80 g/kg, and they were then shade-dried for 12 hours. The ridge was seeded with a 15-cm seed spacing on one side. In accordance with the soil's moisture content, irrigation was applied on the day of sowing, on the 3DAS, and subsequently as and when moisture was deficit. In accordance with treatment schedule, foliar sprays of organic liquid manures were applied at 30, 45, and 60 DAS. Except for the use of synthetic chemicals, all other traditional methods were carried out in accordance with the TNAU recommendations package of practices.

The initial soil samples were collected from the experimental field in in zig zag manner at 0 to 30 cm depth before treatment imposition. Then composite samples were prepared and analyzed for physical and chemical properties. The results were presented in the Table 1.

Table 1. Initial Soil and irrigation water characteristics of the experimental field.

Comment [JR4]: Format table 1 according to General Guideline for Authors

| Particulars | Values | Methods | Reference |
|--|-----------|--|-----------------------------|
| A. Textural composition (Moisture free basis) | | | |
| Mechanical Analysis | | | |
| Clay (%) | 47.6 | Robinson International pipette method | Piper (1966) |
| Silt (%) | 22.5 | | |
| Fine sand (%) | 11.7 | | |
| Coarse sand (%) | 18.2 | | |
| Textural class | Clay loam | USDA, Triangle | Piper (1950) |
| II. Chemical characteristics | | | |
| pH | 8.27 | Potentiometric pH meter | Jackson (1973) |
| Electrical conductivity (dS/m) | 0.25 | EC meter | Jackson (1973) |
| Organic carbon (%) | 0.45 | Walkley and Black rapid titration method | Jackson (1973) |
| Available N (kg/ha) | 218 | Alkaline permanganate method | Subbiah and Asija (1956) |
| Available P ₂ O ₅ (kg/ha) | 18.2 | Olsen's method | Olsen (1954) |
| Available K ₂ O (kg/ha) | 402 | Flame photometer method | Stanford and English (1949) |
| B. Irrigation water characteristics | | | |
| pH | 7.5 | Potentiometric pH meter | Jackson (1973) |
| Electrical conductivity (dS/m) | 1.2 | Electrical conductivity bridge CM -180 | Jackson (1973) |

Soil samples were collected from 0 to 30 cm depth at different places on 0, 30, 45, 60 DAS and at harvest stage of the crop in each plot. The composite soil samples were collected, mixed and kept for shade drying. A representative soil sample was made (1 kg) by quartering method and packed into labelled polythene bags. This processed sample is used for analysis of soil chemical properties namely soil pH, EC, OC, available nitrogen, phosphorous, potassium using the methods mentioned in Table 1.

Five plants samples were taken randomly from sampling row on each plot for destructive sampling during 30, 45, 60 DAS and 90 DAS. Shade dried for three days, then oven dried for one day at 65⁰C for obstinate dry weight, then powdered and utilized for the analysis of nutrients and method of analysis followed as mentioned in Table 2.

Table 2. Methods adopted for plant analysis

| Element | Method | Reference |
|------------|---|---------------------------|
| Nitrogen | Micro Kjeldahl method | Humphries (1956) |
| Phosphorus | Vanadomolybdo phosphoric yellow colour method | Piper (1966) |
| Potassium | Flame photometer method | Muhr <i>et al.</i> (1965) |

Comment [JR5]: Format table 2 according to General Guideline for Authors

Nutrient uptake was computed based on dry weight basis of aerial biomass at harvest stage using the following formulae.

Uptake of nutrient (kg/ha) = $\frac{\text{Nutrient content (\%)} \times \text{Dry matter production (kg/ha)}}{100}$

100

Quality parameters such as crude fibre content in green pods was determined by the method suggested by Association of Official Analytical chemists Inc., USA. (AOAC, 1975). Two grams of pod sample (W_1) were grounded then refluxed with 1.25% H_2SO_4 and then washed, again refluxed with 1.25% NaOH for 30 minutes. Later, sample was dried out, to take initial weight (W_2). Ignite the sample in a muffle furnace and again had taken weight (W_3). Loss in weight was considered as crude fibre content and expressed on the basis of using following relationship.

$$\text{Crude fibre (\%)} = \frac{(W_2 - W_3)}{W_1} \times 100$$

Where, W_1 = Initial weight of sample, W_2 = Weight of refluxed sample and W_3 = Weight of ignited sample.

Protein percentage was enumerated by assessing the total nitrogen using Kjeldahl method in pod (Humphries, 1956) and then nitrogen percentage is multiplied a factor 6.25. Ascorbic acid content was measured with the help of procedure given by Horwitz *et al.* (1970) and expressed in mg/100 grams of fresh sample. Moisture content of pods measured in percentage form

selected 10 pods from each treatment. Initial weight is noted using standard balance, then dried in oven at 65°C for 3 days for complete moisture loss. Thereafter final dry weights were noted and substituted in the following formula.

$$\text{Moisture content (\%)} = [\text{Fresh weight (g)} - \text{Dry weight (g)} / \text{Fresh weight (g)}] \times 100$$

The experimental data was statistically analyzed by analysis of variance using Agres software. The level of significance was $P=0.05$.

Results and Discussion:

The mean statistically analyzed data of soil pH, EC and organic carbon on 30, 45, 60 DAS and at harvest stages and moisture percentage in cluster bean pods were not changed significantly with application of various organic manures. Whereas, significantly higher soil available nitrogen (260.8 kg/ha, 254.0 kg/ha and 250.6 kg/ha on 30, 45 and 60 DAS, respectively), available phosphorus (23.4 kg/ha, 21.8 kg/ha, 21.3 kg/ha and 16.4 kg/ha on 30 DAS, 45 DAS, 60 DAS and 90 DAS, respectively) and available potassium (424.5 kg/ha, 413.5 kg/ha and 406.5 kg/ha on 30, 45, 60 DAS and 90 DAS, respectively) and also higher uptake of nitrogen (41.4 kg/ha), phosphorous (7.1 kg/ha) potassium (23.5 kg/ha) nutrients by cluster bean at harvest stage were noted with RDF + TNAU pulse wonder (T_{11}), which were statistically on par with organic farmers' practice (T_1) and FYM @ 25 t/ha in addition to foliar spray of 3% *Panchagavya* (T_2). Whereas, at 90 DAS available nitrogen (238.6 kg/ha) was found higher with organic farmers' practice (T_1). But, it was on parity with T_{11} and T_2 (Table 3.). This may be owing to rapid and direct availability of available nutrients from urea, single super phosphate and muriate of potassium. Whereas, with application of organic treatments like addition FYM and *Jeevamrutham* had resulted in higher amount organic manure, microbial population, organic acids and enzymes like dehydrogenase, urease and phosphatase in soil helps in conversion of unavailable form of nutrients to available form of nutrients for better plant uptake. *Jeevamruth* has large amount of free living N fixers,

Comment [JR6]: In relation to the results and discussion
3. Review all citations in the according
AuthorGuidelines
4. Redraw all tables.
in the according
AuthorGuidelines

symbiotic N fixers and associated N fixer and phosphorous solubilizers (Devakumar *et al.*, 2008) that might have led to more number of nodules, higher amount of nitrogen fixation and phosphorous solubilization (Yadav and Chhipa, 2007). All this collective action might have resulted into higher availability of nitrogen, phosphorous and potassium in soil. Naga (2013), (2014), Manohar *et al.* (2018a), Prabhavathi (2014) and Lodha (2016a). Lower available nitrogen, phosphorus and potassium and lower uptake of these nutrients by cluster bean plants was observed under absolute control (T₁₂).

The application of farm yard manure improves soil quality by altering bulk density, enhancing aeration, enhancing microbial life in the soil, and increasing plant nutrient availability. During microbial decomposition of FYM in organic treatments, microbes feed on FYM and release organic acids and organic collides with greater cation exchange sites (Yadav *et al.*, 2019), which results in faster mineralization of organic waste in the soil. Eventually, the availability of nitrogen increase (Yadav and Chhipa, 2007) along with increase solubility of native phosphates and reduction of Potassium fixation (Bharadwaj and Omanwar, 1994). The added organic collides with greater cation exchange sites improved available potassium (Yadav *et al.*, 2019) by holding applied K and released K from non-exchangeable K. (CHOGATAPUR, 2015; LODHA, 2016b; Prabhavathi, 2014)

Statistically more crude fibre content in pod (4.7%) and protein percentage (4.1%) in vegetable pod of cluster bean was obtained in RDF in addition to TNAU pulse wonder (T₁₁), but it maintained statistically parity with organic farmers' practice (T₁) and FYM @ 25 t/ha in addition to foliar spray of Panchagavya @ 3 % (T₂). As mentioned earlier higher rapid availability of nutrients from soil and foliar application and better micro and macro nutrient uptake might have promoted higher synthesis of protein and crude fibre. Reddy (2011), Prabhavathi (2014) and Choudhary (2015).

Ascorbic acid represents vitamin C content, which is very essential for human. Significantly higher ascorbic acid (52.7 mg/100g) was noted with organic farmer's practice (T1), which was statistically on par with FYM @ 25 t/ha in addition to foliar spray of 3% *Panchagavya* (T2) and RDF addition of TNAU pulse wonder (T₁₁). The reason behind higher accumulation of ascorbic acid was due to better uptake of nutrient especially K (quality element). Potassium has close relationship with metabolism of carbohydrate and ascorbic acid synthesis Boraiah *et al.* (2017); (El-Sayed and El Sayed, 2013) and (Vanitha *et al.*, 2014). Lower crude fibre (3%), protein percentage (2.6%) and lowest ascorbic acid (33.8 mg/100g) was recorded under absolute control (T₁₂) than others. Moisture percentage in pods was statistically analyzed had not shown any significant difference among the treatments with application of organic manures.

The higher significant yield (11084 kg/ha) was attained in RDF + TNAU pulse wonder (T₁₁) and followed by organic farmers' practice (T1) and FYM @ 25 t/ha + foliar spray of 3% *Panchagavya* (T2) as statistically on par treatments. Whereas, significant lowest yield (7102 kg/ha) was registered in absolute control (T₁₂). Compared to the absolute control, the yield of the RDF with TNAU pulse wonder was 56% higher. However, compared to the absolute control (T₁₂), organic farmers' practices (T1) and FYM @ 25 t/ha, along with a foliar spray of 3% *Panchagavya* (T2), produced yields that were 43% and 42% higher, respectively.

Vegetable clusterbeans gained more nutrients from the soil as a composite consequence. Panchagavya, Jeevamrutham and FYM might had enhanced plant vegetative growth with sustainable release of macro and micro nutrients. Eventually, higher photosynthate produced may get partitioned and translocated into pods for enhancing yield. The composite effect of all above discussed had made organic treatments statically on par with inorganic treatment. (Manohar et al., 2018b; RAJPOOT, 2006).

The reason behind higher yield parameters in inorganic treatment was due to quick available form of nutrient supplied to the plant through soil and foliar application enhanced growth parameters like plant height, number of leaves, leaf area/plant along with higher chlorophyll content resulted into synthesis more quantity of photosynthates. The synthesized food material then might have transferred from source (leaves and stems) to sink successfully (pods) with minimum losses. Thus, resulted in increased yield parameters. (Prabhavathi, 2014)

Conclusion:

Although, inorganic treatment as a nutrient source had shown higher NPK nutrient availability in soil, higher NPK nutrient uptake in plant and higher crude fibre content, protein percentage in pods and higher yield was observed. The organic farmers' practice of FYM, *jeevamrutham* and foliar spray of *panchagavya* was recommended for better fruit quality and sustainable clusterbean production by considering the negative effects caused by chemical farming to environment and life on the earth,

Table 3. Effect of organic manures on pod quality parameter, soil available nitrogen, phosphorous, potassium and their uptake by clusterbean. Comment [JR7]:

| Treatment | Available nitrogen (kg/ha) | | | | Available phosphorus (kg/ha) | | | | Available potassium (kg/ha) | | | | Plant uptake | | | Quality parameters | | | Pod Yield (kg/ha) |
|-------------|----------------------------|--------|--------|--------|------------------------------|--------|--------|--------|-----------------------------|--------|--------|--------|------------------|------------------|------------------|--------------------|-----------------|-------------------------|-------------------|
| | 30 DAS | 45 DAS | 60 DAS | 90 DAS | 30 DAS | 45 DAS | 60 DAS | 90 DAS | 30 DAS | 45 DAS | 60 DAS | 90 DAS | N uptake (kg/ha) | P uptake (kg/ha) | K uptake (kg/ha) | Crude protein (%) | Crude Fibre (%) | Ascorbic acid (mg/100g) | |
| T1 | 253.1 | 250.6 | 246.7 | 238.6 | 21.6 | 20.0 | 19.6 | 15.1 | 419.9 | 411.0 | 404.3 | 394.5 | 37.0 | 6.3 | 23.5 | 3.8 | 4.3 | 52.7 | 10215 |
| T2 | 256.9 | 249.9 | 243.1 | 236.6 | 21.4 | 19.9 | 19.5 | 15.0 | 417.0 | 406.0 | 399.0 | 390.0 | 39.3 | 6.7 | 22.4 | 3.7 | 4.3 | 48.6 | 10145 |
| T3 | 234.0 | 227.1 | 224.9 | 214.9 | 19.4 | 18.0 | 17.7 | 13.6 | 386.5 | 371.0 | 375.5 | 354.5 | 31.6 | 5.4 | 19.3 | 3.4 | 3.9 | 43.7 | 9190 |
| T4 | 232.6 | 228.9 | 223.1 | 215.4 | 19.6 | 18.2 | 17.9 | 13.7 | 384.8 | 368.7 | 373.8 | 355.0 | 34.5 | 5.9 | 20.9 | 3.4 | 3.9 | 44.2 | 9283 |
| T5 | 234.0 | 227.5 | 223.5 | 213.7 | 19.3 | 17.9 | 17.5 | 13.5 | 382.0 | 368.5 | 371.0 | 352.7 | 31.8 | 5.4 | 19.1 | 3.4 | 3.8 | 43.4 | 9125 |
| T6 | 234.3 | 228.7 | 224.1 | 214.2 | 19.6 | 18.2 | 17.8 | 13.7 | 379.7 | 367.6 | 368.7 | 353.6 | 34.7 | 5.9 | 17.9 | 3.4 | 3.9 | 44.0 | 9256 |
| T7 | 225.7 | 224.0 | 218.4 | 209.4 | 17.0 | 16.8 | 15.5 | 11.9 | 367.6 | 356.6 | 356.6 | 340.6 | 24.1 | 4.1 | 20.2 | 3.0 | 3.4 | 38.3 | 8042 |
| T8 | 227.8 | 223.5 | 219.1 | 210.1 | 17.2 | 15.9 | 15.6 | 12.0 | 365.5 | 354.5 | 354.5 | 339.5 | 25.9 | 4.4 | 21.3 | 3.0 | 3.4 | 38.6 | 8115 |
| T9 | 229.8 | 224.1 | 221.6 | 212.5 | 17.3 | 16.0 | 15.7 | 12.1 | 370.5 | 359.5 | 359.5 | 343.5 | 24.7 | 4.2 | 20.1 | 3.0 | 3.4 | 38.8 | 8164 |
| T10 | 230.4 | 226.9 | 223.1 | 214.1 | 19.1 | 17.8 | 17.4 | 13.4 | 371.1 | 359.1 | 358.1 | 344.1 | 29.8 | 5.1 | 20.9 | 3.3 | 3.8 | 43.1 | 9055 |
| T11 | 260.8 | 254.0 | 250.6 | 220.2 | 23.4 | 21.8 | 21.3 | 16.4 | 424.5 | 413.5 | 406.5 | 387.0 | 41.4 | 7.1 | 21.0 | 4.1 | 4.7 | 48.3 | 11084 |
| T12 | 198.3 | 194.8 | 189.1 | 182.6 | 15.0 | 14.9 | 13.7 | 10.5 | 327.0 | 316.0 | 311.0 | 301.0 | 21.0 | 3.6 | 17.4 | 2.6 | 3.0 | 33.8 | 7102 |
| S. Ed | 12.0 | 11.8 | 11.1 | 12.0 | 0.9 | 1.1 | 0.9 | 0.7 | 19.2 | 19.9 | 18.8 | 17.6 | 1.6 | 0.3 | 1.1 | 0.2 | 0.2 | 2.3 | 498 |
| CD (P=0.05) | 24.9 | 24.5 | 23.1 | 25.0 | 2.0 | 2.2 | 1.9 | 1.5 | 39.7 | 41.2 | 38.9 | 36.5 | 3.4 | 0.6 | 2.3 | 0.4 | 0.4 | 4.7 | 1033 |

Reference:

- Alexander, D., Rajan, S., Rajamony, L., Ushakumari, K. and Kurien, S., 2009. The adhoc Package of Practices recommendations for organic farming. ORGANIC FARMING.
- AOAC, 1975. Official methods of analysis, 222. Association of Official Analytical Chemists Washington, DC.
- Ballal, A. and Kadam, A., 2016. Okra crop growth and yield responses to different organic sources of nitrogen. International Journal of Agricultural Science, 8(48): 2042-2044.
- Bharadwaj, V. and Omanwar, P., 1994. Long term effects of continuous rotational cropping and fertilization on crop yields and soil properties-II. Effects on EC, pH, organic matter and available nutrients of soil. Journal of the Indian Society of Soil Science, 42(3): 387-392.
- Boraiah, B., Devakumar, N., Shubha, S. and Palanna, K., 2017. Effect of Panchagavya, Jeevamrutha and cow urine on beneficial microorganisms and yield of capsicum (*Capsicum annum* L. var. grossum). Int. J. Curr. Microbiol. App. Sci, 6(9): 3226-3234.
- CHOGATAPUR, S.V., 2015. Response of Clusterbean (*Cyamopsis tetragonoloba* (L.) Taub.) Genotypes (Gum) to Plant Density And Bioinoculants, University of Agricultural Sciences, Dharwad.
- Choudhary, R.S., 2015. Effect of Organic Manures and Mineral Nutrients on Soil Properties and Yield of Clusterbean [*Cyamopsis tetragonoloba* (L.) Taub.]. Sri Karan Narendra Agriculture University. .
- Devakumar, N. et al., 2008. Activities of Organic Farming Research Centre, University of Agricultural Sciences, Bengaluru, Karnataka, India.
- El-Sayed, H. and El Sayed, A., 2013. Exogenous application of ascorbic acid for improve germination, growth, water relations, organic and inorganic components in tomato (*Lycopersicon esculentum* Mill.) plant under salt-stress. New York Sci. J, 6(10): 123-139.
- Horwitz, W., Chichilo, P. and Reynolds, H., 1970. Official methods of analysis of the Association of Official Analytical Chemists. Official methods of analysis of the Association of Official Analytical Chemists.
- Humphries, E., 1956. Mineral components and ash analysis, Moderne Methoden der Pflanzenanalyse/Modern Methods of Plant Analysis. Springer, pp. 468-502.
- Jackson, M., 1973. Soil chemical analysis, pentice hall of India Pvt. Ltd., New Delhi, India, 498: 151-154.
- Lodha, b., 2016a. Maximization of Resource Use Efficiency for Higher Yield of Clusterbean [*Cyamopsis tetragonoloba* (L.) Taub] College of Agriculture, Gwalior.
- LODHA, B., 2016b. Maximization of Resource Use Efficiency for Higher Yield of Clusterbean [*Cyamopsis tetragonoloba* (L.) Taub], RVSKVV, Gwalior (MP).
- Manohar, C.V.S., Sharma, O. and Verma, H., 2018a. Nutrient status and yield of clusterbean [*Cyamopsis tetragonoloba* (L.) Taub] as influenced by fertility levels and liquid biofertilizers. J. Pharmacogn. Phytochem, 7: 1840-1843.
- Manohar, C.V.S., Sharma, O. and Verma, H., 2018b. Nutrient status and yield of clusterbean [*Cyamopsis tetragonoloba* (L.) Taub] as influenced by fertility levels and liquid biofertilizers. Journal of pharmacognosy and phytochemistry, 7(5): 1840-1843.
- Muhr, G.R. et al., 1965. Soil testing in India United states Agency for International Development Mission to India. New Delhi.
- Naga, S.D., 2013. Effect of integrated nutrient Management in clusterbean [*Cyamopsis Tetragonoloba* (L.) Taub.], Swami Keshwanand Rajasthan Agricultural University.
- Olsen, S.R., 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture.
- Piper, C., 1950. Soil and plant analysis. Adelaide University. Hassel Press, Australia. 368p.
- Piper, C., 1966. Mechanical analysis of soil by International Robinson's Pipette method. Soil and plant analysis.

Comment [JR8]: Review all references

Correct form

According Author Guidelines

Asiedu EA, Powell AA, Stuchbury T. Cowpea seed coat chemical analysis in relation to storage seed quality. *African Crop Science Journal*. 2000, 8: 283-294.

- Prabhavathi, M., 2014. INTEGRATED NUTRIENT MANAGEMENT IN CLUSTERBEAN (*Cyamopsis tetragonoloba* L. Taubert), ACHARYA NG RANGA AGRICULTURAL UNIVERSITY, RAJENDRANAGAR, HYDERABAD.
- Prabhavathi, M., 2014 Integrated Nutrient Management in Clusterbean (*Cyamopsis tetragonoloba* (L.) Taubert), Acharya NG Ranga Agricultural University, college of agriculture, Rajendranagar, Hyderabad.
- RAJPOOT, J.S., 2006. Effect of organic manure and PSB on the productivity of clusterbean.
- Reddy, N., 2011. Studies on effect of inm practices on growth, yield and quality of cluster bean [*Cyamopsis tetragonoloba*(L.)Tuab.]. Andhra Pradesh Horticultural University.
- Stanford, G. and English, L., 1949. Use of the flame photometer in rapid soil tests for K and Ca.
- Subbiah, B. and Asija, G., 1956. Alkaline permanganate method of available nitrogen determination. *Current Science*, 25: 259-260.
- Vanitha, M., Radha, S. and Vijayakumari, B., 2014. Herbals and biofertilizer as a nutrient supplement for improving biochemical parameters of cluster bean (*Cyamopsis tetragonoloba* L. var. PNB). *Journal of Natural Product and Plant Resources*, 4(1): 29-33.
- Yadav, K. and Chhipa, B., 2007. Effect of FYM, gypsum and iron pyrites on fertility status of soil and yield of wheat irrigated with high RSC water. *Journal of the Indian Society of Soil Science*, 55(3): 324-329.
- Yadav, S.K., Benbi, D.K. and Toor, A.S., 2019. Effect of long-term application of rice straw, farmyard manure and inorganic fertilizer on potassium dynamics in soil. *Archives of Agronomy and Soil Science*, 65(3): 374-384.

UNDER PEER REVIEW