

Effects of Native Rhizobia on Soybean [*Glycine max* (L.) Merrill] Production and Soil Properties in Daloa, Center-West of Côte d'Ivoire

ABSTRACT

The field experiment study was conducted at the experimental station of Jean Lorougnon Guédé University in Daloa to evaluate the potential effects of native rhizobia strains on soybean production and soil properties. The experiment was layout in a randomized complete block design with three replications. The treatments include seven native strains, one exotic strain (*Bradyrhizobium japonicum* IRAT FA3), one synthetic fertilizer NPK 12-22-22 formulation and one negative control (no fertilizer, no inoculation). Agronomic parameters (plant height, plant dry biomass, nodule number plant⁻¹, pod number plant⁻¹, pod weight plant⁻¹, seed yield ha⁻¹) and soil parameters (pH, total nitrogen, organic carbon, organic matter and available phosphorus) were measured. Results showed that all tested native rhizobia enhanced nodules number, plant growth and soybean yield as compared to negative control. Among these native rhizobia, RSC119 and RSC504 induced more nodules than the introduced strain *B. japonicum* IRAT FA3. RSC119, RSC309 and RSC508 produced more pods and seed yield than the introduced strain *B. japonicum* IRAT FA3 and the synthetic fertilizer NPK 12-22-22. RSC508 induced the highest pod number plant⁻¹ (102 pods) and seed yield (4.2 t ha⁻¹). Moreover, these local bacteria had positive effects on soil properties following the soybean's harvest. Among these bacteria, RSC119 enhanced mostly total nitrogen, organic matter and available phosphorus, then RSC508 significantly reduced soil acidity. This study therefore suggests that local rhizobia are effectiveness and could be use as inoculum to improve soybean productivity and soil properties restoration.

Keywords: native rhizobia, inoculation, soybean, soil properties, Côte d'Ivoire.

1. INTRODUCTION

Soybean (*Glycine max* L.) is one of the most important annual grain legume in the world. Its seed are an excellent source of protein (40%) and edible oil (20%) [1]. It also appears as the best crop to improve livelihoods and income for rural smallholder farmers in various developing countries. Soybean also increases soil fertility and productivity for its ability to fix and reduce atmospheric nitrogen (N₂) into biological usable forms in symbiosis with rhizobia [2]. Soybean-rhizobia symbiosis could provide easy and inexpensive way to enhance soil fertility and improve crop production while reducing synthetic fertilizers need in agricultural practices [3]. Successful soybean-rhizobia symbiotic relationship is dependent for a large part on the effectiveness and competitiveness of bacterial strains used as inoculant [4]. Moreover, to ensure profitable production of soybeans outside its domestication zone, research recommends the exogenous supply of bacterial strains capable of fixing nitrogen such as inoculum [5]. In Côte d'Ivoire, an exotic strain of *Bradyrhizobium japonicum* (IRAT FA3) was used as inoculant for soybean production since two decades [6]. However, the response of soybean genotypes to this exotic bacterial strain was variable to agroecological conditions [7]. Thus, soybean vulgarization in some areas of the country with no history to soybean cultivation, requires effective strains. Recent studies reported that isolation and selection of native efficient rhizobia strains on different soybean cultivars [8, 9]. Authors revealed that native rhizobia are able to improve soybean cultivars nodulation and growth better than introduced strain *B. japonicum* IRAT FA3 under controlled conditions. The capacity of these efficient local strains of rhizobia which are

adapted to local agroecological conditions deserves to be evaluated in field conditions in order to promote them as elite strains for inoculum formulation. Thus, the objective of this study was to evaluate the symbiotic and agronomic potential of native rhizobia strains on soybean and their background effects on soil properties on farm.

2. MATERIAL AND METHODS

The field experiment was conducted at the experimental station of Jean Lorougnon Guede University in Daloa. It is located in the Centre-West of Côte d'Ivoire at 6°84' N latitude and 6°41' W longitude. The soils of Daloa area were for the most part slightly to moderately desaturated reworked ferralitic type and ferrisols [10, 11]. Average rainfall was between 1400 and 1600 mm per year with an annual average of 1300 mm. The average annual temperature was 25.6°C [12]. The experimental site previous cultivation was cassava (*Manihot esculenta* Crantz) and it had no history of soybean cultivation nor inoculation with Bacteria Fixing Nitrogen (BNF) since the three (3) last years. Soybean [*Glycine max* (L.) Merrill] cultivar Piramama obtained from the Crops Research Station of the National Center for Agronomic Research (CNRA) in Bouaké (Côte d'Ivoire), was used as plant material in this experiment. Rhizobial strains used in this work consisted of seven (07) local rhizobia (RSC115, RSC119, RSC309, RSC324, RSC502, RSC504 and RSC508) isolated from soybean and one (01) exotic strain (*Bradyrhizobium japonicum* IRAT FA3) used as reference. Local and exotic rhizobia were provided respectively by the Laboratory of Agrovalorization of Jean Lorougnon Guede University in Daloa (Côte d'Ivoire) and the Central Laboratory of Soils, Water and Plants (LCSEP) of The CNRA in Bouake (Côte d'Ivoire).

Experiment was carried out according to a randomized complete block design with three replications. The main plot was inoculation with 10 modalities of which seven (07) native strains (RSC115, RSC119, RSC309, RSC324, RSC502, RSC504 and RSC508), one (01) exotic strain (*Bradyrhizobium japonicum* IRAT FA3), one (01) synthetic fertilizer NPK 12 22 22 formulation used as positive control (TN) and one (01) without treatment used as negative control (T0). Each plot size measured 3 m x 2 m with an alley of 1 m between plots. Seeds were sown with a spacing of 50 cm between rows and 20 cm between pockets. Symbiotic and agronomic parameters were measured at different phenological stages. Plant nodulation and growth parameters measured were nodules number, plant height and dry biomass. These parameters were assessed at the beginning of the flowering stage from five plants randomly selected from each plot. Plant height was measured with a meter rule. Nodules were carefully detached from the root of plants previously dug up and counted. Then, fresh biomass (root and shoot) of these same plants were oven-dried at 70 °C for 72 h and their dry weight recorded. At maturity, the average number of pods per plant was counted on five randomly selected plants. Then, seeds yield was determined. Analyses of composite soil sample taken before and 30 days after harvest were used to assess the impact of inoculation on soil quality. Analysed parameters were granulometry by densimetric method using Robinson's pipette [13], pH_{H2O} with a pH meter according to a soil/distilled water ratio of 1: 2.5 [14], organic carbon content by the Walkley and Black [15] method, then converted to organic matter (OM) using the factor 1.724, total nitrogen according to Kjeldahl method [16] and available phosphorus by modified Olsen method [17]. Data collected were subjected to analysis of variance (ANOVA) using STATISTICA 7.1 version. Fisher's Least Significant Difference (LSD) was used to separate means at a significance level of 5 %.

3. RESULTS AND DISCUSSION

3.1 Effect of inoculation on soybean nodulation and growth parameters

Inoculation had a significant influence ($P \leq .05$) on the number of nodules, plant height, and dry biomass of soybean (Table 1). In terms of the different rhizobia tested, RSC119 and RSC504 induced higher number of nodules (42 and 50 nodules respectively). These were 20 % to 43 % higher than the nodules observed on the reference strain IRAT FA3 (35 nodules per plant). On the other hand, nodules were observed on the roots of the negative control plants (3 nodules per plant) and positive control (5 nodules per plant). These nodules had a whitish colouration on inside. Results showed that native rhizobia strains tested were capable of inducing nodule formation on the soybean root system. On the other hand, a few nodules with a whitish internal colouring were observed on control plants of this cultivar. The presence of few nodules in non-inoculated plots reflects the lack of efficient indigenous rhizobia strains in the soils of the field trial. The formation of nodules on the uninoculated plants could also be due to indigenous rhizobia of the genus *Bradyrhizobium* which usually enter into symbiosis with cowpea (*Vigna unguiculata*) often grown in the trial area. However, the nodules formed were non-functional because the bacteria do not have nitrogen-fixing genes specific to soybean. Similar conclusions were drawn by previous studies, such as Nzabi et al. [18] and Argaw [19], who worked on various soybean genotypes in Ethiopia and Kenya respectively. Additionally, the study also found that several native rhizobia strains (RSC119, RSC309 and RSC504) can significantly increase nodule number compared to the exotic *Bradyrhizobium japonicum* IRAT FA3 strain used as soybean inoculum in Côte d'Ivoire. These results confirm observations made by Amani et al. [9] realized on pots under controlled conditions with the same strains. Moreover, works carried out in Bouaké, in the center of Côte d'Ivoire by N'Gbesso et al. [20] on the same soybean cultivar revealed that local bacterial strain 26D4 was able to increase nodule dry weight compared to introduced strain IRAT FA3. Other works carried out in the province

of Kabul in Afghanistan showed that native strains of *Ensifer* (GS4 and GE6W) and *Bradyrhizobium* (GE3) induced formation of more nodules on soybean varieties Stime3300 and Enrei compared to the reference strain USDA110 [21].

All rhizobia tested improved soybean height compared with negative control (T0). Native rhizobia RSC119 and RSC504 induced the highest plant heights at flowering with 44.4 cm and 44.5 cm respectively. Their effect on plant growth was similar to that of chemical fertilizer. Like these bacteria, RSC309, RSC324 and RSC508 had positive impact on soybean growth. Moreover, tested rhizobia strains except RSC115 increased plant dry biomass compared with negative control (T0). Local rhizobia RSC119 and RSC504 induced 4.8 g and 4.4 g of plant biomass respectively. In addition, the dry biomass produced by these bacteria was statistically similar to biomass induced by synthetic fertilizer (4.8 g). Regarding plant growth, parameters such as height and dry biomass were significantly influenced by native rhizobia tested. Indeed, all of these rhizobia improved the height and dry biomass of soybeans compared to the negative control. These rhizobia generally induced the same effects as the synthetic fertilizer NPK 12 22 22 for these different parameters. The improvement in height and biomass observed in this work would be due to the addition of nitrogen during the symbiotic fixation of atmospheric nitrogen. These results confirm those of Alam et al. [22] in Bangladesh. These authors showed that the height and dry biomass of soybean genotypes Shohag, BARI Soybean6, MTD10 and BGM02026 were improved by the local strain of *Rhizobium* sp. BARIGm901 compared to the uninoculated control. Regar et al. [23] similarly observed such enhancements in the Rajasthan region of India, leveraging a local rhizobial strain. Other work showed that inoculation practice improved plant height and biomass as well as synthetic nitrogen fertilization [24].

Table 1: Effect of local rhizobia, introduced strain and synthetic fertilizer NPK 12 22 22 on soybean nodules number, height and dry biomass of soybean at flowering stage

| Treatments | Nodules number | Plant height (cm) | Plant dry biomass (g) |
|------------|----------------|-------------------|-----------------------|
| RSC115 | 33±3e | 32,50±1,5e | 2,7±0,5f |
| RSC119 | 42±4b | 44,40±2,33ab | 4,8±0,4a |
| RSC309 | 37±2cd | 41,30±3,2cd | 3,9±0,3bc |
| RSC324 | 33±2de | 40,20±1,6cd | 3,1±0,3de |
| RSC502 | 27±4f | 39,40±1,1d | 3,1±0,5de |
| RSC504 | 50±3a | 44,50±2,3ab | 4,4±0,6ab |
| RSC508 | 38±3c | 41±3,2cd | 3,4±0,4cd |
| IRAT FA3 | 35±3cde | 42,60±1,2bc | 3,5±0,3cd |
| TN | 5±4g | 46,20±2a | 4,8±0,2a |
| T0 | 3±3d | 31,12±1,8e | 2,4±0,3f |
| Mean | 30±15 | 40,3±5,1 | 3,6±0,9 |
| CV (%) | 49,4 | 12,7 | 24 |
| LSD (5 %) | 4,4 | 3,1 | 0,6 |

In each column, means followed by the same letter are not significantly different at 5% threshold according to Fisher's LSD test. T0: Untreated control; TN: Synthetic fertilizer NPK 12 22 22.

3.2 Effect of inoculation on soybean pods and grain yield

Local tested rhizobia improved significantly ($P = .008$) both number and weight of pods per plant compared to negative control (Table 2). The number and weight of pods varied respectively from 44 to 102 pods per plant and from 30.9 to 56.1 g for all treatments. Native rhizobia RSC508 induced both the most pods (102 pods) per plant and the highest pod weight (56.1 g). RSC119 and RSC309 also boosted pod production per plant compared with synthetic fertilizer and reference strain IRAT FA3. These rhizobia induced the production of 99 and 96 pods per plant respectively, compared to pods induced by synthetic fertilizer (80 pods per plant) and reference strain IRAT FA3 strain (85 pods per plant). Soybeans inoculated with local rhizobia showed an increase in both number and weight of pods compared with the negative control. Furthermore, plants inoculated in this study with local rhizobia RSC119, RSC309 and RSC508 produced more pods than the introduced strain *B. japonicum* IRAT FA3 and the synthetic fertilizer NPK 12 22 22. According to Abdel-Fattah et al. [25], the inability of synthetic fertilizer to increase the number and weight of pods compared with inoculation is due to its low input to the soil. However, soybean plants inoculated with rhizobial strains may produce pod numbers that are statistically similar to those fertilized with synthetic fertilizers [26]. These results corroborate those obtained by Argaw [19] who compared the local Ethiopian *Bradyrhizobium japonicum* TAL-379 strain with an exotic *Bradyrhizobium* sp. strain UK isolate on soybean nodulation and growth. Also, soybean seeds yield per hectare was significantly influenced by

inoculation. Seeds yield per hectare varied from 0.7 t/ha to 4.3 t/ha. Seed yields were improved more by native rhizobia RSC119, RSC309 and RSC508. These rhizobia produced higher yields than synthetic fertilizer. Among these, RSC508 induced the highest yield of 4.3 t/ha compared with 3.5 t/ha for the positive control and 0.7 t/ha for the negative control. Patra et al. [27] achieved similar results with soybean cultivar PK-416 in India. According to Shahid et al. [28], the variation in seed yield between inoculated cultivars is largely due to the combined effect of bacterial isolates on vegetative and production parameters, which are themselves influenced by environmental conditions. Other authors mention that the increase in seed yield of different soybean varieties by rhizobial bacteria is linked to the low nitrogen content of tropical soils. This low nitrogen content in soils is at the origin of the initiation of symbiosis between the soybean and its symbiont [29, 30]. However, according to N'Cho et al [31] and Dabesa and Tana [32] inoculation of soybean seeds with the *Bradyrhizobium* strain alone did not increase soybean yields compared with synthetic fertilizer. They suggested that the bacterial inoculum should be combined with a phosphorus-based fertilizer to increase soybean yield.

Table 2: Effect of native rhizobia, exotic strain IRAT FA3 and synthetic fertilizer NPK 12 22 22 on the number and weight of pods and seed yield of soybean

| Treatments | Pods | | Seeds yield (t/ha) |
|------------|--------|------------|--------------------|
| | Number | Weight (g) | |
| RSC115 | 64±7c | 37,1±2,6f | 3,2e |
| RSC119 | 99±4a | 52,9±1,5ab | 3,7c |
| RSC309 | 96±8a | 50,9±6,2bc | 3,9b |
| RSC324 | 86±8b | 47,4±4,4cd | 3,5d |
| RSC502 | 68±3c | 32,6±1,9gh | 2,6f |
| RSC504 | 69±4c | 35±1,3fg | 3,2e |
| RSC508 | 102±9a | 56,1±4,1a | 4,3a |
| IRAT FA3 | 85±4b | 48,8±0,8cd | 3,6cd |
| TN | 80±3b | 45,8±1,9e | 3,5d |
| T0 | 44±4d | 30,9±1,81i | 0,7g |
| Mean | 79±18 | 43,7±9,1 | 3±0,5 |
| CV (%) | 23 | 20,9 | 36,3 |
| LSD (5 %) | 10,2 | 4,2 | 0,2 |

In each column, means followed by the same letter are not significantly different at 5% threshold according to Fisher's LSD test. T0: Untreated control; TN: Synthetic fertilizer NPK 12 22 22.

3.3 Background effects of local rhizobia on soil properties

Analyses carried out on soil sample taken 30 days after soybean harvest showed that inoculation had a significant influence ($P = 0.006$) on the physico-chemical properties of the soil (Table 3). All tested rhizobia tended to decrease the pH (H_2O) of the soil by 0.3 to 0.5 unit, except RSC508 strain which increased the pH by 0.2 unit compared to the initial soil state. The pH level induced by these bacteria was between 5.5 and 6.4. Results related to pH showed that some local rhizobia tended to make the environment more acidic. This soil acidity is always satisfactory for good biological activity. These results are not surprising because during rhizobia-legume symbiosis process, the different partners produce organic acids which tend to acidify the environment. In contrast, the local rhizobium RSC508 increased soil pH. This increase in soil pH is linked to several factors including the ability of the bacteria to induce the production of a large quantity of biomass. Total Nitrogen (Nt), Organic Carbon (OC) and Organic Matter (OM) were significantly increased by native rhizobia RSC119 compared with the other treatments and the initial state of the soil. This strain induced 3.12 % of OM against 1.27 % for the initial soil. In addition, it induced the highest level of Available Phosphorus (Av.P) compared to all the treatments, with 64 mg.kg⁻¹. Moreover, RSC508 and RSC502 tended to maintain soil phosphorus levels. The C/N ratio was generally between 11 and 12 for all treatments except with RSC324, RSC502 and T0. Results obtained show that native rhizobia strains used as inoculum during soybean crop establishment, have significant beneficial effects on the physico-chemical parameters of the soil 30 days after harvest. However, while soil organic matter (OM) and total nitrogen (Nt) levels increased, they did not reach

the minimum threshold (2%) required for total nitrogen. These observations are in line with those made by N'Goran et al. [33] with an association of yam and soybean. The increase in these elements in the soil is due, among other things, to the early fall of soybean leaves before harvest. N'Diaye et al. [34] attributed the increase in soil organic carbon and Nt levels to a supply of organic matter coming from all parts of the plant. For these authors, soil carbon and nitrogen are closely linked to soil organic matter (OM) which gives the latter physico-chemical properties that promote the sustainable functioning of ecosystems. They pointed out that the different parts of a plant with different biochemical qualities have different levels of degradation, giving the soil a continuous supply of organic matter. This contribution of legumes to OM is responsible for the increase in corn productivity, as demonstrated by Kouassi et al. [35]. Natural fallows generally play this same role at ground level, but it will still be necessary to observe for a longer period or even several years [36]. The C/N values obtained are generally between 9 and 12, reflecting effective mineralization of organic matter and intense microbial activity at the soil level. These results show the capacity of native rhizobia to induce significant production of soybean matter and an intensification of microbial activity. According to Konaté et al. [37], the acceleration of microbial biomass activity stimulates the decomposition and mineralization of pre-existing organic matter in the soil. Assimilable phosphorus and total phosphorus levels decreased with all treatments. This drop in assimilable P could be linked to the sandy texture of the site soils. The sandy texture exposes the soil to leaching and the loss of mineral elements in general. This assertion partly justifies the results obtained by Kouadio et al. [38], who maintain that the leaching phenomenon to which sandy soils are subjected during the rainy seasons exposes them to nutrient depletion.

Table 3: Background-effects of native rhizobia inoculated to soybean on soil quality

| Treatments | pH _{H2O} | OC (%) | N (%) | C/Nt | OM (%) | Av.P (mg.kg ⁻¹) |
|--------------|-------------------|---------|-------|------|--------|-----------------------------|
| Initial soil | 6,4a | 0,7ef | 0,06b | 12c | 1,27cd | 66a |
| RSC115 | 5,9b | 0,62f | 0,06b | 11d | 1,07d | 56bcd |
| RSC119 | 6b | 1,81a | 0,15a | 12c | 3,12a | 64a |
| RSC309 | 6,1b | 0,92cde | 0,08b | 11d | 1,58bc | 55bcd |
| RSC324 | 6,1b | 0,76def | 0,06b | 14a | 1,31cd | 54cd |
| RSC502 | 6b | 0,74ef | 0,06b | 13b | 1,27cd | 60abc |
| RSC504 | 6,1b | 0,96bcd | 0,08b | 12c | 1,64bc | 50de |
| RSC508 | 6,6a | 1,15b | 0,10b | 12c | 1,98b | 61ab |
| IRAT FA3 | 6,4a | 0,98bc | 0,08b | 12c | 1,68bc | 56bcd |
| TN | 6,4a | 0,76def | 0,07b | 11d | 1,31cd | 46e |
| T0 | 6,1b | 0,88cde | 0,07b | 13b | 1,51cd | 52de |
| Mean | 6,2 | 0,9 | 0,08 | 12 | 1,6 | 56 |
| CV (%) | 8,1 | 33,3 | 20 | 16,7 | 31,2 | 10,7 |
| LSD (5 %) | 0,3 | 0,2 | 0,05 | 1,0 | 0,4 | 6 |

In each column, means followed by the same letter are not significantly different at 5% threshold according to Fisher's LSD test. T0: Untreated control; TN: Synthetic fertilizer NPK 12 22 22.

4. CONCLUSION

The present study evaluated the performance of native rhizobia on soybean productivity and soil fertility in Côte d'Ivoire. Results indicated that native rhizobia strains are efficient and competitive to enhance soybean nodulation, growth and

yield. Native rhizobia strains RSC119 and RSC504 induced more nodules formation than the induced strain *Bradyrhizobium japonicum* IRAT FA3. Also, native rhizobia strains RSC119, RSC309, RSC504 and RSC508 induced more pods number and seed yield than synthetic fertilizer NPK 12 22 22 formulation. The highest number of pods per plant and seed yield were obtained with RSC508. Moreover; local rhizobia improved soil properties such as soil organic matter and total nitrogen on short periods after soybean harvest. Among these bacteria, the strain RSC508 increased soil pH. Results showed that soybean inoculation with ineffective and competitive local rhizobia strains significantly increased soybean productivity and helps to restore soil fertility over short periods.

UNDER PEER REVIEW

REFERENCES

1. Masciarelli O, Llanes A, and Luna V. A new PGPR co-inoculated with *Bradyrhizobium japonicum* enhances soybean nodulation. *Microbiological Research*. 2014; 169: 609-615.
2. Mahamood J, Abayomi YA and Aduloju M O. Comparative growth and grain yield responses of soybean genotypes to phosphorous fertilizer application. *African Journal of Biotechnology*. 2009; 8(6): 1030-1036.
3. Alori ET, Dare MO and Babalola OO. Microbial Inoculants for Soil Quality and Plant Health. *Sustainable Agriculture Reviews*. 2017 ; 22 : 281-307.
4. Nakei MD, Venkataramana PB and Ndakidemi PA. Soybean-nodulating rhizobia: ecology, characterization, diversity, and growth promoting functions. *Frontiers in Sustainable Food Systems*. 2023 ; 6 : 1-17. doi: 10.3389/fsufs.2022.1085843
5. Kim DH, Kaashyap M, Rathore A, Das RR, Parupalli S, Upadhyaya HD, Gopalakrishnan S, Gaur PM, Singh S, Kaur J, Yasin M and Varshney RK. Phylogenetic diversity of *Mesorhizobium* in chickpea. *Journal of Biosciences*. 2014 ; 39 : 513-517.
6. N'Gbesso MFDP, N'guetta ASP, Kouamé C and Foua KB. Evaluation of the efficiency of seed inoculation in 11 soybean genotypes (*Glycine max* L. Merrill) in the savannah zone of Côte d'Ivoire. *Science & Nature*. 2010; 7(1): 1812-074.
7. N'Zi JC, Koua AP, Kouassi KD, Kahia J, Kouassi JL, N'Guetta ASP and Kouamé C. Effect of inoculating seeds with *Bradyrhizobium japonicum* on the agronomic performance of five varieties of soybean (*Glycine max*) in Côte d'Ivoire. *African Journal of Agricultural Research*. 2015 ; 10(37) : 3671-3677.
8. Amani K, Konaté I, N'Gbesso MFDP, Attien YP, Fondio L, Filali-Maltouf A and Tidou AS. Phenotypic and symbiotic diversity of rhizobia isolated from root nodules of soybean [*Glycine max* (L.) Merrill] in Côte d'Ivoire. *International Journal of Current Microbiology and Applied Sciences*. 2019 ; 8(3) : 766-774.
9. Amani K, Fondio L, Konaté I, N'Gbesso MFDP, Beugré GAM, Tidou AS and Filali-Maltouf A. Response of indigenous rhizobia to the inoculation of soybean [*Glycine max* (L.) Merrill] varieties cultivated under controlled conditions in Côte d'Ivoire. *Advances in Microbiology*. 2020 ; 10(3) : 110-122.
10. Brou YT. Climate variability, deforestation and agro-demographic dynamics in Côte d'Ivoire. *Drought*. 2010; 21(1): 1-6.
11. Konaté Z, Abobi HDA, Soko FD and Yao-Kouamé A. Effects of soil fertilization using household solid waste composted in landfills on the yield and chemical quality of lettuce (*Lactuca sativa* L.) . *International Journal of Biological and Chemical Sciences*. 2018; 12(4): 1611-1625.
12. Brou YT, Akindès F and Bigot S. La variabilité climatique en Côte d'Ivoire : entre perceptions sociales et réponses agricoles. *Cahiers Agricultures*. 2005 ; 14(6): 533-540.
13. Gee GW and Bauder JW. Particle-size analysis, in *Methods of soil analysis*. Part 1: Physical and mineralogical methods. American society of Agronomy, Soil Science Society of America, Madison. 1986; 383-411.
14. Anderson JM and Ingram JSI. *Tropical Soil Biology and Fertility: A Handbook of Methods*. CAB International, Wallingford, UK, 2nd edition. 1993.
15. Walkley A and Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sciences*. 1934; 37: 29-36.
16. Bremner JM. Nitrogen-total in *Methods of soil analysis*, part 3 : Chemical methods. Soil Science Society of America Inc, American Society of Agronomy, Inc., Madison, Wisconsin, USA. 1996.

17. Dabin B. On a method for analyzing phosphorus in tropical soils. Symposium on tropical soil fertility, Antananarivo, Madagascar, November 19-25. 1967; 1:99-115.
18. Nzabi AW, Makini F, Mutai E, Gesare M and Mgwagi G. Influence of indigenous and introduced rhizobial strains on soybean grain yield in South West Kenya. Kenya Agricultural Research Center, Kisii. 2000.
19. Argaw A. Response of soybean to inoculation with *Bradyrhizobium* spp. in saline soils of Shinille plains, Eastern Ethiopia. East African Journal of Sciences. 8(2): 79-90, 2014.
20. N'Gbesso MFDP, Fondio L, Coulibaly ND and N. C. Kouamé, "Symbiotic effectiveness of five local strains of rhizobia on soybean growth parameters," International Journal of Biological and Chemical Sciences, vol. 11, no. 5, pp. 2327-2340, 2017.
21. Habibi S, Ayubi AG, Ohkama-Ohtsu N, Sekimoto H, and Yokoyama T. Genetic characterization of soybean rhizobia isolated from different ecological zones in North-Eastern Afghanistan. Microbes and Environments. 2017; 32(1): 71-79.
22. Alam F, Bhuiyan MAH, Alam SS, Waghmode TR, Kim PJ and Lee YB. Effect of *Rhizobium* sp. BARIRGm901 inoculation on nodulation, nitrogen fixation and yield of soybean (*Glycine max*) genotypes in gray terrace soil. Bioscience, Biotechnology and Biochemistry. 2015; 79(10): 1660-1668.
23. Regar MK, Meena RH, Gajanand J and Mundra SL. Effect of Different Rhizobial Strains on Growth and Yield of Soybean [*Glycine max* (L.) Merrill]. International Journal of Current Microbiology and Applied Sciences. 2017 ; 6(11) : 3653-3659.
24. Caliskan S, Arslan M, Uremis I and Caliskan ME. The effect of row spacing on yield and yield components of full season and double-cropped soybean. Turkish Journal of Agriculture and Forestry. 2007; 31(3): 147-154.
25. Abdel-Fattah I, El-Shaarawi A, Sawsan ZS, Abou-Taleb M and Ahmed EG. Effect of inorganic nitrogen and *Bradyrhizobium japonicum* inoculation on growth and yield of soybean. Australian Journal of Basic and Applied Sciences. 2011; 5(10): 436-447.
26. Jarecki W, Buczek J and Bobrecka-Jamro D. Response of soybean (*Glycine max* (L.) Merr.) to bacterial soil inoculants and foliar fertilization. Plant Soil and Environment. 2016; 62(9): 422-427.
27. Patra RK, Pant LM and Pradhan K. Response of soybean to inoculation with rhizobial strains: Effect on growth, yield, N uptake and soil N status. World Journal of Agricultural Science. 2012; 8(1): 51-54.
28. Shahid QM, Saleem MF, Khan HZ and Anjum SA. Performance of soybean (*Glycine max* L.) under different phosphorus levels and inoculation. Pakistan Journal of Agricultural Sciences. 2009; 46(4): 237-241.
29. Lamptey S, Ahiabor BDK, Yeboah S and Asamoah C. Response of soybean (*Glycine max*) to rhizobial inoculation and phosphorus application. Journal of Experimental Biology and Agricultural Sciences. 2014; 2(1): 72-77.
30. Kumar NR and Reddy RS. Screening of *Bradyrhizobium* isolates for plant growth promoting properties in vitro conditions. International Journal of Current Microbiology and Applied Science. 2018; 7(10): 2232-2237.
31. N'Cho CO, Yusuf AA, Ama-Abina JT, Jemo M, Abaidoo RC and Savane I. Effects of commercial microbial inoculants and foliar fertilizers on soybean nodulation and yield in northern Guinea savannah of Nigeria. International Journal of Advance Agricultural Research. 2013 ; 1 : 66-73.
32. Dabesa A and Tana T. Response of soybean (*Glycine max* L. (Merrill)) to *Bradyrhizobium* inoculation, lime, and phosphorus applications at bako, western Ethiopia. International Journal of Agronomy. 2021 ; 1-12.
33. N'Goran KE, Kassin KE, Zohouri GP, N'Gbesso MFDP and Yoro GR. Agronomic performance of yam-legume crop associations in the Central-West of Côte d'Ivoire. Journal of Applied Biosciences. 2011; 43:2915-2923.
34. N'Diaye O, Diop AT, Akpo LE and Diène M. Dynamics of soil carbon and nitrogen content in Ferlo farming systems: Case of Dahra CRZ. Journal of Applied Biosciences. 2014; 83:7554-7569.

35. Kouassi NJ, Tonessia DC, Seu JG, Soko DF, Ayolié K. Influence of delayed sowing of maize (*Zeamays L.*) and Bambara groundnut (*Vignasubterranea (L.) Verdc.*) on their production in savannah zones from Ivory Coast. *Journal of Applied Biosciences*. 2016; 102:9745-9755.
36. Gnahoua GM, Kouassi FY, Angui PKT, Balle P, Olivier R and Peltier R. Effects of *Acacia mangium*, *Acacia auriculiformis* and *Chromolaena odorata* fallows on soil fertility and yam (*Dioscorea spp.*) yields in forest zone of Ivory Coast. *African Agronomy*. 2008; 20(3): 291-301.
37. Konaté Z, Gala TJB, Messoum FG, Sekou A, Yao-Kouamé A, Camara M and Kéli ZJ. Alternatives to mineral fertilization of soils in rainfed upland rice cultivation: contributions of soybean and cowpea crops to the fertility of hyperdystric ferralsol in the central-west of Côte d'Ivoire. *Journal of Applied Biosciences*. 2012; 54:3859-3869.
38. Kouadio KP, Yoboué KE, Kouadio KKH, Bini YYP and Yao-Kouamé A. Morpho-pedological characteristics of Ahoué soils in the Brofodoumé Sub-Prefecture, South-East Ivory Coast. *Africa Science*. 2019; 15(5): 140-150.

UNDER PEER REVIEW