

## Original Research Article

# Estimation of avoidable yield losses due to pest complex in cowpea [*Vigna unguiculata* (L.) Walp.]

### ABSTRACT

A field experiment was conducted during the *Kharif* seasons of 2022 and 2023 at the research farm of Rajasthan Agricultural Research Institute, Durgapura, Jaipur, Rajasthan, India, to evaluate the yield losses caused by a pest complex on cowpea. The pooled study of both ~~the~~ year revealed that the highest total avoidable loss was observed in the untreated control (NP) at 40.90%, followed by the PVS treatment at 24.45%. The lowest avoidable loss was recorded in the PRS treatment at 13.49%. The treatment PTS showed the highest increase in cowpea seed yield over control, with a 69.24% increase, followed by PRS with a 46.29% increase. Conversely, the PVS treatment exhibited the lowest yield increase at 27.83%. The maximum yield of 1049.70 kg ha<sup>-1</sup> was achieved with the PTS treatment, while the lowest yield of 620.40 kg ha<sup>-1</sup> was recorded in the NP treatment. The PVS and PRS treatments yielded 793.10 kg ha<sup>-1</sup> and 908.20 kg ha<sup>-1</sup>, respectively.

**Keywords:** *Pest complex, cowpea, yield losses, treatments.*

### 1. INTRODUCTION

Cowpea [*Vigna unguiculata* (L.)] (Family: Leguminaceae) is one of the most important principle pulse crop of tropics and commonly known as crowdel pea, Chala, Chola or Choli, Chavli, Lobia, southern pea and black eyed bean. In Rajasthan, cowpea is grown on 79000 hectares area with 27000 metric tonnes production and average yield of 337 kg per hectare (Anonymous, 2022). The important insect species attacking cowpea crop are aphid, *Aphis craccivora* Koch; jassid, *Empoasca fabae* (Harris); whitefly, *Bemisia tabaci* (Genn.); thrips, *Megaleurothrips distalis* Karny; and spotted pod borer, *Maruca vitrata* (Fab.) resulting in heavy yield losses (Prasad *et al.*, 1983 and Satpathy *et al.*, 2009). The spotted pod borer is one of the most important pests of cowpea and causes severe yield losses (up to 60%) in the tropics and sub tropics (Singh *et al.*, 1978). *Maruca vitrata* attacks cowpea during the reproductive phase. The female moth lays eggs on or near the flower buds (Sharma, 1998). The larvae of spotted pod borer are known to cause damage by webbing the leaves, bud, flower and pods together and feed inside them. The whitefly, *Bemisia tabaci* (Genn.), feeds on plant sap and excretes a sugary substance known as honeydew. This honeydew promotes the growth of sooty mold, which can significantly weaken plants and impair their ability to photosynthesize. Infested plants typically exhibit yellowing leaves and stunted growth. In addition to causing direct damage through feeding, whiteflies are also vectors for transmitting

Formatted: Font: Not Bold

Comment [U011]: Write your conclusion and recommendation at the end of the abstract

Comment [U012]: Write in alphabet order

Comment [U013]: In which country?

Comment [U014]: Include Cowpea bruchid (*Callosobruchus maculatus*)

plant viruses (Gerling, 1990). The leafhopper, *Empoasca fabae*, feeds on the cell sap from the lower surface of leaves and injects toxic substances, leading to symptoms such as yellowing and curling of leaf margins, as well as stunted plant growth. Severe infestations can cause leaf burning and subsequent leaf drop, resulting in a significant decrease in yield ranging from 40 to 60 per cent (Narke and Suryawanshi, 1987).

**Comment [U015]:** Please kindly indicate the aims and objectives of this research at the end of the introduction.

## 2. MATERIALS AND METHODS

The experiment was conducted using a simple Randomized Block Design (RBD). Cowpea genotype CPD-119 was evaluated to estimate avoidable losses due to a pest complex in cowpea, with each treatment replicated five times. Each treatment was sown in plots measuring 1.2 x 3 m<sup>2</sup>, maintaining row to row and plant to plant distances of 30 cm and 10 cm, respectively.

Treatments

- T<sub>1</sub>: Protection against insect pests throughout crop season (PTS).
- T<sub>2</sub>: Protection against insect pests throughout vegetative stage (before flowering stage) (PVS).
- T<sub>3</sub>: Protection against insect pests from flowering to harvesting stage (PRS).
- T<sub>4</sub>: No protection against insect pests (NP).

**Table 1. Details of treatments**

Treatment	Insecticides	Dosage
T <sub>1</sub>	Imidacloprid 17.8 SL and chlorantraniliprole 18.5 SC	20g a.i./ha
T <sub>2</sub>	Imidacloprid 17.8 SL	20g a.i./ha
T <sub>3</sub>	Chlorantraniliprole 18.5 SC	20g a.i./ha
T <sub>4</sub>	Unprotected	-

\* Imidacloprid 17.8 SL was applied 30 days after sowing and chlorantraniliprole 18.5 SC was applied at the 50 per cent flowering stage.

The percentage of avoidable loss in cowpea seed yield was calculated separately for different protection levels based on the cowpea seed yield (kg/ha).

The yield data of cowpea seeds from protected and unprotected plots was recorded. The yield difference in protected plots over unprotected plots was calculated and the avoidable loss was determined using the following formula. (Pradhan, 1964).

$$\text{Avoidable Loss (\%)} = \frac{\text{Yield in treated plot} - \text{Yield in untreated control plot}}{\text{Yield in treated plot}} \times 100$$

$$\text{Increase in yield (\%)} = \frac{\text{Yield in treated plot} - \text{Yield in untreated control plot}}{\text{Yield in untreated plot}} \times 100$$

### 3. RESULTS AND DISCUSSION

#### 3.1 *Experiment at Kharif, 2022*

##### 3.1.1 Yield

The yield of cowpea worked out separately in each treatment. From these data, yield of cowpea seed was converted to  $\text{kg ha}^{-1}$  and these data were analyzed.

Significant differences were also observed among different treatments. The highest yield ( $1054.60 \text{ kg ha}^{-1}$ ) was achieved in the treatment of PTS, whereas it was lowest ( $631.80 \text{ kg ha}^{-1}$ ) in the treatment of NP. The yield of PVS and PRS treatments was  $810.20$  and  $923.60 \text{ kg ha}^{-1}$ , respectively. The descending order of treatments on the basis of total yield of cowpea seed was found to be  $\text{PTS} > \text{PRS} > \text{PVS} > \text{NP}$ .

##### 3.1.2 Avoidable loss

The per cent avoidable loss in cowpea seed yield was worked out in different protection levels separately on the basis of cowpea seed yield ( $\text{kg ha}^{-1}$ ).

The results explicated that the highest avoidable loss was noticed in the treatment NP (40.09%) and it was followed by PVS (23.17%) treatment, whereas, it was lowest in the treatment of PRS (12.42 %). The descending order of treatments on the basis of percent avoidable loss in cowpea seed yield was  $\text{NP} > \text{PVS} > \text{PRS}$ .

##### 3.1.3 Per cent increase in seed yield of cowpea over control

Per cent increase in cowpea seed yield over control was worked out in different levels of protection on the basis of cowpea seed yield.

The overall increase in cowpea seed yield over control was maximum in treatment of PTS (66.92 %). It was followed by the treatment PRS (46.19 %). The minimum per cent increase in cowpea seed yield was observed in treatment of PVS (28.24%) as the crop was kept unsprayed during the most vulnerable stage *i.e.*, reproductive stage of cowpea. The descending order of different protection level based on per cent increase in cowpea seed yield over control was  $\text{PTS} > \text{PRS} > \text{PVS}$  indicating the turn the order of effectiveness of these treatments.

#### 3.2 *Experiment at Kharif, 2023*

##### 3.2.1 Yield

The yield of cowpea worked out separately in each treatment. From these data, yield of cowpea seed was converted to  $\text{kg ha}^{-1}$  and these data were analyzed.

Significant differences were also observed among different treatments. The highest yield ( $1044.80 \text{ kg ha}^{-1}$ ) was achieved in the treatment of PTS, whereas it was lowest ( $609 \text{ kg ha}^{-1}$ ) in the treatment of NP. The yield of PVS and PRS treatment was  $776$  and  $892.80 \text{ kg ha}^{-1}$ , respectively. The

Formatted: Font: (Default) Arial, Italic, Font color: Black

Comment [U016]: How was it converted?

Comment [U017]: Kindly state the table you are writing and discussing on

Comment [U018]: Please kindly state the pest of disease (s) responsible for the loss

Formatted: Font: (Default) Arial, Italic

descending order of treatments on the basis of total yield of cowpea seed was found to be PTS > PRS > PVS > NP.

### **3.2.2 Avoidable loss**

The per cent avoidable loss in cowpea seed yield was worked out in different protection levels separately on the basis of cowpea seed yield ( $\text{kg ha}^{-1}$ ).

The results explicated that the highest avoidable loss was noticed in the treatment NP (41.71%) and it was followed by PVS (25.73%) treatment, whereas, it was lowest in the treatment of PRS (14.55 %). The descending order of treatments on the basis of percent avoidable loss in cowpea seed yield was NP > PVS > PRS.

### **3.2.3 Per cent increase in seed yield of cowpea over control**

Per cent increase in cowpea seed yield over control was worked out in different levels of protection on the basis of cowpea seed yield.

The overall increase in cowpea seed yield over control was maximum in treatment of PTS (71.56 %). It was followed by the treatment PRS (46.39 %). The minimum per cent increase in cowpea seed yield was observed in treatment of PVS (27.42%) as the crop was kept unsprayed during the most vulnerable stage, i.e., reproductive stage of cowpea. The descending order of different protection level based on per cent increase in cowpea seed yield over control was PTS > PRS > PVS indicating the turn the order of effectiveness of these treatments.

## **3.3 Pooled (Kharif, 2022 and 2023)**

### **3.3.1 Yield**

The yield of cowpea worked out separately in each treatment. From these data, yield of cowpea seed was converted to  $\text{kg ha}^{-1}$  and these data were analyzed.

Significant differences were also observed among different treatments. The highest yield ( $1049.70 \text{ kg ha}^{-1}$ ) was achieved in the treatment of PTS, whereas it was lowest ( $620.40 \text{ kg ha}^{-1}$ ) in the treatment of NP. The yield of PVS and PRS treatments was  $793.10$  and  $908.20 \text{ kg ha}^{-1}$ , respectively. The descending order of treatments on the basis of total yield of cowpea seed was found to be PTS > PRS > PVS > NP.

### **3.3.2 Avoidable loss**


The per cent avoidable loss in cowpea seed yield was worked out in different protection levels separately on the basis of cowpea seed yield ( $\text{kg ha}^{-1}$ ).

The results explicated that the highest total avoidable loss was noticed in the treatment NP (40.90%) and it was followed by PVS (24.45%) treatment, whereas, it was lowest in the treatment of PRS (13.49 %). The descending order of treatments on the basis of percent avoidable loss

Comment [U019]: How was it converted?

incowpea seedyieldwasNP >PVS > PRS.

UNDER PEER REVIEW



**Table: 2 Seed yield and avoidable losses due to pest complex in different protection levels in cowpea during *Kharif*, 2022 and 2023**

S. No	Protection levels	Treatments	Seed yield (kg/ha)			Avoidable loss in cowpea seed yield (%)			Increase in cowpea seed yield over control (%)		
			2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
1.	Protection against insect pests throughout crop season (PTS).	Imidacloprid 17.8 SL and chlorantraniliprole 18.5 SC	1054.6	1044.8	1049.7	-	-	-	66.92	71.56	69.24
2.	Protection against insect pests throughout vegetative stage (before flowering stage) (PVS)	Imidacloprid 17.8 SL	810.2	776	793.1	23.17	25.73	24.45	28.24	27.42	27.83
3.	Protection against insect pests from flowering to harvesting stage (PRS)	Chlorantraniliprole 18.5 SC	923.6	892.8	908.2	12.42	14.55	13.49	46.19	46.39	46.29
4.	No protection against insect pests (control) (NP).	-	631.8	609	620.4	40.09	41.71	40.90	-	-	-
S.Em ±			34.57	36.15	36.91						
C.D. (P=0.05)			106.51	111.37	107.73						

**Comment [U0110]:** This table is not scientific, kindly present it in a scientific format

### 3.3.3 Per cent increase in seed yield of cowpea over control

Per cent increase in cowpea seed yield over control was worked out indifferent levels of protection on the basis of cowpea seed yield.

The overall increase in cowpea seed yield over control was maximum intreatment of PTS (69.24 %). It was followed by the treatment PRS (46.29 %). The minimum per cent increase in cowpea seed yield was observed in treatment of PVS (27.83%) as the crop was kept unsprayed during the most vulnerable stage i.e., reproductive stage of cowpea. The descending order of different protection level based on per cent increase in cowpea seed yield over control was PTS > PRS > PVS indicating the turn the order of effectiveness of these treatments.

The results are agreement with those of Anusha and Balikai (2015) who reported the apparent losses caused by pod borers and sucking pests in cowpea to be between were 47.23 to 62.52 per cent. The present findings align with Kanhere *et al.* (2012), who reported an 84.25 per cent yield improvement in protected plots over the unprotected plots, with an avoidable loss of 45.73 per cent due to pod borer damage. Similarly, Duraimurugan and Tyagi (2014) was also reported 32.97 per cent yield losses in green gram. These results are also conform Thisity with the findings of Shukla (2005) and Rathwa *et al.* (2018). The varied damage caused by major insect pests of cowpea was due to the varied biotic and abiotic factors of various localities.

**Comment [U0111]:** This discussion is too small, kindly explain your result better

### 4. CONCLUSION

The PTS treatment had the highest yield (1049.70 kg ha<sup>-1</sup>), with the lowest in NP (620.40 kg ha<sup>-1</sup>). The PVS and PRS treatments yielded 793.10 kg ha<sup>-1</sup> and 908.20 kg ha<sup>-1</sup>, respectively. The highest total avoidable loss was in NP (40.90%) followed by PVS (24.45%), and the lowest in PRS (13.49%). The PTS treatment also showed the highest yield increase over control (69.24%) followed by PRS (46.29%), with PVS showing the lowest increase (27.83%).

**Comment [U0112]:** This conclusion is too shallow, please kindly recast

### REFERENCES

1. Anonymous. Agricultural Statistics of Rajasthan. Directorate of Economics and Statistics, Department of Planning, Rajasthan, Jaipur. 2022;96.
2. Anusha CH, Balikai RA. Avoidable loss estimation due to pod borers in C-152 and DC-15 varieties of cowpea. Journal of Experimental Zoology India. 2012;18(1):349-353.
3. Duraimurugan P, Tyagi K. Pest spectra, succession and its yield losses in mungbean and urdbean under changing climatic scenario. Legume Research. 2014;37:212-222.
4. Gerling D. Natural enemies of whiteflies: predators and parasitoids, Whiteflies: their bionomics. Pest status and management. Intercept Andover, United Kingdom. 1990;147-185.
5. Kanhere RD, Patel VN, Umbarkar PS. Estimation of yield losses caused by insect pests infesting cowpea. Insect Environment. 2012;17(4):182-183.

6. Narke CG, Suryawanshi DS. Chemical control of major pests of okra. *Pesticides*. 1987;21(1):37-42.
7. Pradhan S. Assessment of losses by insect pests of crops and estimation of insect population. *Entomology in India*. Entomology Society of India, New Delhi. 1964;17-58.
8. Prasad D, Singh KM, Katiyar R.N. Succession of insect pests in early maturing high yielding varieties of pea, *Pisum sativum* Linn. *Indian Journal of Entomology*. 1983;45(4):451-455.
9. Rathwa MG, Patel VN, Makwana AB. Estimation of yield losses due to spotted pod borer, *Maruca testulalis* in cowpea. *International Journal of Chemical Studies*. 2018;6(2):322-324.
10. Satpathy S, Shivalingaswami TM, Kumar A, Raj AB, Rai M. Efficacy of biopesticides and new insecticides for managements of cowpea pod borer, *Maruca vitrata*. Symposium on international conference on grain legumes: Quality improvement value addition and trade, held on February 14-16, 2009 at IIPR, Kanpur. 2009;292-293.
11. Sharma HC. Bionomics, host plant resistance and management of the legume pod borer, *Maruca vitrata*-a review. *Crop Protection*. 1998;17(5):373-386.
12. Shukla NP. Bionomics of spotted pod borer *Maruca vitrata* (Fabricius), pest succession and management of pest complex of cowpea. Ph.D. Thesis (unpublished) submitted to Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar; 2005.
13. Singh SR, Van Emden HF, Taylor TA. Resistance to pests of cowpea in Nigeria. *Pests of Grain Legumes: Ecology and Control*, London, New York: Academic press. 1978;414.