

Short communication

Study on the abundance and diversity of soil invertebrate macro-fauna across different land use types at the GKVK Campus, Bengaluru

Abstract

At the University of Agricultural Sciences, GKVK, Bengaluru, four different land use types—roadside soil, forest ecosystem, paddy ecosystem ZARS, and botanical garden are taken into consideration in this study to analyze the abundance, diversity, and community composition of soil invertebrate macro-fauna. Three soil macrofauna sampling sites are selected from each land use category. The transect has a randomly selected beginning point, and tiny land units (25*25*30 cm) are excavated at 5-meter intervals perpendicular to the slope. A total of 12 units (4 treatments * 3 replications) of various land use types were sampled in May 2021. Berlese-Tullgren funnels are used for excising and collecting soil invertebrates. This study's results indicate that intensive land cultivation leads to quantitative alterations in the abundance and diversity of soil invertebrate macrofauna (SIMF) communities.

Keywords Soil macrofauna, Berlese-Tullgren funnel, land use type, abundance and diversity

Introduction

Soil macrofauna diversity, abundance, and community composition are all impacted by land use type. Changes in land use and intensification have an influence not just on plant communities but also on soil food webs and the connections between above- and below-ground populations (Wardle *et al.*, 2004). Annual cropping techniques don't provide permanent soil cover, it will disturb the soil and reduce the diversity and abundance of soil fauna communities (Barrios, 2007; Rossi *et al.*, 2010). Soil macro-fauna are also impacted by unsustainable land management techniques such as overgrazing, fire, deforestation, pollution, soil erosion, and fertility depletion (Bignell *et al.*, 2005). The development of soil macrofauna is facilitated by forest land and garden fields, which have more soil cover and less soil disturbance. These characteristics are absent from grazing areas and crop-cultivated fields, which leads to greater soil compaction, deterioration, and a lack of food and cover that are necessary for the survival of soil macrofauna (Moreira *et al.*, 2008).

It is well established that residue inputs and soil management techniques affect the distribution, abundance, and diversity of soil invertebrate macro-fauna (SIMF) (Brown *et al.*, 2004; Manhaes *et al.*, 2013). It is therefore assumed that there are variations in the effects of various land

Commented [sa1]:

Commented [sa2]: A lot of Grammar mistake in the paper so remove these mistake.

Commented [sa3]: Article wrong so need to write again properly

Commented [sa4]: Check the sentence and improve it?

Commented [sa5]: Provide a brief context about the abundance and diversity of soil invertebrate macro-fauna.

Commented [sa6]: Reference must be provided at least 2015 or above

Commented [sa7]:

management techniques on SIMF. Prior studies conducted at the UAS, GKVK, campus Bangalore, have revealed a knowledge gap about local SIMF assemblages in several land use categories. To discover how land use changes, impact the abundance and diversity of SIMF.

Commented [sa8]: Improve the abstract quality it's not give complete attention to the topic

Commented [sa9]: Also clearly state the specific objective

Materials and Methods

Description of the study area

The study was carried out in four primary land use types: roadside soil; botanical garden; forest ecosystem; and paddy ecosystem ZARS at UAS, GKVK, Bangalore, which is situated in the northern region of the Bangalore district; Geographically, the region is located between Latitude: 13° 04' 55.92" N and Longitude: 77° 34' 34.57" E.

Sampling Methods and Experimental Design:

Sampling was carried out in May 2021, during the beginning of light rainfall, when soil macrofauna activity is normally heightened. Every category of land use (Plate 1-3) has 3 sampling locations. Using an excavator tool soil was dug to a depth of 25 × 25 × 30 cm at 10-meter intervals along a transect perpendicular to the slope and with randomly placed starting points (Anderson & Ingram, 1993). Three sample monoliths were taken from each type of land use to create a total of twelve sampling points. Using Berlese funnels (Muvengwi *et al.*, 2018) (Plate 4), the macro-fauna was separated from the soil, collected in vials with 70% alcohol, then identified and counted. After the extraction of macrofauna, the extracted soil was put back in its original location to reduce soil disturbance in selected land use types.

Commented [sa10]: And then write that

Identification of soil macro-fauna

The collected soil invertebrates were identified and counted. According to Eaton and Kaufman (2007), identification keys and visual aids such as photographs were used in the laboratory to identify species of soil invertebrate macro-fauna (SIMF). Four different land use categories were evaluated for macro-fauna abundance (Table 1). The abundance and diversity of each macrofauna within the four land use groups were calculated.

Results

Commented [sa11]: Need more work in results. Properly discuss with findings.

There was a total 55 soil macrofauna were collected from four land use categories (Table 2). While earthworms and Collembola were collected in the forest, roadside, and botanical areas but absent from the cultivated paddy habitat. Ants and spiders were identified in all 4 land use types (Fig. 1). The abundance of SIMF was observed in the order of forest land > botanical garden > paddy field > roadside soil. Overall, the abundance of SIMF was lowest in cultivated field and roadside soil and highest in forest land (49.09%) followed by the botanical garden (29.09%) (Table 1).

Commented [sa12]: Was a total of 55

Discussion

Composition and occurrence of SIMF community

Under unsustainable land use management methods, the reaction of the soil invertebrate macrofauna (SIMF) community to environmental disturbance became apparent. Notably, SIMF groups showed preferences in the habitats of forests and botanical gardens, possibly as a result of less soil disturbance and/or higher quality of litter. The positive relationship between vegetation and the diversity of soil macro-fauna suggests that higher availability of energy and nutrients enhances the growth and activity of soil macrofauna (Negasa *et al.*, 2017). This preference could be explained by the higher levels of organic matter found in garden and forest areas, which offer beneficial substrates for soil macrofauna (Beare *et al.*, 1997). Conversely, paddy-cultivated fields had lower organic matter content and greater soil acidity conditions will lower the SIMF. Consistent with findings by Decaens *et al.* (1994), soil macro-fauna abundance tends to decrease to low levels in crop-cultivated lands.

Commented [sa13]: Change reference and add latest

Commented [sa14]:

Commented [sa15R14]: Discuss the reason about these lands have not soil invertebrate macro-fauna? Reason give

Conclusion

The findings of this study indicate that intensive cultivation of land leads to quantitative alterations in the abundance and diversity of soil invertebrate macrofauna (SIMF) communities. These changes are attributed to specific management practices that result in habitat destruction and the removal of organic substrate, consequently diminishing the availability of food sources for associated SIMFs. Among the four land use classes examined, forested areas exhibit relatively high diversity of SIMFs followed by botanical garden, because of low soil disturbance and availability food sources in this land use type. crop cultivated fields and roadside soil display lower diversity in invertebrates because land disturbance by human activity. Therefore, it is advisable to adopt sustainable cropping systems that uphold acceptable levels of SIMF abundance while

Commented [sa16]: need more hard work on the discussion and especially in the conclusion.

minimizing human disturbances. Achieving this goal necessitates incorporating knowledge of biological processes into the design of land management systems. Additionally, further in-depth studies are crucial to identifying the optimal combination of land use and varying management practices across different land types, ensuring the most effective sustainability of SIMF populations.

Commented [sa17]: Summarize the key findings and their implications abundance and diversity of soil invertebrate macro-fauna across different land use types at the GKVK Campus, Bengaluru

Table 1: Composition and distribution of soil invertebrate macro-fauna in different land use types at GKVK, Bangalore

Taxa	Distribution			
	Paddy Ecosystem ZARS	Botanical Garden	Forest Ecosystem	Roadside Soil
Earthworm				
Collembola	-	+	-	+
Spider	-	+	+	-
Ant	+	+	+	+
Millipede	+	+	+	+

+ indicates present, - indicates absent

Table 2: Abundance of each taxon in different land use types

Land use types	Earthworm	Collembola	Spider	Ants	Millipede	Total % of arthropods
Paddy ecosystem	0	0	5	2	0	12.72%
Botanical garden	3	4	8	1	0	29.09%
Forest ecosystem	0	9	11	4	3	49.09%
Roadside soil	2	0	1	1	2	10.90%






Plate 2: Setting of Berlese Funnel for the Estimation of Soil Invertebrate population from collected samples

Commented [sa18]: Describe pictures properly with finding in discussions.

Commented [sa19]:



desdica

Commented [sa20]:

Fig.1. Density of soil invertebrate macro-fauna in different land use types at GKVK, Bangalore

REFERENCE

- Anderson, J. M., & Ingram, J. S. I. (Eds.) (1993). *Tropical soil biology and fertility: A handbook of methods*, Wallingford: CAB International.
- Barrios, E. (2007). Soil biota, ecosystem services and land productivity. *Ecological Economics*, 64: 269–285.
- Beare, M. H., Reddy, M. V., Tian, G., & Srivastava, S. C. (1997). Agricultural intensification, soil biodiversity and agroecosystem function in the tropics: The role of decomposer biota. *Applied Soil Ecology*, 6: 87–108.
- Bignell, D. E., Tondoh, J., Dibog, L., et al. (2005). Below-ground biodiversity assessment developing a key functional group approach in bestbet alternatives to slash and burn. In C. A. Palm, S. A. Vosti, P. A. Sanchez, & P. J. Ericksen (Eds.), *Slash and burn agriculture: The search for alternatives*, 119–142. New York: Columbia University Press p.
- Brown, G. G., Moreno, A. G., Barois, I., et al. (2004). Soil macro-fauna in SE Mexican pastures and the effect of conversion from 551 native to introduced pastures. *Agro Ecosystems & Environment*, 103: 313–327
- Decaens, T., Lavelle, P., Jimenez, J. J., et al. (1994). Impact of land management on soil macrofauna in the oriental llanos of Colombia. *European Journal of Soil Biology*, 30: 157–168.
- Eaton, E. R., & Kaufman, K. (2007). *Kaufman field guide to insects of North America*. New York: Hillstar Editions.
- Manhaes, C. M. C., Gama-Rodrigues, E. F., Moço, M. K. S., et al. (2013). Meso-and macrofauna in the soil and litter of leguminous trees in a degraded pasture in Brazil. *Agroforestry Systems*, 87(5): 993 –1004.
- Moreira, F. M., Huising, E. J., & Bignell, D. (2008). *A handbook of tropical soil biology: Sampling and characterization of below-ground biodiversity*. London: Earthscan.

Muvengwi, J., Davies, A.B., Parrini, F. and Witkowski, E.T., (2018). Geology drives the spatial patterning and structure of termite mounds in an African savanna. *Ecosphere*, 9(3): 02148.

Negasa, T., Ketema, H., Legesse, A., Sisay, M., & Temesgen, H. (2017). Variation in soil properties under different land use types managed by smallholder farmers along the toposequence in southern Ethiopia. 290(20170): 40-50.

Rossi, J. P., Celini, L., Mora, P., et al. (2010). Decreasing fallow duration in tropical slash-and-burn agriculture alters soil 660 macro invertebrate diversity: A case study in southern French Guiana. *Agriculture, Ecosystems & Environment*, 135: 148–154.

Wardle, D. A., Bardgett, R. D., Klironomos, J. N., et al. (2004). Ecological linkages between aboveground and belowground biota. *Science.*, 304(5677): 1629 – 1633.

Commented [sa21]: Manage properly format table specially

Commented [sa22]:

UNDER PEER REVIEW