

**PRE-HOSPITAL ANTIBIOTICS USE IN THE RIVERS STATE UNIVERSITY
TEACHING HOSPITAL PAEDIATRIC OUTPATIENT CLINIC**

ABSTRACT

Background

Antibiotics are prescription-only medications but can be procured over-the-counter. This over-the-counter use of antibiotics is of global health concern as it has led to antibiotics resistance, increased serious side effects, increased cost as well as cycle of abuse/misuse.

Comment [EAMS1]: over the counter

Comment [EAMS2]: over the counter

Comment [EAMS3]: antibiotic

Comment [EAMS4]: sever

Comment [EAMS5]: as and a

Objective of the study

To determine the prevalence of pre-hospital use of antibiotics.

Methodology

An observational study involving 401 caregiver/child pair seen at the Paediatric Outpatient clinic of a tertiary health facility over 6 months. Data on demography, clinical symptoms, and weight of the children were obtained using a semi-structured questionnaire and informed consent obtained from the caregivers.

Comment [EAMS6]: six

Comment [EAMS7]: Data on the children

...

Comment [EAMS8]:

Results

A total of 401 caregiver/child pairs participated in the study with male predominance. Eighty nine (22.2%) of the children received antibiotics before presentation. The most common reasons for administering antibiotics were fever and cough. Majority (69.9%) gave syrup formulation. The reasons for choice of formulation were age of child, available formulation at home and it been more effective. Amoxicillin, augmentin and cefpodoxime were the most common antibiotics given and in 75% of cases, were not prescribed by a doctor.

Comment [EAMS9]: The majority

Comment [EAMS10]: the choice

Comment [EAMS11]: the age

Comment [EAMS12]: the child

Comment [EAMS13]: the availability of

Comment [EAMS14]: its being more effective

Comment [EAMS15]: , in

Comment [EAMS16]: the appropriate

Comment [EAMS17]: doses

Comment [EAMS18]: underdosed

Comment [EAMS19]: and

Comment [EAMS20]: an overdose.

They also received vitamin C, antimalarials and cough syrup before presentation. About 62% of the children received appropriate dose, 32.4% underdose while 5.6% received overdose.

Most of the children were diagnosed with malaria and respiratory tract infections in the hospital.

Conclusion

Pre-hospital antibiotics use and abuse/misuse is a common practice among the respondents.

Comment [EAMS21]: antibiotic

Education of the populace is advocated to curb this menace of antibiotic abuse with its attendant development of drug resistance.

Key words

Abuse, antibiotics, Paediatric, Pre-hospital

Introduction

Antibiotics are medications used for the treatment of infections caused by bacteria thus playing a crucial role in the reduction of morbidity and mortality in children. They are commonly used especially in developing countries, as infectious diseases account for one of the commonest causes of morbidity and mortality. [1] Rational use of these drugs is of utmost concern as overuse or misuse has led to the growth of antimicrobial resistance (AMR) globally which threatens its' ability to treat common infections. The World Health Organization (WHO) has documented that more than 50% of antibiotic use globally is inappropriate. [2] It is noteworthy that antibiotics could be misused through self-medication/pre-hospital administration with/without poor adherence to treatment. [3]

Comment [EAMS22]: to treat

Comment [EAMS23]: , thus

Comment [EAMS24]: , thus

Comment [EAMS25]: reducing

Comment [EAMS26]: used ,

Comment [EAMS27]: most common

Pre-hospital antibiotic use refers to the practice of the use of antibiotics without prior formal consultation and prescription by a doctor. These antibiotics are either purchased over the counter or the left overs of a previous prescription. This practice is common in adults and even children both in developed and developing countries of the world with prevalence rates between 1% and 69%. [4-11] Surprisingly, more than 50% of antibiotics which are meant to

Comment [EAMS28]: , which

Comment [EAMS29]: its

Comment [EAMS30]: over

Comment [EAMS31]: using

Comment [EAMS32]:

Comment [EAMS33]: in both

Comment [EAMS34]:

be prescription drugs are purchased overthecounter, without **doctor's** prescription, from pharmacies, patent medicine stores and from even street vendors worldwide. [12] Pre-hospital use of antibiotics thus lack professional supervision leading to its inappropriate use or misuse. These drugs are either given unnecessarily, in over-dose or in under-dose. Various studies carried out in Jordan [13] and Europe [14] documented that pre-hospital antibiotics **given** were used for treating viral infections **of** which antibiotics are actually ineffective. [15] The integrated management of childhood illnesses (IMCI) discourages the use of antibiotics **to** children with infections of viral causes such **as** common cold and acute watery diarrhoea. Pre-hospital antibiotic use by parents on their children could have far-reaching consequences which span from side effects from the antibiotics to the promotion of the spread of antimicrobial resistance **which** could lead to treatment failure, deteriorating clinical conditions and even death. [16,17] It could also lead to unnecessary medical costs and delays in seeking proper medical care resulting in increased morbidity and mortality. The pre-hospital use of antibiotics has **therefore** become a major public health problem. [18] Various factors have been associated with pre-hospital use of antibiotics. These include **high** cost of medical consultation, long waiting **time** in hospitals, limited supply of medicines, unacceptable practices/behaviours of health care professionals, long distance to hospitals, high cost of transportation to health care facilities, numerous hospital **bottle necks**, lack of health insurance and poverty. [19,20] In addition, inadequate regulation of the distribution and sale of drugs **including antibiotics especially in low- and middle-income countries** account for this practice. [21] The type of pre-hospital antibiotics used varies **with** geographic location, age **as well as** the prevalent symptom(s). A cross-sectional population-based survey carried out in Indonesia [22] reported the commonest pre-hospital antibiotics as amoxicillin and ampicillin **whereas** in Uganda, [8] amoxicillin, erythromycin and metronidazole were the most common pre-

Comment [EAMS35]: a docors

Comment [EAMS36]:

Comment [EAMS37]: , of

Comment [EAMS38]: for

Comment [EAMS39]: as the

Comment [EAMS40]: , hreference,

Comment [EAMS41]:

Comment [EAMS42]: ,

Comment [EAMS43]: the high

Comment [EAMS44]: times

Comment [EAMS45]: bottlenecks

Comment [EAMS46]: , including antibiotics ,

Comment [EAMS47]: countries,

Comment [EAMS48]: depending on

Comment [EAMS49]: and

Comment [EAMS50]: , whereas

hospital antibiotics given to their febrile children. In Ibadan, [23] south-west Nigeria, ampicillin-cloxacillin drop was the commonest pre-hospital antibiotic given by mothers to their under-5 children followed by ampicillin and cefuroxime whereas an earlier study carried out in the same centre showed cotrimoxazole as the most common pre-hospital antibiotics followed by penicillins and cephalosporins. [9]

Comment [EAMS51]: the ampicillin-cloxacillin

There is a paucity of information in Nigeria regarding the pre-hospital use of antibiotics in children by mothers/caregivers and in Rivers State, no study has been carried out in this regard considering the fact that the rise in the use of pre-hospital antibiotics is now a major global health problem. [24] The present study was therefore carried out to ascertain the prevalence and pattern of pre-hospital antibiotic administration by mothers to their sick children attending the Paediatric Outpatient Clinic of Rivers State University Teaching Hospital(RSUTH) in south south Nigeria. Findings from this study will not only add to the body of knowledge in Nigeria and the world at large but would give useful information that will help formulate policies for educational interventions on the ills of pre-hospital antibiotic use which would inturn curb this menace.

Comment [EAMS52]: lack

Comment [EAMS53]: mothers

Comment [EAMS54]:

Comment [EAMS55]: southwestern

Comment [EAMS56]: will

Comment [EAMS57]: use,

Comment [EAMS58]: , in

Comment [EAMS59]: inturn,

Materials and Methods

It was a descriptive cross-sectional study design carried out over six months from January 1st to June 30th 2023, among mothers/caregivers attending the Paediatric outpatient clinic of the Rivers State University Teaching Hospital (RSUTH) in Port Harcourt, south-south Nigeria. The RSUTH, a state-owned tertiary hospital located in the Government reserved area of the State is a 375-bed hospital which receives referral from all the Primary Health Care centres and general hospitals in the 23 local government areas of the state as well as from private hospitals and neighbouring states. The hospital consists of both non-clinical and clinical departments.

Comment [EAMS60]: a referral

Comment [EAMS61]: neighbouring

The Paediatric department is one of the clinical departments of the hospital and consist of the Paediatric outpatient clinic, children emergency room, special care baby unit and the childrens' ward. The Paediatric outpatient clinic is open 5 days of the week, Mondays to Fridays from 8am to 4pm and consists of the specialist clinics and the general paediatric clinics. Each clinic day is run by two to three consultants, resident doctors, house officers, nurses of various cadres, other non- medical support staff and sees 42 patients on average each day. Health talks are usually given on each clinic day by the nurses before the commencement of each clinic day thereafter vital signs and anthropometric measurements are taken.

Comment [EAMS62]: hospital's clinical departments and consists

Comment [EAMS63]: five

Comment [EAMS64]:

Comment [EAMS65]: daily

Comment [EAMS66]: .

Comment [EAMS67]: Thereafter,

A research assistant was recruited for the study and was trained on the aim and objectives of the study, inclusion and exclusion criteria as well as the proper administration of the questionnaire to participants. A convenient sampling method was deployed for the study.

Comment [EAMS68]: study's aim and objectives, inclusion and exclusion criteria, and proper questionnaire administration to participants.

Ethical clearance was obtained from the Rivers State Hospitals Management Board Ethics Research Committee. Mothers/caregivers were duly educated about the research and a verbal consent whether or not to participate in the study was obtained.

Comment [EAMS69]: and caregiver

All mothers/caregivers whose children were within the age group one month to 16 years were consecutively recruited for the study. The inclusion criteria included all mothers/caregivers of children one month to 16 years irrespective of their symptoms who gave consent to participate in the study whereas mothers/caregivers who did not give consent to participate in the study were excluded.

Comment [EAMS70]: of one

Comment [EAMS71]: which

A pre-tested structured questionnaire developed by the researchers was administered by the researchers and/or research assistant to the mothers/caregivers. Data collected included biodata of the patients and mothers/caregivers as well as questions on the pattern of pre-hospital antibiotic administration to their children. Socio-economic class of the participants were determined using Oyedeji's classification. [25]

Comment [EAMS72]: and

Comment [EAMS73]: assistants

Comment [EAMS74]: the biodata

Comment [EAMS75]: and

Comment [EAMS76]: Socioeconomic

Data was entered into an Excel sheet and analysed using SPSS software version 23. Data was presented as percentages, tables and figures.

RESULTS

SOCIO-DEMOGRAPHIC CHARACTERISTICS

Table 1: Socio-demographic Characteristics of Child

Variable	Frequency (n = 401)	Percent
Sex		
Male	235	58.6
Female	166	41.4
Child Age Group (months)		
1 – 10	89	22.2
11 – 20	51	12.7
21 – 30	43	10.7
31 – 40	41	10.2
41 – 50	24	6.0
>50	153	38.2
Mean age: 28.26 ± 3.80 months		
Childs Weight Group (Kg)		
<4	12	3.0
4.1 – 14.0	176	43.9
14.1 – 24.0	121	30.2
24.1 – 34.0	32	8.0
>34.0	60	15.0
Mean weight: 14.94 ± 2.06 Kg		

Table 2: Parents' Socio-demographic Characteristics

Variable	Frequency (n = 401)	Percent
Mother's Age Group (years)		
<30	80	20.0
30 – 40	225	56.1
>40	96	23.9
Mean Age: 35.48 ± 1.26 years		
Mothers' Occupation		
Business/trader	166	41.4
Civil servant	59	14.7
Public Servant	58	14.5
Student	19	4.7
Hose wife/Unemployed	30	7.5
Artisan	33	8.2
Professional	36	9.0
Mothers' Level of Education		
Primary	5	1.2
Secondary	123	30.7
Tertiary	273	68.1
Father's Age Group (years)		
28 – 37	125	31.2

38 – 47	186	46.4
>47	90	22.4
Mean age: 42.11 ± 7.43 years		
Father's Occupation		
Business/ Trader	146	36.4
Civil servant	88	21.9
Public servant	34	8.5
Professional	52	13.0
Artisan	40	10.0
Unemployed	6	1.5
Private company employee	35	8.7
Level of Education		
Primary	4	1.0
Secondary	111	27.7
Tertiary	286	71.3
Socioeconomic Class (OYEDEJI)		
Class 1	102	25.5
Class 2	195	48.6
Class 3	86	21.5
Class 4	17	4.2
Class 5	1	0.2
Socioeconomic Class (OLUSANYA)		
Class 1	274	68.4
Class 2	126	31.4
Class 3	1	0.2

Table 3: History of Antibiotics Usage

Variable	Frequency (n = 401)	Percent
Administered Antibiotics		
Yes	89	22.2
No	312	77.8
Reason for Antibiotics (Multiple response, n = 147)		
Rash	8	9.3
Pain	11	12.8
Catarrh	13	15.1
Cough	42	48.8
Vomit	7	8.1
Fever	54	62.8
Diarrhoea	12	14.0
Type of Antibiotics formulation (n = 162)		
Tablet	18	20.2
Syrup	62	69.7
Capsule	8	9.0
Injection	1	1.1
Name of Antibiotic Administered (n =		

89)		
Amoxyl/Augmentin	44	49.5
Azithromicin	5	5.6
Oral Cephalosporins	21	23.5
Inj Ceftriaxone	1	1.1
Septrin	4	4.5
Ampiclox	7	7.9
Erythromycin	2	2.2
Ciprofloxacin	2	2.2
Ampicilin	3	3.3

Table 4: Reason for Choice of Formulation

Variable	Frequency (n = 211)	Percent
Reason for Choice of Formulation		
Age of child	154	73.0
Available medication	26	12.3
Most effective	14	6.6
Prescribed	8	3.8
Previous Experience	5	2.4
Other reasons	4	1.9
Other reasons (n = 4)		
Weight of Child	1	25.0
No reason	2	50.0
Has not administered drug to child before	1	25.0




Figure 1: Reason for Choice of Formulation Used

Table 5: Duration of Medication

Variable	Frequency (n =84)	Percent
----------	--------------------	---------

Duration of Antibiotic Medication (n = 84)		
< 7 days	64	76.2
≥ 7 days	20	23.8

Table 6: Person Responsible for Prescription

Person Responsible for Prescription	Frequency (multiple responses n = 199)	Percent
Doctor	50	25.1
Pharmacist	42	21.1
Chemist	14	7.0
Self-Experience	72	36.2
Nurse	17	8.5
others	4	2.0

Table 7: appropriate antibiotic dosing

Appropriate Antibiotic Dose (n = 71)		
Yes	44	62.0
Under-dose	23	32.4
Overdose	4	5.6

Discussion

Majority of those recruited into the study were males (58.6%) similar to the report from Uganda where males made up 56.2% of the participants.[26] The prevalence of pre-hospital administration of antibiotics in the study was 22.2% which is lesser than the 39.5% reported in Uganda [8], 40.1% reported in Kenya[26], 46.7% reported in Enugu[27] and Yemen[28]. The study in Uganda [8] and Kenya [26] were among children below 5 years of age while ours involved all children. Children less than 5 years have lower immunity and are more likely to present with acute infections for which parents will give antibiotics. The study by Ekwuochi et al in Enugu[27] was among children with diarrhoea, a symptom which many Gignorantly think should be treated with antibiotics. In Yemen, the study was much earlier at the turn of the century and may have reduced now. The high rate of antibiotic use before presentation in the hospital is likely due to the ease of procurement of antibiotics over the

Comment [EAMS77]: ,and

Comment [EAMS78]: five

Comment [EAMS79]: , while

Comment [EAMS80]: five

Comment [EAMS81]: years old

Comment [EAMS82]: was conducted

Comment [EAMS83]:

Comment [EAMS84]: in Yemen at

counter in Nigeria and other low income countries. The lesser prevalence in our study could be due to the present economic hardship whereby parents can not afford to buy the medications on their own hence waiting to get a proper prescription from the hospital.

Comment [EAMS85]: and hence

Comment [EAMS86]: wait

The most common antibiotics used in this study were amoxicillin/amoxicillin -clavulacin acid, oral cephalosporins and Ampicillin -cloxacacin combination which was similar to the report from Uganda[8], where the most commonly used antibiotic was amoxicillin 33/83 (39.8%), followed by erythromycin 18 (21.7%), metronidazole 14 (16.9%), ciprofloxacin 13 (15.7%) and ampicillin 6 (7.2%). The report from Yemen [28] identified trimethoprin sulfamethoxazole and amoxicillin- clavulanic acid as the most commonly prescribed antibiotics.

The most common reasons for the use of antibiotics in these children included fever, cough and catarrh while diarrhoea, rash and vomiting were the least reasons for pre-hospital were

Comment [EAMS87]: using

Comment [EAMS88]: ,

the least reasons. This is similar to the report from Uganda[8] where children with fever also had catarrh, cough, diarrhoea and vomiting necessitating a prehospital antibiotics therapy.

Comment [EAMS89]: reasons

Comment [EAMS90]: [8] ,

Comment [EAMS91]: ,

The above symptoms in children are often viral in nature and do not usually require antibiotics therapy however, ignorance may be a factor in making parents give antibiotics unnecessarily.

Comment [EAMS92]:

Comment [EAMS93]:

Comment [EAMS94]: antibiotic

Among those that received antibiotics, 62% of them received an appropriate dose while 32.4% received underdose and 5.6% got an overdose of administered medication.

Comment [EAMS95]: who

Comment [EAMS96]: ,

Comment [EAMS97]: an underdose ,

Inappropriate dose may worsen the present antimicrobial resistance

Comment [EAMS98]: .

Conclusion

Pre-hospital antibiotics use and abuse/misuse is a common practice among the respondents.

Comment [EAMS99]: Prehospital

Most of the drugs were procured from pharmacy shops and chemist shops without a prescription. Intervention by the different ministries of health through education of the

Comment [EAMS100]: antibiotic

populace to curb the menace of drug abuse with its attendant development of drug resistance as well as putting mercenaries in place to prosecute those selling and buying OTC drugs. The practice should be prohibited. This public enlightenment can be carried out by the different ministries of health.

Comment [EAMS101]:

Comment [EAMS102]: he different ministries of health can carry out this public enlightenment.

References

1. WHO 2008. World Health Statistics 2008. Mortality and burden of disease. Geneva: WHO 36-64
2. World Health Organization. WHO Policy Perspectives on Medicines-Promoting rational use of medicines: core components. Geneva: World Health Organization; 2002. Available at: https://apps.who.int/iris/bitstream/handle/10665/67438/WHO_EDM_2002.3.pdf
3. Kardas P, Devine S, Golembesky A, Roberts C. A systematic review and meta-analysis of misuse of antibiotic therapies in the community. Int J Antimicrob Agents 2005; 26: 106-13
4. Knopf H, Wolf IK, Sarganas G, Zhuang W, Rascher W, Neubert A. Off-label medicine use in children and adolescents: Results of a population-based study in Germany. BMC Public Health 2013; 13: 631

Comment [EAMS103]: Please review the web or remove

5. Schillie SF, Shehab N, Thomas KE, Budnitz DS. Medication overdose leading to emergency department visits among children. *Am J Prev Med* 2009; 37: 181-7 doi: 10.1016/j.amepre.2009.05.018
6. Maltezou HC, Dedoukou X, Asimaki H, Kontou I, Ioannidou L, Mitromara K, Theodoridou K, Katerelos P, Theodoridou M. Consumption of antibiotics by children in Greece: A cross-sectional study. *Int JPediatr Adolesc Med* 2017; 4: 108-11 doi: 10.1016/j.ijpam.2017.04.002
7. Jasim A. Parental self medication of antibiotics for children in Bagdad city. *Int J Pharm Pharm Sci* 2014; 6: 485-9
8. Nyeko R, Otim F, Obiya EM, Abala C. Pre-hospital exposures to antibiotics among children presenting with fever in Northern Uganda: a facility-based cross-sectional study. *BMC Pediatr* 2022; 22: 322 doi: 10.1186/s12887-022-03375-2
9. Fehintola FA. Pre-hospital and prescription use of antibacterial drugs at a secondary health centre in Ibadan, Nigeria. *Afr J Pharm Pharm* 2009; 3(4): 120-3
10. Richman PB, Garra G, Eskin B, Nashed AH, Cody R. Oral antibiotic use without consulting a physician: a survey of ED patients. *Am J Emerg Med* 2001; 19(1): 57-60
11. Al-Ayed MSZ. Parents' knowledge, attitudes and practices on antibiotic use by children. *Saudi J Med Med Sci* 2019; 7: 93 doi: 10.4103/sjmms.sjmms_171_17
12. Cars O, Hogberg L. Innovating for antibacterial resistance.
13. Al-Azzam SI, Al-Husein BA, Alzoubi F, Masadeh MM, Al-Horani MA: Self-medication with antibiotics in Jordanian population. *Int J Occup Med Environ HEALTH* 2007; 20(4): 373-80
14. Grigoryan L, Burgerhof JG, Degener JE, Deschepper R, Lundborg CS, Monnet DL, Scicluna EA, Birkin J, Haaijer-Ruskamp FM. Attitudes, beliefs and knowledge

- concerning antibiotic use and self-medication: a comparative European study. *Pharmacoepidemiol Drug Saf* 2007; 16(11): 1234-43
15. Shimizu Y. Stop overuse and misuse of antibiotics: combat resistance. World Health Organization 10 November 2017. Available at: who.int/westernpacific/news/item/10-11-2017-stop-overuse-and-misuse-of-antibiotics-combat-resistance Cited 05 March 2024
 16. Goosesens H. Antibiotic consumption and link to resistance. *Clin Microbiol Infect* 2009; 15: 12-5 doi: 10.1111/j.1469-0691.2009.02725.x
 17. Simonsen GS, Tapsall JW, Allegranzi B, Talbot EA, Lazzari S. The antimicrobial resistance containment and surveillance approach – a public health tool, *Bulletin of the World Health Organization* 2004; 82(12): 928-34
 18. Togoobaatar G, Ikeda N, Ali M, et al. Survey of non-prescribed use of antibiotics for children in an urban community in Mongolia. *Bulletin of the World Health Organization* 2010; 88(12): 930-6
 19. Hadi U, Duerink DO, Lestari ES, Nagelkerke NJ, Werter S, Keuter M, Suwandojo E, Rahardjo E, Broek PVD, Gyssens IC. Survey of antibiotic use of individuals visiting public healthcare facilities in Indonesia. *Int J Infect Dis* 2008; 12: 622-9
 20. Saradamma RD, Higginbotham N, Nichterc M. Social factors influencing the acquisition of antibiotics without prescription in Kerala state, south India. *Soc Sci Med* 2000; 50: 891-903
 21. Bin Abdulhak AA, Al Tannir MA, Almansor MA, Almohaya MS, Onazi AS, Marei MA, Aldossary OF, Obeidat SA, Obeidat MA, Raiz MS, Tleyjeh IM. Non-prescribed sale of antibiotics in Riyadh, Saudi Arabia: A cross sectional study. *BMC Public Health* 2011; 11: 538-42

22. Widayah A, Suryawati S, de Crespigny C, Hiller JE. Self-medication with antibiotics in Yogyakarta city, Indonesia: A cross sectional population based survey. *BMC Res Notes* 2011; 4: 491 doi: 10.1186/1756-0500-4-491
23. Adisa R, Orherhe O, Fakeye TO. Evaluation of antibiotic prescriptions and use in under-five children in Ibadan, south western Nigeria. *Afr Health Sci* 2018; 18(4): 1189-201
24. Al Rasheed A, Agoub U, Alkhashan H, Abdelhay O, Alawwad A, Al About A, Al Battal S. Prevalence and predictors of self-medication with antibiotics in Al Wazarat Health Center, Riyadh city, KSA. *BioMed Res Int* 2016;8. 10.1155/2016/3916874
25. Oyedeji GA. Socioeconomic and cultural background of hospitalized children in Ilesa. *Niger J Paediatr* 1985; 12: 111-7
26. Al Rasheed A, agoub U, Alkhashan H, Abdelhay O, Alawwad A, Al About A, Al Battal S: Prevalence and Predictors of Self-Medication with Antibiotics in Al Wazarat Health Center, Riyadh City, KSA. *BioMed Res Int.* 2016:8 (kenya study)
27. Uchenna Ekwochi, Josephat M. Chinawa, Ikechukwu Obi, Herbert A. Obu, Samuel Agwu, Use and/or Misuse of Antibiotics in Management of Diarrhea Among Children in Enugu, Southeast Nigeria, *Journal of Tropical Pediatrics*, Volume 59, Issue 4, August 2013, Pages 314–316
28. Mohanna M. Self-Medication with antibiotics in children in Sana'a City, Yemen. *Oman Med J.* 2010; 25: 41-43.