

***Balantidium Coli Colitis* Revealing a Common Variable Immunodeficiency: A Case Report**

Formatted: Font: Italic

Formatted: Tab stops: 3.15", Centered

Abstract:

Balantidium Coli is a large protozoan parasite, measuring up to 150 micrometers, initially identified in 1857 and later renamed in 1861-1862. It is an animal parasite, with its primary reservoir being pigs (domestic pigs/wild boars), but it can also be found in other animal species such as dogs, cats, hyenas, buffaloes, camels, monkeys, and others. Humans are an accidental host, as they represent a dead-end in the parasitic life cycle. Infection by this parasite is typically asymptomatic or presents with mild symptoms. It becomes pathogenic only in cases of primary or secondary immunodeficiency, where it typically manifests as colitis; however, urinary tract involvement is also possible. In our case, *Balantidium Coli* infection presented multiple challenges. It is not a well-known parasite in our region, particularly since pig farming is not practiced in Morocco. The parasite was found in both the urine and stool of the patient, who exhibited clear signs of infection. This led to the "late" discovery of Common Variable Immunodeficiency (CVID) in the patient, a primary immunodeficiency typically diagnosed before the age of 4 and characterized by a deficiency in immunoglobulin production, particularly affecting the IgA and IgG fractions.

Formatted: Justified

Comment [PF1]: Show some brief global, regional burden of the parasite

Introduction:

Balantidium Coli is a protozoan parasite measuring about 150 micrometers, with pigs as its primary reservoir. Humans are accidental hosts, and contamination can occur through direct contact by ingesting the parasite's cyst via dirty hands or indirectly by consuming contaminated food or water. It is the largest pathogenic protozoan in humans and is responsible for a zoonosis [1]. It typically causes colitis, first described and identified by Malmsten in 1857 in two patients presenting with dysentery. It was initially named *Paramoecium* then and was later renamed *Balantidium* by Leuckart in 1861 and Stein in 1862 [2].

Formatted: Justified

Common Variable Immunodeficiency (CVID) is a primary immunodeficiency characterized by a deficiency in the production of immunoglobulin G and immunoglobulin A, leading to hypogammaglobulinemia. It is usually diagnosed before the age of 4, following recurrent infections, particularly of the respiratory tract [3]. We report the case of a patient in whom *Balantidium Coli* colitis revealed a Common Variable Immunodeficiency.

Clinical Case:

The patient was a 23-year-old woman, born from a first-degree consanguineous marriage. She had a history of recurrent respiratory and digestive infections (more than three episodes per year) that responded to probabilistic antibiotic therapy (amoxicillin + clavulanic acid) since early childhood. She also had chronic iron deficiency anemia since the age of 14,

Formatted: Justified

treated with iron supplementation (80 mg/day) without improvement in ferritin levels. Additionally, she had a history of *Helicobacter pylori* gastritis diagnosed at age 19, treated with amoxicillin 1 g twice daily, clarithromycin 500 mg twice daily, in combination with omeprazole 20 mg twice daily for 15 days. She had an appendectomy at age 12 and primary amenorrhea. The patient received three doses of the Sinopharm COVID-19 vaccine.

The illness began six months before her hospitalization, with the onset of diffuse abdominal pain in the colonic region, associated with mucus-bloody diarrhea, occurring 6 to 8 times per day, all evolving in a febrile context with weight loss of 18 kg over six months. The condition progressed with the appearance of vomiting, lumbar pain, hematuria, and frequent urination, prompting the patient to seek emergency care at the Ibn Sina University Hospital Center in Rabat...[which country?](#)

On clinical examination, the patient was found to be cachectic with growth retardation, measuring 1.42 meters tall and weighing 27 kg, with a body mass index of 13.4. She was dehydrated, with a skin fold persisting for more than two seconds. The patient was somnolent and confused, with a Glasgow Coma Score of 13/15, hypotensive at 91/62 mmHg, tachycardic at 100 beats per minute, and tachypneic at 27 breaths per minute, with pale conjunctiva and a temperature of 38.7°C. Pulmonary examination revealed rhonchi, and abdominal examination revealed a palpable spleen tip with minimal splenomegaly (15 cm liver span) and tenderness in the colonic region without clear guarding or rigidity, with an appendectomy scar in the right iliac fossa. On rectal examination, the glove was covered with mucus streaked with blood.

The patient was urgently hospitalized for an etiological workup and appropriate management. The patient's laboratory workup revealed hypochromic microcytic anemia with a hemoglobin level of 6 g/dL, moderate leukocytosis at 11,000/mm³, predominantly neutrophils at 8,500/mm³, with moderate eosinophilia at 620/mm³, and normal lymphocytes at 1,580/mm³ with mild thrombocytosis at 480,000/mm³. Coagulation tests were normal. The inflammatory workup showed an elevated C-reactive protein (CRP) at 98 mg/L, with fibrinogen at 5.6 g/L. Procalcitonin was negative at 0.11 ng/L. The electrolyte panel showed hyponatremia at 124 mmol/L and hypokalemia at 2.8 mmol/L. Ferritin was severely depleted at 3 µg/L, total protein was decreased at 57 g/L, and hypoalbuminemia was noted at 23 g/L, with fasting blood glucose at 0.58 g/L and triglycerides at 0.28 g/L.

The patient thus presented with three major problems:

- **Severe dehydration and malnutrition with probable malabsorption.**
- **A probable digestive and/or urinary infection, given the fever, digestive, and urinary symptoms, potentially due to an intracellular bacterium, virus, or parasite (given the eosinophilia); a combination of the three is also possible.**
- **An immunosuppressive condition that could explain the recurrent infections as well as the profound deterioration in the general state, such as an acquired or primary immunodeficiency, a chronic inflammatory or autoimmune disease, an underlying neoplastic condition, or a metabolic disorder like cystic fibrosis.**

Our priority was to rehydrate the patient and correct the electrolyte disturbances, alongside antibiotic therapy with intravenous metronidazole 500 mg three times a day. The patient

underwent a QuantiFERON test, which came back negative, and a urine cytobacteriological examination revealed leukocyturia at 130,000/mL and hematuria at 24,000/mL without acanthocytes, with cystic protozoa present in the culture. Fecal calprotectin was strongly positive at 180 µg/g of stool, and the stool culture was sterile, while three parasitological stool examinations isolated ciliated protozoa measuring between 80 and 120 µm, suggesting *Balantidium coli*. Given this unusual finding in our context, PCR confirmation was performed on urine and stool, which came back positive.

Since *Balantidium coli* is rarely pathogenic in humans and is thus considered an opportunistic pathogen, an underlying immunosuppressive condition was investigated. HIV 1 and 2 serology was performed and was negative, which was expected since an opportunistic infection would occur at stages B or C (AIDS), and the patient did not present with lymphopenia. Cystic fibrosis was investigated due to the early onset of recurrent respiratory and digestive infections, with [a normal sweat-chloride test \(32mmol/L\) coming back normal at 32 mmol/L](#), and [a negative the search for the](#) CFTR gene mutation [was negative](#). Inflammatory Bowel Disease (IBD) was considered, although the patient had no erythema nodosum, oral or nasal mucosal ulcerations, anal fissures, or arthralgias; ANCA and ASCA tests [wereing was](#) negative, the enteroscanner showed no abnormalities, upper gastrointestinal endoscopy revealed non-specific antrum-fundus gastritis with [a positive Helicobacter pylori](#) [positive](#) without ulcerations or signs of malignancy, while rectosigmoidoscopy was unremarkable, and total ileocolonoscopy was normal. The histopathological study of biopsies revealed chronic interstitial ileitis in acute vasculo-exudative flare without specificity or malignancy signs, and non-specific interstitial colitis without malignancy signs. The uroscan was also unremarkable.

The possibility of primary immunodeficiency was then considered. Serum protein electrophoresis revealed hypoalbuminemia at 23 g/L and profound hypogammaglobulinemia at 2.4 g/L. Lymphocyte immunophenotyping showed no abnormalities, and 24-hour proteinuria was negative at 0.11 g/24h. Quantitative measurements of immunoglobulins revealed severely reduced IgG at 1.86 g/L and IgA at 0.34 g/L, while IgM levels were normal. The diagnosis of Common Variable Immunodeficiency (CVID) was therefore established according to [revised international \(PAGID\) criteria in 2016](#). The patient was placed on therapeutic human immunoglobulin infusion at 0.8 g/kg during the first course, followed by 0.4/kg/month. For [herthe](#) *Balantidium coli* infection, she was treated with doxycycline 400 mg/day on the first day, then 200 mg/day for 21 days, with good clinical and biological improvement.

Discussion:

Our clinical case presents several atypical features that are worth highlighting. *Balantidium coli* is a large protozoan parasite, approximately 150 µm in size, commonly found in tropical regions, with higher prevalence in Asia. The typical reservoir is pigs, but it can also be found in other species such as dogs, cats, rats, hyenas, and monkeys [1-4]. Humans are an accidental host, with transmission occurring either directly through ingestion of the parasite cyst or indirectly through consumption of contaminated food or water [1]. The individuals most frequently affected are pig farmers and those in close contact with them, as demonstrated in a study conducted in Ghana [5].

Comment [PF2]: Refence this

Comment [PF3]: Repeated severally

This ~~brings is the us to the~~ first atypical case aspect in the Moroccan context: this parasite ~~is has~~ not been known to be present here, Moreover, there is no and we do not have pig farming since pigs are the common host where the parasite is commonly found [6-7]. On ~~Upon~~ further investigation, it was -we discovered that the patient came from the Sekhirate region, residing in a small village without running water, which is known for the presence of wild boars that could potentially carry the parasite. A review of the literature also revealed that cattle can be affected, as shown in a study on a species of buffalo in Thailand [8], as well as ostriches [4] and camels [9]. This therefore suggests the need for an epidemiological investigation to conduct large-scale screening in various cattle and camel farms, screening of wild boar populations, and exploration of the groundwater for possible contamination.

It is important to note that different direct examination techniques can be used to detect *Balantidium coli*, with the best being Lugol's iodine staining [10]. However, direct examination has limitations, However which is why PCR has demonstrated greater accuracy [1, 11].

The second atypical aspect is the presence of the protozoan in the patient's urine. *Balantidium coli* is generally asymptomatic or even harmless to humans, but it can become pathogenic in immunocompromised patients, where it typically causes digestive symptoms, primarily colitis [1-12]. However, in our patient, the parasite was not only found in the urine but also showed significant clinical and biological impact. This prompted us to search for similar cases in the literature, where we found two reports: the first involving a pregnant woman in Ethiopia [13], and the second in an Iranian patient with bladder cancer [14]. Notably, all three cases involved women, even though the carriage of *Balantidium coli* shows a slight male predominance according to the Ghana study [5]. It appears that urinary localization is more common in females.

The third atypical aspect is the late diagnosis of Common Variable Immunodeficiency (CVID), which is the most common primary immunodeficiency disorder. This condition is more familiar to pediatricians since it is generally diagnosed before the age of four [3]. This delay in diagnosis can be attributed to the limited access to healthcare, due in part to the significant shortage of doctors in Morocco, as well as the lessack of awareness about CVID among general practitioners, highlighting the need for specialized training.

Conclusion:

Balantidium coli is an animal protozoan parasite, primarily hosted by pigs but can also be found in other species. Humans serve as an accidental host. This parasite is rarely pathogenic in humans, requiring an immunosuppressive condition to become harmful, which classifies it as an opportunistic pathogen. When pathogenic, Balantidium coli typically causes colitis but can also have urological manifestations. It is importantnt The key takeaway is always to consider an underlying immunosuppressive condition, whether primary or acquired, as this parasitosis may be indicative of such a state.

Comment [PF4]: Repeated !

Comment [PF5]: BTW, what was the underlying immunosuppression condition of this patient?

References:

[1]- Microbiology of Waterborne Diseases (Second Edition), Microbiological Aspects and Risks 2014, Pages 277-286, Microbiology of Waterborne Diseases, Chapter Fifteen -

Balantidium coli, Rachel M. Chalmers, Head of Cryptosporidium Reference Unit, Public Health Wales, Singleton Hospital, Swansea, UK, <https://doi.org/10.1016/B978-0-12-415846-7.00015-9>

[2]- Balantidiasis, Updated: Aug 25, 2023, Author: Subhash Chandra Parija, MD, MBBS, PhD, DSc, FRCPath; Chief Editor: Pranatharthi Haran Chandrasekar, MBBS, MD, <https://emedicine.medscape.com/article/213077-overview?form=fpf>

[3]- Revue Francophone des Laboratoires Volume 2023, Issue 549, February 2023, Pages 38-46, Les déficits immunitaires communs variables : définition, physiopathologie et diagnostic biologique Common variable immunodeficiency, definition, frequency and diagnosis Emma Chambrey a, Marie-Nathalie Kolopp-Sarda a b, Christophe Malcus c, [https://doi.org/10.1016/S1773-035X\(23\)00025-4](https://doi.org/10.1016/S1773-035X(23)00025-4)

[4]- Source: National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Parasitic Diseases and Malaria, Last Reviewed: June 6, 2019, <https://www.cdc.gov/dpdx/balantidiasis/index.html>

[5]- Aninagyei et al. BMC Infect Dis (2021) 21:1047, RESEARCH ARTICLE Prevalence and risk factors of human Balantidium coli infection and its association with haematological and biochemical parameters in Ga West Municipality, Ghana Enoch Aninagyei^{1*}, Salifu Nanga², Desmond Omame Acheampong³, Rita Mensah⁴, Mercy Nelly Boadu⁵, Henrietta Terko Kwansa⁶, Bentum¹ and Clement Okraku Tettey¹, <https://doi.org/10.1186/s12879-021-06731-2>

[6]- Veterinary World, EISSN: 2231-0916 Available at www.veterinaryworld.org/Vol.14/April-2021/33.pdf RESEARCH ARTICLE Open Access Prevalence of Balantidium coli (Malmsten, 1857) infection in swine reared in South Italy: A widespread neglected zoonosis Filippo Giarratana¹, Luca Nalbone¹, Ettore Napoli¹, Vincenzo Lanzo² and Antonio Panebianco¹ 1. Department of Veterinary Science, University of Messina, Polo Universitario dell'Annunziata, 98168 Messina, Italy; 2. Freelance Veterinary Professional, 89024 Polistena (RC), Italy. Corresponding author: Luca Nalbone,

[7]- Pathogens Article Prevalence of Swine Gastrointestinal Parasites in Two Free-Range Farms from Nord-West Region of Romania Mihai-Horia Băieș¹, Zsolt Boros¹, Călin Mircea Gherman^{1,*}, Marina Spînu², Attila Mathe³, Stefan Pataky³, Menelaos Lefkaditis⁴ and Vasile Cozma^{1,5} 1 Department of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Mănăștur Street 3-5, 400372 Cluj-Napoca, Romania 2 Department of Infectious Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Mănăștur Street 3-5, 400372 Cluj-Napoca, Romania 3 Agricultural Research and Development Station of Turda, Agriculturii Street 27, 401100 Turda, Romania 4 Laboratory of Microbiology and Parasitology, Department of Veterinary Medicine, School of Health Sciences, University of Thessaly, 43100 Karditsa, Greece 5 Academy of Agricultural and Forestry Sciences Gheorghe Ionescu-Sisești (A.S.A.S.), Mărășești Boulevard 61, 011464 Bucharest, Romania, <https://doi.org/10.3390/pathogens11090954>

- [8]- Veterinary World, EISSN: 2231-0916 Available at www.veterinaryworld.org/Vol.13/August-2020/8.pdf RESEARCH ARTICLE Open Access Gastrointestinal parasite infections in fighting bulls in South Thailand Domechai Kaewnoi, Ratchakul Wiriyaprom, Saowakon Indoung and Ruttayaporn Ngasaman Faculty of Veterinary Science, Prince of Songkla University, Hatyai, Songkhla, Thailand. Corresponding author: Ruttayaporn Ngasaman
- [9]- Sazmand et al. Parasites Vectors (2019) 12:610 <https://doi.org/10.1186/s13071-019-3863-3> REVIEW Zoonotic parasites of dromedary camels: so important, so ignored Alireza Sazmand1* , Anja Joachim2 and Domenico Otranto1,3
- [10]- Veterinary World, EISSN: 2231-0916 Available at www.veterinaryworld.org/Vol.14/April-2021/9.pdf RESEARCH ARTICLE Open Access Comparison between five coprological methods for the diagnosis of *Balantidium coli* cysts in fecal samples from pigs Juan Carlos Pinilla1 , Andrea Isabel Pinilla2 and Angel Alberto Florez1 1. Department of Veterinary Medicine, Faculty of Exact, Natural and Agricultural Sciences, University of Santander, Bucaramanga, Colombia; 2. Department of Microbiology, Faculty of Health, Industrial University of Santander, Bucaramanga, Colombia.
- [11]- Identification of Zoonotic *Balantidium coli* in Pigs by Polymerase Chain Reaction-Restriction Fragment Length Polymorphism (PCR-RFLP) and Its Distribution in Korea by Jae-Won Byun 1, Jung-Hyun Park 2, Bo-Youn Moon 1, Kichan Lee 1, Wan-Kyu Lee 2 ORCID, Dongmi Kwak 3 ORCID and Seung-Hun Lee 2,
- [12]- Korean J Parasitol Vol. 58, No. 1: 47-49, February 2020 ☐ CASE REPORT <https://doi.org/10.3347/kjp.2020.58.1.47> •Received 11 December 2019, revised 6 February 2020, accepted 12 February 2020.) © 2020, Korean Society for Parasitology and Tropical Medicine This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Dysentery Caused by *Balantidium coli* in China Peixia Yu*, JianRong Rong, Yan Zhang, Jingjing Du Department of Clinical Laboratory Medicine, Shanxi Bethune Hospital & Shanxi Academy of Medical Sciences, Taiyuan, 030009, Shanxi, PR China
- [13]- Almaw A, Berhan A, Solomon Y, Malkamu B, Eyayu T, Workineh L, Mekete G, Yayehrad AT. *Balantidium coli*; Rare and Accidental Finding in the Urine of Pregnant Woman: Case Report. Int Med Case Rep J. 2022 Mar 22;15:105-109. doi: 10.2147/IMCRJ.S355536. PMID: 35345500; PMCID: PMC8957295.
- [14]- DOI: 10.1002/ccr3.7494 CASE REPORT Uninvited guest, *Balantidium coli* in urine in a patient with bladder cancer: A case report and review of the literature Farnaz Farmani1 | Neda Soleimani1,2 | Mohammad Razeghi1 | Amir Zamani1 | Sahand Mohammadzadeh1,2 | Davoud Soleimani3