

# HIIT IMPACTS THE BDNF AND PHOSPHORILATED TAU IN ALZHEIMER'S DISEASE CORRELATED WITH BLOOD LACTATE: A RANDOMIZED CONTROL TRIAL

**Aims:** To examine the effects of high-intensity body weight interval training (PMED) on brain-derived neurotrophic factor (BDNF), and phosphorylated-TAU protein (p-TAU) in Alzheimer's disease.

**Study design:** An experimental randomized control trial.

**Place and Duration of Study:** The recruitments were made in Vila Vicentina Community Center (CTIV), the Elderly Support Center of the University of Pará (UEPA), the Senior Living Center (CCI) and SESC Porto Velho, Brazil between 2016 and 2018.

**Methodology:** Seventy-eight seniors diagnosed with AD were divided into three groups (n=29): GCP (High-Intensity Interval Training) MICT (moderate-intensity continuous training), and GCS a sedentary control group. The HIIT group engaged in 30 minutes of exercise at 80-90% of maximal heart rate, while the MICT group walked for 30 minutes at 60% of maximal heart rate. Both exercise groups performed their respective activities three times per week for three months, composing 36 exercise sessions. Pre- and post-intervention measurements of BDNF and p-TAU were conducted.

**Results:** The GCS group showed no changes in neurochemical factors ( $p > 0.05$ ). In contrast, the HIIT group experienced increased BDNF and decreased p-TAU concentrations in the blood ( $p < 0.0001$  for both). The MICT group also showed similar improvements ( $p < 0.01$  for both). Between the experimental groups the HIIT had a higher effect on BDNF ( $p < 0.01$ ) and p-TAU ( $p < 0.01$ ) than MICT. In both exercise groups, blood lactate concentration had a strong correlation with BDNF and p-TAU levels ( $p < 0.01$ ).

**Conclusion:** Exercise increased BDNF and reduced p-TAU concentrations, improving neurochemistry in AD. Additionally, the intensity of exercise plays a significant role in the interaction between muscle and brain.

**Keywords:** *Aging. Exercise. Mental Health. Neurophysiology, Neuroscience*

## 1. INTRODUCTION

No human brain disorder has received more attention in recent years than AD disease (AD). AD is characterized by memory impairment, cognitive decline, and progressive dementia with altered neurochemical markers [1]. Although the role of inflammation in AD is not fully understood, exploring the connection between inflammatory factors, and myokines with the disease is important.

Muscle activity, especially during intense physical exercise, plays a key role due to its neuroendocrine regulation [2,3]. Myokines, substances released by muscles after vigorous exercise, such as lactate, interleukin-6, and irisin [4,5], are linked to brain plasticity and function. These substances are produced from minutes to days after intense exercise and have significant effects on brain [3].

The pathophysiology of AD involves a reduction in cerebral cortex choline acetyltransferase, neurotransmitters, and neuromodulators, along with changes in their synthesizing enzymes, modifications in beta-amyloid, and alterations in neuronal morphology (Cummings et al., 2012). Reduced production of brain-derived neurotrophic factor (BDNF) and increased production of p-TAU [6] are linked to neuron and axon death in AD due to the hyperphosphorylation of TAU protein [7].

Physical activity is one of the few effective interventions for neurodegenerative diseases and other cognition- and memory-related problems, including AD [8]. Many studies reported that physical activity can counteract the neurobiological effects of AD. For example, reduced functional decline in the hippocampus [9], and improved mitochondrial health in the brain [10], both associated with improvements in many aspects of the brain as his electrophysiology, astrocyte remodeling in the long-term exercise exposition [11] could be benefited to the physical exercise. However, the intensity of the exercise intervention can make a significant difference due to the release of lactate and myokines like IL-6 and irisin [12]. For example, intense exercise can improve mitochondrial health in the brain and lead to astrocyte remodeling with long-term exposure [11]. These changes can result in important cognitive benefits, justifying the use of this approach in our study.

Previously, our group displayed that the HIIT is able to improve the cognition pf healthy adults [13], for frailty older adults [14], the glycemic control of older adults with metabolism impairments [15], and in the cognition and dementia symptoms of frailty older adults [16]. In conjunct, this allows us to explore more about the effect of HIIT in neurodegenerative impairments related to AD.

Since PMED has shown positive results in treating memory-related, cognitive, and mental illnesses more effectively than moderate or low-intensity physical exercise, this study aims to provide insights beyond traditional aerobic or resistance training. It is crucial to explore whether the intensity and type of exercise affect the efficacy of physical exercise, as this approach is poorly explored in the literature regarding the neurobiological impairments caused by AD. This study specifically aims to examine the effects of PMED on brain-derived neurotrophic factor BDNF and p-TAU in AD.

## **2. MATERIAL AND METHODS**

### **2.1 2.1 Experimental Approach to The Problem**

For determine the effects of physical exercise, and non-invasive brain stimulation, both in electrophysiological behavior of the brain, the executive functions, and dementia symptoms in older adults with AD we planning two exercise approach to stimuli two levels of blood lactate concentration in response to the exercise. We controlled the exercise intensity across the heart rate, and lactate. Then, we measured the BDNF and p-TAU, and performed the correlation test to determine possible influence of the this myokine and the brain chemistry.

### **2.2 Study Type, and Participants and Ethical Statement**

The study involved 1096 elderly patients with AD from several centers: Vila Vicentina Community Center (CTIV), Elderly Support Center of the University of Pará (UEPA), Senior Living Center (CCI), and SESC Puerto Velho, Brazil. However, only 88 elderly patients met all the inclusion and exclusion criteria. Recruitment was conducted by the National Institute

for Disability Research (NINCDS) Recruitment Center until completion of the Participation Diagnosis. Participants were randomly assigned to groups using computer-based randomization with BioStat 5.3 software.

### **2.3 Procedures for inclusion and exclusion**

Nurses, trained in caring for patients with mental and exacerbating illnesses, referred older patients to the study. The diagnosis of AD followed the NINCDS criteria [17] and was confirmed by the Brazilian health system. To be included, subjects needed to score between 9 and 10 on the Edmonton Frailty Scale (EFS) [18]. Exclusion criteria included depression, severe sensory deficits, or other neurological diseases. Ineligible individuals included those unable to complete tasks due to physical limitations, incapable of understanding the task content or participating in research, or unwilling to sign the informed consent form.

Following the selection process, 34 older adults with AD of both sexes comprised the final sample, randomly assigned to two groups ( $n= 29$  each): GCP ( $71.22 \pm 3.66$  years old) (received only PMED), and GCS ( $70.77 \pm 4.83$  years old) (no intervention). All laboratory and statistical procedures were conducted in a blinded manner. After the completion of raw data collection, field researchers were no longer exposed to the data to ensure unawareness of individual or collective results until statistical processing. An independent statistician processed the data to prevent bias and enhance result confidence.

### **2.4 BDNF and p-TAU quantification**

Blood collection was conducted by a nurse, extracting 8 mL from the brachial vein into vacuum tubes. Following collection, the blood was promptly stored at  $-20^{\circ}\text{C}$  for up to 72 hours before processing. All blood samples were obtained between 7 and 8 am without prior exercise on the same day.

Assays were performed using peripheral blood samples according to the standard protocol for Peripheral Blood Mononuclear Cells (PBML) at Fiocruz-RO (Bioassay for Leishmania and Malaria Platform FIOCRUZ-RO). BDNF and TAU assays utilized a commercially available sandwich ELISA kit (Chemicon International, USA), following the manufacturer's instructions. Samples were diluted 1:2 and plated on microtiter plates (96 flat-bottom wells) with a standard curve ranging from 7.8 to 500 pg of BDNF (sensitized), incubated for 24 hours. Plates were washed four times with wash buffer and incubated with rabbit anti-BDNF antibody (1:1000 dilution) for 3 hours at room temperature.

Following washing, a second incubation with peroxidase-conjugated anti-rabbit antibody (1:1000 dilution) was carried out at room temperature for 1 hour. BDNF levels were determined by measuring absorbance at 450 nm after the addition of streptavidin, substrate, and prepared solutions. The standard curve demonstrated a direct relationship between optical density (OD) and BDNF concentration, where higher OD values corresponded to higher BDNF concentrations in the analyzed samples. P-TAU levels were analyzed using the same method as for BDNF.

### **2.5 Physical exercise protocols (PMED)**

### **2.5.1 High-Intensity interval training**

The entire program consists of 30-minute classes, structured as follows: 5 minutes for warm-up, 20 minutes for core exercises, and 5 minutes for technical cool-down. Participants begin in a seated position and transition between various standing positions throughout the session.

The core component of the PMED program includes exercises targeting lower body, abdominal, upper body strength and resistance, balance, and overall motor coordination. Intensity levels vary in cycles. Participants have the autonomy to pause and rest if they feel fatigued or uncomfortable, as the exercise regimen is self-regulated. The program is designed to induce fatigue, with resting indicating maximum intensity reached, ensuring a progressive and high-intensity exercise experience.

Training intensity reaches up to 80%, determined using a hypothetical repetition-maximum protocol from the American College of Sports Medicine (ACSM) (Communications et al., 2002), followed by straightforward percentage calculations. During stress tests, exercises are performed at a standardized intensity until concentric failure is achieved. New loads are adjusted based on these outcomes every two weeks. The formula used is: [20 reps at standard load \* (percent work) = reps].

### **2.5.2 Moderated-Intensity Interval Training**

Performed at 50% and 60% of maximum heart rate measured with a finger oximeter (GTECH, model LED, Brazil) in a 30 x 15 m room next to an urban network basic health unit, a walking program of 40 minutes was performed to promote the cool down, a 5-minute relaxation session was performed using very low-intensity stretching techniques similar to yoga. All maneuvers take place in a delineated, flat area of the parking lot. All possible obstacles have been removed to avoid accidents.

## **2.6 Statistical Procedures**

Before participant recruitment, sample size and power calculations were conducted with a focus on changes in heart rate variability. The data did not exhibit a common standard deviation ( $p > 0.05$ ). A sample size of 14 subjects per group was determined necessary to achieve at least 80% power to detect mean differences between groups. The actual achieved power was 91.3%. The power calculation method was previously described (Wedell-Neergaard et al., 2019).

Statistical analysis proceeded through three phases. First, the Kolmogorov-Smirnov test (Dallal-Wilkinson-Lille p-value) assessed data normality, confirming a parametric distribution ( $p > 0.05$ ). Data were then depicted descriptively using mean and standard deviation plots. Second, following confirmation of normality, the Kruskal-Wallis test with Dunn's post hoc test (5% significance level) was employed to identify differences within and between groups in the post-intervention phase. The Spearman correlation test was used. All statistical procedures were conducted using GraphPad Prism 5.0.

### **3. RESULTS AND DISCUSSION**


#### **3.1 The group displayed a homogenous distribution of age, marital status, and formal instruction.**

Population demographic profiles were defined using the following variables: (1) baseline mental state (MMSE), (2) gender, (3) age, (4) income, (5) social income, (6) marital status. Overall, distributions were similar between the groups for almost the entire population profile. As there were no significant differences among most of these variables, statistical checks for normality of distribution were performed only between age and baseline mental status, the only variables showing significant differences between groups. Patients were predominantly in the 65 to 70 age group (50.0%), followed by the 71 to 75 age group (36.6%) and 76 years or older (13.3%). Regarding marital status, more than half of the participants were married (67.83%), followed by divorced (16.6%) or widowed (15.0%).

#### **3.2 The intensity control displays adequate exercise control**

This study investigates the impact of high-intensity body weight interval training (PMED) on brain-derived neurotrophic factor (BDNF) and phosphorylated-TAU protein (p-TAU) in AD and the possible participation of the lactate in crosstalk phenomenon in response to the intensity of the physical exercise. GCP and MICT were found to significantly increase BDNF levels and decrease p-TAU, suggesting a beneficial effect on neurochemistry related to AD pathology. Additionally, GCP overperformed the MICT. To both, the correlation test found strong correlation of the lactate. These findings highlight the potential of high-intensity exercise as a short-term treatment for neurobiological impairments in AD.


The baseline heart rate and blood lactate showed no difference between all groups ( $p > 0.05$ ). However, both reached the planned cardiac frequency. The cardiac frequency measured during the exercise showed differences for all data points in MICT and GCP to Before ( $p > 0.0001$ ). The comparison between groups displays differences between MICT and GCP ( $p > 0.01$ ). Similar results were found to blood lactate. Both, MICT and GCP exhibit differences to Before ( $p > 0.0001$ ) and between then ( $p > 0.01$ ). These data are summarized in figure 1A and 1B.



A and B a=  $p < 0.0001$  vs Before, b=  $p < 0.01$

**Figure 1: Exercise Control.** GCP (n=29, experimental group with AD receiving HIIT), MICT (n=29, with AD receiving walking intervention), and GCS (n=29, negative sitting control without intervention). Figure 1A= Cardiac Frequency; and 1B= Blood Lactate. The Two Way ANOVA with Tukey's Post Hoc test both set up at 5% of significance was used to identify the within-group and between-group possible difference.

Regarding BDNF variants, GCS showed no difference post-intervention ( $p > 0.05$ ). GCP increased post-intervention ( $p < 0.0001$ ) (Fig. 2A). Examining p-TAU (Fig. 2B), GCS values displayed no difference post-intervention ( $p > 0.05$ ). GCP exhibited a decrease in p-TAU ( $p < 0.0001$ ) (Fig. 1B). Although not necessary we performed the size effect showing large effect in regards both molecules ( $d = 0.92$  and  $0.95$ , respectively).




a= 0.0001 vs baseline and b= 0.01 vs MICT-Post

**Figure 2: BDNF and p-TAU quantification.** GCP (n= 29 AD receiving PMED), and GCS (AD n= 29, without intervention). ELISA was used to quantify BDNF and p-TAU proteins before and after the intervention. The Kruskal-Wallis test was followed by DUNN's post hoc test with a significance of 5%. **Note:** (ns= non-significant; \* =  $p < 0.0001$  for intragroup Pre- vs Post-test of GCP and the Post-test of GCP vs all other data points).

Previous studies have shown that exercise, including high-intensity interval training, modulates BDNF levels and enhances neuroplasticity [19]. Our results align with these findings, indicating exercise-induced BDNF release may be related to the exercise intensity (Lima et al., 2017; Silva et al., 2022) which can lead to an avenue of new approach to investigate the mechanisms under these observations. These facts, allow us to affirm that the exercise-based interventions exhibit significant positive impacts on neurochemical impairment in AD. The effect sizes observed in BDNF, and TAU protein levels underscore the relevance of these interventions. Overall, the exercise demonstrates the ability to counteract the common pathological chemical changes in the brain caused by AD contributing to the field's understanding.

### **3.4 The correlation test display strong correlation between blood lactate with BDNF and pTAU in response the high-intensity exercise**

The analysis of the correlation between Lactate and BDNF revealed several significant relationships across different treatment groups. In the GCS group, the correlation was moderate and not statistically significant ( $p=0.8727$ ). In the MICT group, the correlation was weak and non-significant ( $p=0.9090$ ). However, in the GCP group, a strong positive correlation was observed ( $p=0.0169$ ). The correlation between Lactate and p-TAU also showed varying results. In the GCS group, there was a moderate positive correlation that was not statistically significant ( $p=0.5142$ ). The MICT group displayed a weak and non-significant correlation ( $p=0.1845$ ). In the GCP group, there was a strong positive correlation, it did reach statistical significance ( $p=0.0323$ ). These data are summarized in the figure 3.



**Figure 3: Correlation Matrix:** GCP (n=29, experimental group with AD receiving HIIT), MICT (n=29, with AD receiving walking intervention), and GCS (n=29, negative sitting control without intervention). The Pearson Correlation set up at 5% of significance was used to determinate the possible correlation between Lactate and BDNF and Lactate and p-TAU.

To elucidate the mechanisms underlying our findings, we hypothesize that HIIT-induced changes in BDNF and p-TAU levels may be mediated through the release of lactate. Specifically, molecules such lactate are known to cross the blood-brain barrier and exert neuroprotective effects [20–22]. For instance, lactate signaling in the brain increases BDNF production, impacting brain function [23], cognition [24], long-term memory [25], and neuroplasticity [4], many of several impairments related to AD which could be an advantage to promote the physical exercise programs to fight against the AD installation and progress. The hypothesis, not investigated here is that the lactate protects against neuronal injury by activating Akt and ERK1/2 signaling pathways, contributing to the neuroprotective effects of physical exercise [12], and in the regulating JAK2/STAT3 and PACAP signaling pathways, countering NMDA-induced neurotoxicity [26,27], generating a protective molecule linked to intense exercise. In conjunct, we suppose that this molecule exported to the muscle under intense physical exercise could compose the main mechanism under the phenomenon observed.

While our study underscores the potential of HIIT in improving neurochemistry in AD, limitations warrant consideration. These include the relatively small sample size, challenges in medication control, and the absence of beta-amyloid investigation, and another myokines as interleukin-6 and irisin, two between many molecules that can be closed involved in this prosses. Furthermore, the potential influence of anti-inflammatory medications on BDNF levels necessitates careful consideration in future studies.

#### **4. CONCLUSION**

Despite these limitations, our findings support the notion that PMED-induced changes in BDNF and p-TAU levels offer promising avenues for combatting neurobiological impairments in AD. Future research should aim to elucidate these mechanisms further and explore personalized exercise protocols that optimize neuroprotective outcomes.

#### **DISCLAIMER OF ARTIFICIAL INTELLIGENCE**

The authors hereby declare that generative AI technology, CHAT GPT 4.0, was used for editing the English language of the manuscript. The request was solely to correct spelling, punctuation, and grammar, without altering, adding, omitting information or data, or changing the meaning of the text.

#### **CONSENT**

All authors declare that 'written informed consent was obtained from the volunteers for publication of this report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

#### **ETHICAL APPROVAL**

The accomplished all ethical parameters provided by Platform Brazil according to the National Health Council decision CNS 466/12. Registration number 2066823 was issued thereafter. In accordance with the ethical principles like the law, all volunteers and their legal representatives signed a free informed consent form before participating in this study as volunteers after eliminating their doubts about this study.

#### **REFERENCES**

1. Raskin, J.; Cummings, J.; Hardy, J.; Schuh, K.; Dean, R. Neurobiology of Alzheimer's Disease: Integrated Molecular, Physiological, Anatomical, Biomarker, and Cognitive Dimensions. *Curr Alzheimer Res*2015, doi:10.2174/1567205012666150701103107.
2. Pal, M.; Febbraio, M. a; Whitham, M. From Cytokine to Myokine: The Emerging Role of Interleukin-6 in Metabolic Regulation. *Immunol Cell Biol*2014, 92, 331–339, doi:10.1038/icb.2014.16.
3. Jin, Y.; Sumsuzzman, D.M.; Choi, J.; Kang, H.; Lee, S.R.; Hong, Y. Molecular and Functional Interaction of the Myokine Irisin with Physical Exercise and Alzheimer's Disease. *Molecules*2018, 23, 1–13, doi:10.3390/molecules23123229.
4. Müller, P.; Duderstadt, Y.; Lessmann, V.; Müller, N.G. Lactate and BDNF: Key Mediators of Exercise Induced Neuroplasticity? *J Clin Med*2020, doi:10.3390/jcm9041136.

5. Müller, P.; Duderstadt, Y.; Lessmann, V.; Müller, N.G. Clinical Medicine Lactate and BDNF: Key Mediators of Exercise Induced Neuroplasticity?, doi:10.3390/jcm9041136.
6. Congdon, E.E.; Sigurdsson, E.M. Tau-Targeting Therapies for Alzheimer Disease. *Nat Rev Neurol* 2018.
7. Pradeepkiran, J.; Reddy, P. Structure Based Design and Molecular Docking Studies for Phosphorylated Tau Inhibitors in Alzheimer's Disease. *Cells*2019, doi:10.3390/cells8030260.
8. Du, Z.; Li, Y.; Li, J.; Zhou, C.; Li, F.; Yang, X. Physical Activity Can Improve Cognition in Patients with Alzheimer's Disease: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Clin Interv Aging* 2018.
9. Intlekofer, K.A.; Cotman, C.W. Exercise Counteracts Declining Hippocampal Function in Aging and Alzheimer's Disease. *Neurobiol Dis*2013, 57, 47–55, doi:10.1016/j.nbd.2012.06.011.
10. Bernardo, T.; Marques-Aleixo, I.; Beleza, J.; Oliveira, P.; Ascensão, A.; Magalhães, J. PHYSICAL EXERCISE AND BRAIN MITOCHONDRIAL FITNESS : THE POSSIBLE ROLE AGAINST ALZHEIMER ' S DISEASE. *Brain Pathology*2016, 26, 648–663.
11. Belaya, I.; Ivanova, M.; Sorvari, A.; Ilicic, M.; Loppi, S.; Koivisto, H.; Varricchio, A.; Tikkanen, H.; Walker, F.R.; Atalay, M.; et al. Astrocyte Remodeling in the Beneficial Effects of Long-Term Voluntary Exercise in Alzheimer's Disease. *J Neuroinflammation*2020, 17, 1–19, doi:10.1186/s12974-020-01935-w.
12. Li, D.J.; Li, Y.H.; Yuan, H. Bin; Qu, L.F.; Wang, P. The Novel Exercise-Induced Hormone Irisin Protects against Neuronal Injury via Activation of the Akt and ERK1/2 Signaling Pathways and Contributes to the Neuroprotection of Physical Exercise in Cerebral Ischemia. *Metabolism*2017, doi:10.1016/j.metabol.2016.12.003.
13. Souza, T.R.; Campos, P.F.; Almeida, M.; Faria, V.M.; Silveira Chaves, B.; Faria, W.M.; Neves, C.M.; Valentim-Silva, J.R. Exercício Progressivo de Curtíssima Duração Possui Potente Efeito Sobre a Memória de Trabalho, Controle Inibitório e Motricidade Fina de Adultos Jovens Sedentários. / Progressive Exercise of Very Short Duration Has a Potent Effect on Working Memory, Inhib. Motricidade2019, 15, 154–163.
14. Da Silva, V.F.; Rabaioli, E.F.; Azevedo, C.M. de; Maffini, P.H.; Freire, I. de A.; Silio, L.F.; Militão, A.G.; Calomeni, M.R.; Romanholo, R.A.; Valentim-Silva, J.R. Estimulação Cerebral Não Invasiva e Efeito Sinérgico Do Exercício Físico Sobre a Fragilidade e Equilíbrio de Idosos. *Revista Eletrônica Acervo Saúde*2020, 12, e2602, doi:10.25248/reas.e2602.2020.
15. Portela, P.F. de M.; Neto, V.G.C.; Monteiro, E.R.; Santos da Silva, R.; da Silva, V.F.; Nogueira, C.J.; Schutz, S.; Scudese, E.; Salvino, A.K.S.; Valentim-Silva, J.R. HIIT Is Most Effective than Mict on Glycemic Control of Older People with Glucose Metabolism Impairments: A Systematic Review and Metanalysis. *Prim Care Diabetes* 2023, 17.
16. Silva, V.F. da; Silva, D.A.S.; Martins, P.C.; Calomeni, M.R.; Freire, I. de A.; Militão, A.G.; Borges, C.J.; Junior, G. de B.V.; Moraes, M.A. de; Silva, A.J.R.M.; et al. Effect of Physical Exercise and Noninvasive Brain Stimulation on Cognition and Dementia of Elderly People with Frailty : A Randomized Study. *International Journal Imaging System and Technology*2022, 1, 1–12, doi:10.1002/ima.22765.
17. Dubois, B.; Feldman, H.H.; Jacova, C.; DeKosky, S.T.; Barberger-Gateau, P.; Cummings, J.; Delacourte, A.; Galasko, D.; Gauthier, S.; Jicha, G.; et al. Research Criteria for the Diagnosis of Alzheimer's Disease: Revising the NINCDS-ADRDA Criteria. *Lancet Neurology*2007, 6, 734–746, doi:10.1016/S1474-4422(07)70178-3.
18. Fabrício-Wehbe, S.C.C.; Schiaveto, F.V.; Vendrusculo, T.R.P.; Haas, V.J.; Dantas, R.A.S.; Rodrigues, R.A.P. Cross-Cultural Adaptation and Validity of the "Edmonton

- Frail Scale - EFS" in a Brazilian Elderly Sample. *Rev Lat Am Enfermagem*2009, doi:10.1590/S0104-11692009000600018.
19. Murawska-Cialowicz, E.; Wojna, J.; Zuwała-Jagiello, J. Crossfit Training Changes Brain-Derived Neurotrophic Factor and Irisin Levels at Rest, after Wingate and Progressive Tests, and Improves Aerobic Capacity and Body Composition of Young Physically Active Men and Women. *Journal of Physiology and Pharmacology*2015.
  20. Wedell-Neergaard, A.-S.; Lehrskov, L.L.; Christensen, R.H.; Pedersen, B.K.; Ellingsgard, H.; Krogh-Madsen, R. Exercise-Induced Changes in Visceral Adipose Tissue Mass Are Regulated by IL-6 Signaling: A Randomized Controlled Trial. *Cell Metab*2019, 29, 844–855.
  21. Severinsen, M.C.K.; Pedersen, B.K. Muscle–Organ Crosstalk: The Emerging Roles of Myokines. *Endocr Rev*2020, 41, 594–609, doi:10.1210/ENDREV/BNAA016.
  22. Brooks, G.A. Lactate Shuttles in Nature. *Biochem Soc Trans*2002, 30, 258–264, doi:10.1042/.
  23. Numakawa, T.; Suzuki, S.; Kumamaru, E.; Adachi, N.; Richards, M.; Kunugi, H. BDNF Function and Intracellular Signaling in Neurons. *HistolHistopathol* 2010.
  24. Vaynman, S.; Ying, Z.; Gomez-Pinilla, F. Hippocampal BDNF Mediates the Efficacy of Exercise on Synaptic Plasticity and Cognition. *European Journal of Neuroscience*2004, doi:10.1111/j.1460-9568.2004.03720.x.
  25. Bekinschtein, P.; Cammarota, M.; Katche, C.; Slipczuk, L.; Rossato, J.I.; Goldin, A.; Izquierdo, I.; Medina, J.H. BDNF Is Essential to Promote Persistence of Long-Term Memory Storage. *Proc Natl Acad Sci U S A*2008, doi:10.1073/pnas.0711863105.
  26. Zhang, H.J.; Zhang, X.F.; Ma, Z.M.; Pan, L.L.; Chen, Z.; Han, H.W.; Han, C.K.; Zhuang, X.J.; Lu, Y.; Li, X.J.; et al. Irisin Is Inversely Associated with Intrahepatic Triglyceride Contents in Obese Adults. *J Hepatol*2013, doi:10.1016/j.jhep.2013.04.030.
  27. Islam, H.; Neudorf, H.; Mui, A.L.; Little, J.P. Interpreting 'Anti-Inflammatory' Cytokine Responses to Exercise: Focus on Interleukin-10. *Journal of Physiology*2021, 599, 5163–5177, doi:10.1113/JP281356.