

# **Evaluation of Human Energy Consumption in paddy cultivation practices**

## **ABSTRACT**

As the majority of the framers are small and marginal in India accounts for almost 85%, the farming practices are usually doing manually. These manual practices demand much energy consumption from the farmers which ultimately results in the reduced efficiency, productivity of work. The objective of the study is to evaluate the Energy Expenditure Rate (EER), Cardiac Cost of Work (CCW), Cardiac Cost of Recovery (CCR), Total Cardiac Cost of Work (TCCW), and Physiological Cost of Work (PCW) in paddy cultivation practices. Sample size of 10 per activity was taken for the present study in the East and West Godavari districts of Andhra Pradesh. For all the components (EER, CCW, CCR, TCCW, and PCW), heart rate was taken at three different intervals viz. resting phase, working phase, and recovery phase. Each activity in the paddy cultivation was standardized to five minutes. Results found that EER was found to be very high (180.71%), (183.11%) in transplanting activity and threshing activity among female-dominated and male-dominated activities respectively. Similarly, CCW (245bpm, 240bpm); CCR (57.85bpm, 56.62bpm); TCCW (302.85bpm, 296.62bpm); PCW (60.57bpm, 59.32bpm) values were found high in the threshing activity which ranked first among male dominated activities and transplanting activity among female dominated activities. It can be concluded that farmers can opt the tools that are less energy-demanding. Such type of tools is drum seeder, improved sickle, Dapog method of paddy transplanting, etc. Extension agents can suggest the farmers about improved tools that are at low operational cost and can give maximum efficiency and productivity.

Keywords: Energy consumption, productivity, Efficiency.

### **1. Introduction:**

During the Green Revolution, the agricultural sector in India reached self-sufficiency by using better inputs such as high-yielding varieties (HYVs) and inorganic fertilizers. Since

then, the agriculture sector had amazing expansion and now it has reached a substantial worldwide footprint in terms of the production of important staple foods like rice, wheat, milk, sugarcane, fruits, and vegetables. Indian agriculture suffers from the paradox of 'Grain Mountains and Hungry Millions' (Sharma, 2011). Currently, rice is grown on 43.79 million hectares (Mha) of cultivated land in India, with a production of 116.4 million tons (Mt) (GOI, 2020). It is essential to ensure that farmers' incomes are expanded to be commensurate with their efforts and investments. Almost all the cultivation practices in the paddy production system were practiced manually by farmers because the majority of farmers in India are small and marginal, so not in a position to mechanize land for agricultural practices. Cultivation of rice involves many activities that require great physical effort including ploughing the field, uprooting the seedlings from the nursery, planting the saplings in the main field, fertilizer application, harvesting, threshing, and winnowing.

According to Varghese's (1994) classification, Weeding is a strenuous activity as it is heavy activity, physiological cost, and energy expenditure are very high, and Moderate to very severe pain was observed in the lower back, upper back, neck, shoulder, knee, and wrists if done manually. Nawi *et.al.*, (2012) done a study to evaluate the human energy consumption of various cultivation practices involved in lowland rice cultivation in Malaysia. Based on recorded average heart rates, fertilizer application was found to be the most strenuous operation, with an average heart rate of 138 beats  $\text{min}^{-1}$ . There were no significant differences in the average heart rates of the farmers among the individual tasks within the very first ploughing, second ploughing, and harvesting operations, with the average heart rates for these three tasks being 116, 106, and 106 beats  $\text{min}^{-1}$ , respectively. The energy expenditures were 3.90, 3.43, and 3.35 kcal  $\text{min}^{-1}$ . filling the seed into the blower tank and broadcasting the seed were regarded as the most critical tasks for the seed broadcasting operation, with average heart rates of 124 and 136 beats  $\text{min}^{-1}$ , respectively. The highest energy expenditure of 418.38 kcal  $\text{ha}^{-1}$  was reported for seed broadcasting, and the lowest energy expenditure of 127.96 kcal  $\text{ha}^{-1}$  was reported for second ploughing. The total seasonal human energy expenditure for rice cultivation was estimated to be 5810.71 kcal  $\text{ha}^{-1}$ , 55.7 percent of which has been spent on pesticidal spray. Though the sample size in the study was relatively small, the results showed that human energy expenditure per unit area (kcal  $\text{ha}^{-1}$ ) was positively linked to the average heart rate of the farmers and negatively linked to the field capacity. Anita *et.al.*, (2019) revealed that among the four traditional agricultural activities, a maximum average heart rate of 107.8 beats  $\text{min}^{-1}$  was observed while farmers performing

weeding with traditionally available tools, and among the activities with improved implements, use of long handle weeder in vegetable cultivation recorded the maximum average heart rate of 101.8 beats min<sup>-1</sup>, followed by use of sapling transplanter for transplanting of tomato recorded the average heart rate of 97.6 beats min<sup>-1</sup>.

## 2. Sampling plan:

The present study was done in the two highest paddy producing districts in Andhra Pradesh namely East Godavari and West Godavari. From each district 5 samples were taken randomly which constitute 10 samples for each activity in paddy cultivation.

## 3. Materials and methods:

For EER, Heartbeat was taken as the parameter and it was taken at three different intervals which yields the percentage increase in the Energy Expenditure from the resting phase to the working phase in the particular activity. Each activity was standardized for five minutes. Sample size is 10 for each activity and includes both males and females. Present study was carried out in west and east Godavari districts of Andhra Pradesh. Results were presented separately for both the males and females according to the activities they performed.

### 2.1 Classification of workload; (Nag *et al.*, 1980)

Physical workload	Physiological parameters	
	Energy Expenditure (KJ/min)	Heartbeats (beats/min)
Very light	Up to 5.0	Up to 90
Light	5.0-7.5	91-105
Moderate	7.6-10.0	106-120
Heavy	10.0-12.5	121-135
Very Heavy	12.6-15.0	136-150
Extremely Heavy	<15.0	Above 151

### 2.2 Energy Expenditure (EE):

EE calculated using the following formula (Varghese *et al.*, 1994)

Energy expenditure (kJ/min) =  $0.159 \times \text{Heart rate at resting phase (beats/min)} - 8.72$ ;  
 $0.159 \times \text{Heart rate at working phase (beats/min)} - 8.72$ ;  $0.159 \times \text{Heart rate at recovery phase (beats/min)} - 8.72$ . (Varghese *et al.*, 1994)

### **2.3 Cardiac Cost of Work (CCW):**

$$\text{CCW} = \Delta \text{HR} \cdot t_A$$

Where, CCW = Cardiac Cost of Work,  $\Delta \text{HR}$  = Mean Working Heart Rate – Mean Resting Heart Rate,  $t_A$  = Duration of Activity

### **2.4 Cardiac Cost of Recovery (CCR):**

$$\text{CCR} = (\text{AHR recovery} - \text{AHR rest}) \cdot t_R$$

Where, CCR = Cardiac Cost of Work, AHR recovery = Average Recovery Heart Rate,  
AHR rest = Average Resting Heart Rate,  $t_R$  = Duration of Recovery

### **2.5 Total Cardiac Cost of Work (TCCW):**

$$\text{TCCW} = \text{CCW} + \text{CCR}$$

Where, CCW = Total Cardiac Cost of Work, CCR = Cardiac Cost of Recovery

### **2.6 Physiological Cost of Work (PCW):**

$$\text{PCW} = \text{TCCW}/t_A$$

Where, TCCW = Total Cardiac Cost of Work,  $t_A$  = Duration of activity

## **4. Results and Discussion:**

### **4.1 Energy Expenditure Rate (EER):**

For EER, Heartbeat was taken as the parameter and it was taken at three different times in a time-bound activity viz, Working phase, Resting phase, and finally Recovery phase. EER is the percentage change in the energy in Kilo Joules per minute from the resting phase to the working phase. Each activity was standardized for five minutes with a sample size of 10 was taken for each activity.

**Table..1 Energy Expenditure Rate values in the paddy cultivation practices done by male farmers (N=10)**

Activities	Resting phase (KJ/min)	Working phase (KJ/min)	Energy Expenditure (EE) (KJ/min) (Mean and SD)	Percent change in EER per minute	Interpretation w.r.t. physical load
Land preparation	4.04	9.87	20.27( $\pm$ 0.87)	143.84%	Extremely Heavy
Irrigation	4.33	10.52	20.00( $\pm$ 2.12)	142.71%	Extremely Heavy
Manure and fertilizer application	4.28	11.59	20.67( $\pm$ 1.87)	170.6%	Extremely Heavy
Pesticidal spray	4.17	11.21	20.89( $\pm$ 1.10)	168.67%	Extremely Heavy
Threshing	4.29	12.14	22.43( $\pm$ 0.92)	183.11%	Extremely Heavy

**Table. 2 Energy Expenditure Rate values in the paddy cultivation practices done by female farmers (N=10)**

Activities	Resting phase (KJ/min)	Working phase (KJ/min)	Energy Expenditure (EE) (KJ/min) (Mean and SD)	Percent change in EER per minute	Interpretation w.r.t. physical load
Uprooting	4.27	10.02	19.93( $\pm$ 1.30)	134.78%	Extremely Heavy
Transplanting	4.33	12.16	23.00( $\pm$ 0.79)	180.71%	Extremely Heavy
Weeding	4.13	11.17	21.29( $\pm$ 0.88)	170.28%	Extremely Heavy
Harvesting	4.14	11.53	22.61( $\pm$ 1.42)	178.26%	Extremely Heavy
Winnowing	4.19	10.82	21.29( $\pm$ 1.30)	158.37%	Extremely Heavy

From the above table 1, it was concluded that EER in the threshing activity was found to be very high (183.11 percent) which indicates that it was the most energy-demanding

activity in paddy cultivation practices. So threshing activity stood first. Followed by manure and fertilizer application activity (170.60 percent), Pesticidal spray (168.67 percent), Land preparation (143.84 percent), and Irrigation (142.71 percent) which were ranked second, third, fourth, and fifth respectively.

From the above table.2, it was concluded that EER in the transplanting activity was found to be very high (180.71 percent) which indicates that it was the most energy-demanding activity in paddy cultivation practices. So, transplanting activity stood first. Followed by harvesting activity (178.26 percent), weeding (170.28 percent), winnowing (158.37 percent), and uprooting (134.78 percent) which were ranked second, third, fourth, and fifth respectively.

#### 4.2 Cardiac Cost of Work (CCW)

**Table 3 Cardiac Cost of Work values in the paddy cultivation practices done by male farmers (N=10)**

Activities done by male farmers	Cardiac Cost of Work (CCW) in beats/minute (Mean and SD)
Land preparation	198 ( $\pm 15$ )
Irrigation	190 ( $\pm 11$ )
Manure and fertilizer application	215 ( $\pm 19$ )
Pesticidal spray	216 ( $\pm 16$ )
<b>Threshing</b>	<b>245 (<math>\pm 21</math>)</b>

It was observed from table 4 that CCW values in the threshing activity among male farmers were found to be high (245 bpm) which ranked first followed by pesticidal spray (216bpm), manure and fertilizer application (215bpm), land preparation (198bpm), and irrigation (190 bpm) activities ranked second, third, fourth, and fifth respectively.

**Table 4 Cardiac Cost of Work values in the paddy cultivation practices done by female farmers (N=10)**

Activities done by female farmers	Cardiac Cost of Work (CCW) in beats/minute (Mean and SD)
Uprooting	181 ( $\pm 14$ )
<b>Transplanting</b>	<b>240 (<math>\pm 22</math>)</b>
Weeding	208 ( $\pm 16$ )

Harvesting	226(±18)
Winnowing	199(±11)

It was observed from above table 5 that CCW values in the transplanting activity among female farmers found to be high (240 bpm) which ranked first followed by harvesting (226 bpm), weeding (208 bpm), winnowing (199 bpm), and uprooting (181 bpm) activities ranked second, third, fourth, and fifth respectively.

### 4.3 Cardiac Cost of Recovery (CCR)

**Table 5. Cardiac Cost of Recovery values in the paddy cultivation practices done by male farmers (N=10)**

Activities done by male farmers	Cardiac Cost of Recovery values in beats/min (Mean and SD)
Land preparation	36.45(±2.3)
Irrigation	28.8(±1.7)
Manure and fertilizer application	33.48(±2.1)
Pesticidal spray	37.8(±3.4)
<b>Threshing</b>	<b>57.85(±4.8)</b>

From the table 6, it was concluded that CCR values in the activities done by male farmers have been found that threshing (57.85 bpm) ranked first followed by pesticial spray (37.80 bpm), land preparation (36.45 bpm), Manure and fertilizer application (33.48 bpm), and irrigation activities (28.8 bpm) which were ranked second, third, fourth, and fifth respectively

**Table 6 Cardiac Cost of Recovery values in the paddy cultivation practices done by female farmers (N=10)**

Activities done by female farmers	Cardiac Cost of Recovery values in beats/min (Mean and SD)
Uprooting	30.1(±1.6)
<b>Transplanting</b>	<b>56.62(±4.7)</b>
Weeding	40.6(±2.9)

Harvesting	53.64( $\pm$ 3.7)
Winnowing	39.3( $\pm$ 1.8)

From the table 7 it was concluded that CCR values in the activities done by female farmers have been found that transplanting activity (56.62 bpm) ranked first followed by harvesting (53.64 bpm), weeding (40.60 bpm), Winnowing (39.30 bpm), and uprooting activities (30.1 bpm) which were ranked second, third, fourth, and fifth respectively.

#### 4.4 Total Cardiac Cost of Work (TCCW)

**Table 7 Total Cardiac Cost of Work values in the paddy cultivation practices done by male farmers (N=10)**

Activities done by male farmers	Total Cardiac Cost of Work values in beats/min (Mean and SD)
Land preparation	234.45 ( $\pm$ 31.3)
Irrigation	218.8 ( $\pm$ 19.3)
Manure and fertilizer application	248.48 ( $\pm$ 22.4)
Pesticidal spray	253.8 ( $\pm$ 23.6)
<b>Threshing</b>	<b>302.85 (<math>\pm</math>18.1)</b>

From the table 8 it was concluded that TCCW values in activities done by male farmers have been found that threshing (302.85 bpm) ranked first followed by pesticidal spray (253.8 bpm), manure and fertilizer application (248.48 bpm), land preparation (234.45 bpm), and irrigation activities (218.8 bpm) which were ranked second, third, fourth, and fifth respectively.

**Table 8 Total Cardiac Cost of Work values in the paddy cultivation practices done by female farmers (N=10)**

Activities done by female farmers	Total Cardiac Cost of Work values in beats/min (Mean and SD)
Uprooting	211.1( $\pm$ 13.1)
<b>Transplanting</b>	<b>296.62(<math>\pm</math>19.6)</b>

Weeding	248.60(±22.1)
Harvesting	279.64(±17.9)
Winnowing	238.30(±19.4)

From the above table 9 it was concluded that TCCW values in activities done by female farmers have been found that transplanting activity (296.62 bpm) ranked first followed by harvesting (279.64 bpm), weeding (248.60 bpm), Winnowing (238.30 bpm), and uprooting activities (211.1 bpm) which were ranked second, third, fourth, and fifth respectively.

#### 4.5 Physiological Cost of Work (PCW)

**Table 9 Physiological Cost of Work values in the paddy cultivation practices done by male farmers (N=10)**

Activities done by male farmers	Physiological Cost of Work values in beats/min (Mean and SD)
Land preparation	46.89(±3.9)
Irrigation	43.76(±4.1)
Manure and fertilizer application	49.69(±3.8)
Pesticidal spray	50.76(±5.2)
<b>Threshing</b>	<b>60.57(±4.9)</b>

From the table 10, it was concluded that PCW values in activities done by male farmers have been found that threshing (60.57 bpm) ranked first followed by pesticidal spray (50.76 bpm), manure and fertilizer application (49.69 bpm), land preparation (46.89 bpm), and irrigation activities (43.76 bpm) which were ranked second, third, fourth, and fifth respectively.

**Table 10 Physiological Cost of Work values in the paddy cultivation practices done by female farmers (N=10)**

Activities done by female farmers	Physiological Cost of Work values in beats/min (Mean and SD)
Uprooting	42.22(±4.1)
<b>Transplanting</b>	<b>59.32(±4.6)</b>
Weeding	49.72(±39)
Harvesting	55.92(±5.3)
Winnowing	47.66(±2.7)

From the table 10, it was concluded that PCW values in activities done by female farmers have been found that transplanting activity (59.32 bpm) ranked first followed by harvesting (55.92 bpm), weeding (49.72 bpm), Winnowing (47.66 bpm), and uprooting activities (42.22 bpm) which were ranked second, third, fourth, and fifth respectively.

#### **5. Discussion:**

The physiological workload of a worker can majorly be reduced using a conoweeder for wetland paddy crops, The physiological cost of work and energy expenditure of respondents was 10.36 beats min<sup>-1</sup> and 7.50 KJ min<sup>-1</sup> respectively against 19.83 and 10.82 if done manually (Awasthi *et al.*, 2018).Mechanization of some tasks could drastically decrease worker physical effort and fatigue and thereby increase production (Nawi *et al.*, 2012).

#### **6. Conclusion and Way forward:**

All the cultivation practices in paddy were found to be very difficult to do. If the farmers continue to do the practices manually, it leads to various health hazards which ultimately results in the reduction of efficiency and productivity of the work. There is a need to conduct the extension programmes, campaigns with the special attention to the drudgery involved cultivation practices. Extension agents should make the farmers aware of improved tools available for the paddy cultivation practices such as improved sickle, weeder, transplanter etc. which helps in reduced energy consumption in doing the cultivation practices.

#### **9. Ethical considerations:**

Data was collected with the informed consent of the framers.

## 10. References:

- Agricultural statistics at a glance, Directorate of Economics and Statistics, Ministry of Agriculture, Government of India; 2022.
- Anitha, R. O., Singh, B. K., & Afifa, J. (2019). An evaluation of drudgery reducing agricultural technologies developed for farm women. *International Journal of Agricultural Science and Research*, 9(2), 35-42.
- Awasthi, N., Singh, A. K., Yadav, C. K., & Kumar, A. (2018). Performance analysis of Conoweeder to combat drudgery involved in weeding of paddy. *The Pharma Innovation Journal*, 7(7), 251-255.
- GOI (2020). Directorate of economics and statistics, ministry of agriculture and farmers' welfare, government of India.
- Joshi, P., Kumar, A., Bisht, K., and Singh, O. P. (2020). Ergonomics assessment of drudgery in wheat production system. *Indian Journal of Extension Education*, 56(4), 155-159.
- Kumari, S., & Gupta, R. (2023). Assessment of musculoskeletal problems of the women involved in peas harvesting. *Journal of Community Mobilization and Sustainable Development*, 18(2), 705-708.
- Nag, P. K., Sebastian, N. C., & Malvankar, M. G. (1980). Occupational workload in Indian agricultural workers. *Ergonomics*, 23: 91-102.
- Nawi, N. M., Yahya, A., Chen, G., Bockari-Gevao, S. M., & Maraseni, T. N. (2012). Human energy expenditure in lowland rice cultivation in Malaysia. *Journal of agricultural safety and health*, 18(1), 45-56.
- Pandey, S., Dubey, S. K., Singh, A., Gautam, U. S., Singh, R., Tripathi, K. M., Saurabh, Awasthi, N., & Kumari, A. R. (2022). Drudgeries and occupational health hazards perceived by the women farmers in Central zone of Uttar Pradesh. *Journal of Community Mobilization and Sustainable Development*, 17(1), 61-66.
- Sadhna, P.; S.K. Dubey; A. Singh; U.S. Gautam; K.M. Tripathi; Saurabh; K.A. Ranjan; A. Singh and A. Nimishi. 2020. Occupational health hazards and drudgeries perceived by the women farmers in Western Zone of Uttar Pradesh. *Journal of Community Mobilization and Sustainable Development*, 15(1): 1-7.

Sanjit K., *et al.* "Ergonomical evaluation of occupational problems of women workers in agricultural tasks." *International Journal of Basic and Applied Physiology* 1.1 (2012): 103-8.

Sharma, D. (2011). India's shameful paradox of plenty. counter currents.org

Shukla, P., Mehta, C. R., Agrawal, K. N., Potdar, R. R., Kumar, M., & Singh, K. (2023). Approach for ergonomic assessment of self-propelled combine harvester seats based on anthropometric body dimensions. *International Journal of Occupational Safety and Ergonomics*, 29(1), 282-293.

Varghese, M. A., Saha, P. N., & Atreya, N. (1994). A rapid appraisal of occupational workload from a modified scale of perceived exertion. *Ergonomics*, 37(3), 485-491.

UNDER PEER REVIEW