

1 Original Research Article

2

3 **MANAGEMENT OF FUSARIUM WILT OF CASTOR (RICINUS COMMUNIS L.)**
4 **CAUSED BY *Fusarium oxysporum f.sp. RICINI* WITH BIORATIONALS AND**
5 **FUNGICIDES UNDER *IN VITRO* CONDITIONS**

6 **Keywords:** Castor, *Fusarium* spp, Biorationals, Fungicides

7 **ABSTRACT**

8 To mitigate the harmful effects of synthetic chemical pesticides, eight biorationals *i.e*
9 ginger, onion, bougainvillea, garlic, chili, turmeric, neem, and aloe vera were tested at
10 different test concentrations of 10%, 15%, and 20% against *Fusarium oxysporum f. sp. ricini*,
11 the pathogen causing wilt in castor. Among all the tested biorationals, chilli found as most
12 effective while aloe vera showed the least pathogen suppression. The study recommended neem
13 and chili as effective biorationals against *F. oxysporum f. sp. ricini*. New fungicide combinations *i.e*
14 Picoxystrobin + Tricyclazole, Fluopyram + Tebuconazole, Boscalid + Pyraclostrobin, and
15 Iprovalicarb + Propineb were tested under *in vitro* condition out of which Picoxystrobin +
16 Tricyclazole and the standard fungicides Carbendazim and Thiophanate-methyl showed cent
17 percent inhibition at all tested concentrations (500, 1000, 1500 and 2000 ppm). Fluopyram +
18 Tebuconazole and Boscalid + Pyraclostrobin also showed high efficacy, while Iprovalicarb +
19 Propineb was the least effective. The study revealed that neem and chili are recommended as
20 effective biorationals against *F. oxysporum f. sp. ricini*. Among fungicides, Picoxystrobin +
21 Tricyclazole proved most effective, suggesting its potential for managing castor wilt.

22 **Keywords:** Castor, *Fusarium* spp, Biorationals, Fungicides

23 **1. INTRODUCTION**

24 Castor (*Ricinus communis*), a versatile and resilient plant, has captivated the interest of
25 mankind for a long time for its numerous applications. This hardy plant exhibits exceptional
26 oil production, yielding 350 and 900 kg of oil per hectare, and demonstrated the adaptability
27 to marginal soils. The oil has numerous industrial applications, as evidenced by the recent
28 surge in global demand, with over 700 documented applications (Chakrabarty *et al.*, 2021).
29 India stands as a prominent global cultivator of castor, holding the foremost position in terms
30 of both cultivation (0.83 million ha) and production (1.57 million tons) [Yamanura and

31 Kumar, 2020]. at all crop growth stages depending on the seasonal conditions [Raouf and
32 Nageshwar Rao, 1996]. Among all diseases *Fusarium* wilt, is one such important disease
33 which is soil and seed-borne.

34 The fungus is 10–20 per cent internally as well as externally seed borne
35 [Chattopadhyay, 2000]. The disease affects all crop growth phases throughout the year,
36 however it spikes during the flowering, spike production, and capsule maturity stage. The
37 disease was initially reported in Morocco [Rieuf, 1953] and in India, at Udaipur and Sirohi of
38 Rajasthan [Nanda and Prasad, 1975]. Description of the symptoms of castor wilt caused by *F.*
39 *oxysporum* f. sp. *ricini* seen as yellowing, sickly appearance of infected plant and marginal
40 leaf necrosis, which later covers the leaves entirely. Drooping of lower leaves leaving limited
41 top leaves followed by irretrievable wilting of the plant leading to rapid death [Nanda and
42 Prasad, 1975]. Transverse and longitudinal sections of the affected roots revealed the
43 presence of fungus in vascular tissues and xylem parenchyma. The tyloses development is
44 observed in xylem vessels of infected roots of castor plants.

45 Recent advancements in integrated disease management have focused on the use of
46 biorational products and novel fungicides to mitigate *Fusarium oxysporum* infections.
47 Biorationals, such as biocontrol agents and plant extracts, offer promising alternatives due to
48 their eco-friendly nature and minimal impact on non-target organisms [Cárdenas-Laverde et
49 al, 2021]. These products harness the natural mechanisms to suppress the fungal growth and
50 enhance plant resistance.

51 The current study aims to evaluate the efficacy of selected biorationals and new
52 fungicide molecules against *F. oxysporum* f. sp. *ricini* under controlled *in vitro* conditions.
53 By assessing their impact on fungal growth inhibition, spore germination, and disease
54 progression, this research seeks to contribute to the development of integrated disease
55 management strategies for sustainable castor bean production. Through this investigation, we
56 aim to provide valuable insights into the potential of biorationals and new fungicides as
57 effective tools for managing *F. oxysporum* f. sp. *ricini* in agricultural settings. This research
58 plays a crucial role in advancing the understanding of alternative disease management
59 practices that balance the efficacy with both environmental and economic sustainability.

60

61

62 2. MATERIALS AND METHODS

63 2.1 Collection and Preparation of inoculum

64 Plants showing characteristic wilt symptoms were collected from the AICRP Castor
65 Research plot, K-block, ZARS, GKVK, UAS, Bengaluru. Tissue isolation technique was
66 followed to isolate the pathogen on Potato Dextrose Agar (PDA) under *in vitro* condition.

67 2.1.1 Preparation of inoculum

68 Semi-cooked sorghum grains (100g in 250 ml conical flask) were autoclaved for
69 approximately 20 minutes at 121°C and 15 psi pressure. Subsequently, the flasks were
70 inoculated with actively growing fungal mycelial culture derived from purified culture on
71 Potato Dextrose Agar (PDA). The inoculated flasks were then placed in an incubator and
72 maintained for 15 days at $27 \pm 2^\circ\text{C}$. Once the sorghum grains were completely colonized by
73 the fungus, they were regularly shaken by hand to achieve a concentration of 1×10^6
74 conidia/ml.

75 2.2 EVALUATION OF BIORATIONALS *IN VITRO*

76 Efficacy of biorationals against the test pathogen were tested by following Poison
77 Food Technique as described by Nene and Thapliyal, 1973. Plant parts of eight selected
78 medicinal crops (100 g) were first washed thoroughly with tap water and then with sterilized
79 distilled water and air dried, weighed plant materials were ground in pestle and mortar. The
80 materials were homogenized for 5 minutes and then filtered through double layered muslin
81 cloth followed by Whatman No. 1 filter paper and filtrates were considered as standard
82 extract (100%). The standard extracts solution was individually incorporated into sterilized
83 medium in 250 ml conical flasks at required quantities of 10%, 15% and 20% concentration
84 after Millipore filtration. The melted PDA was poured into 90 mm sterilized Petri plate and
85 PDA without extracts was maintained as control. All plates were replicated three times and
86 the data was analysed statistically using Complete Randomized Design (CRD). Plates were
87 inoculated with 5 mm mycelium discs of seven days old fungal culture and incubated at
88 $26 \pm 10^\circ\text{C}$ for 5 days. The radial growth of the mycelium was measured after a day of
89 incubation and percent growth inhibition was calculated (Vincent, 1947). The plates were
90 incubated until the growth of the control plate was completely covered by test fungus. Each
91 treatment was replicated thrice. The efficacy of different biorationals was expressed as per

92 cent inhibition of mycelial growth over control and calculated by using the formula [Vincent,
93 1947] as follows

$$94 \quad I = (C-T)/C \times 100$$

95 where ,

96 I = Per cent inhibition in growth of pathogen

97 C = Radial growth of the test fungus in control (mm) and

98 T = Radial growth of the test fungus in treatment (mm)

99 **2.3 EVALUATION OF NEW FUNGICIDE MOLECULES *IN VITRO***

100 Poison food technique was followed to evaluate the efficacy of fungicide molecules
101 under *in vitro* for their effect on fungus development [Nene and Thapliyal, 1973]. PDA
102 medium was prepared and autoclaved, various test fungicides of concentrations 500 ppm,
103 1000 ppm, 1500 ppm and 2000 ppm were prepared adding to the same medium. Initially the
104 test fungicides were dissolved in 100 ml of sterile molten PDA medium and 15 ml of PDA
105 was poured into sterile Petri plates under aseptic conditions and allowed to solidify. Mycelial
106 discs of *Fusarium* (7 mm diameter) were cut from 7-day old culture plate and placed it in the
107 centre of Petri plate containing PDA medium amended with fungicide. The PDA medium
108 (without fungicide) inoculated with test fungus alone served as control (check). The plates
109 were incubated at room temperature. Each treatment was replicated thrice. The diameter of
110 the fungal colonies in the treatments was measured when the growth in the control plate was
111 full. The colony diameter was measured, compared with control and reduction in the growth
112 was taken as a measure of fungi toxicity. Per cent inhibition of the pathogen in different
113 chemical treatments over control was calculated by the following formula.

$$114 \quad I = (C-T)/C \times 100$$

115 where ,

116 I = Per cent inhibition in growth of pathogen

117 C = Radial growth of the test fungus in control (mm) and

118 T = Radial growth of the test fungus in treatment (mm)

119 **3. RESULTS AND DISCUSSION**

120 **3.1 *In vitro* evaluation of biorationals and new fungicide molecules against *Fusarium***
121 ***oxysporum* f. sp. *ricini* wilt pathogen of castor**

122 To reduce the harmful effect of synthetic chemical pesticides, the current study was
123 planned with biorationals which are biological products extracted from the plant source and
124 are used directly in the process of pathogen inhibition. In the current investigation, a total of
125 eight biorationals were tested viz., ginger, onion, bougainvillea, garlic, chilli, turmeric, neem
126 and aloe vera at three different concentrations of 10, 15, and 20 per cent against *F.*
127 *oxysporum* f. sp. *ricini* pathogen causing wilt in castor.

128 The data from the Table 1, Figure 1 A and Plate 1A clearly showed that Neem (T₆)
129 showed a mean maximum inhibition of 46.30 % followed by chilli (T₇) with the mean
130 inhibition per centage of 46.05 however both are on par with one another. Lowest mean
131 inhibition percentage was recorded in T₈ i.e Aloe vera (28.02%). Among all the biorationals
132 tested at three different concentrations i.e 10, 15 & 20 %. Neem @ 10 & 15 % concentration
133 showed maximum inhibition of 54.07 & 53.70 % respectively followed by ginger at 15%
134 concentration (53.33%) and with inhibition percentage of 51.85 and 51.11 % in chilli at 10 &
135 15 % respectively.

136 Efficacy of four new fungicides along with two standard fungicides as checks
137 (Thiophante methyl 70 % WP and Carbendazim 50 % WP) were evaluated against *Fusarium*
138 *oxysporum* f. sp. *ricini* at four different concentrations viz., 500, 1000, 1500 and 2000 ppm
139 by following Poisoned Food Technique as described in Materials and Methods (Section 2.3)

140 The data from the Table 2, Figure 1B and Plate 1 B revealed that among the tested
141 combi products against the test pathogen, picoxytrsobin 6.78%+Tricyclazole 20.33% was
142 found highly effective and completely inhibited the mycelia growth of the test pathogen at all
143 four different concentrations tested and it was superior over their combi products treatments
144 tested in reducing the radial growth of the pathogen whereas, Fluopyram 17,7% +
145 Tebuconazole 17.7% was found to be the next best with per cent inhibitions pof 87.41 (1500
146 ppm); 87.04 (1000 & 2000 ppm) and 86.67 (500 ppm) followed by Boscalid 25.2 % +
147 Pyraclostrobin 12.8 % with the inhibition of 83.70% (2000 ppm) and 80.74 % (1500 ppm).
148 Lowest inhibition per Cent of 31.85 (500 ppm), 14.81 (1000 ppm), 30.74 (1500 ppm) and
149 41.48 (2000 ppm) were recorded with Iprovalicarb 5.5 % + propineb 61.25%.

150 It is evident from the results that the inhibition in the growth of the test pathogen
151 increased with the increase in the concentration of the fungicides. Two standard check
152 fungicides (Thiophante Methyl and carbendazim) also recorded cent per cent inhibition at
153 500, 1000, 1500 and 2000 ppm.

154 **DISSCUSSION**

155 Evaluation of biorationals and new fungicide molecules *in vitro* against *F. oxysporum*
156 f. sp. *ricini* has provided insightful data on their efficacy in controlling the wilt disease in
157 castor. The current investigation aimed to identify the environmentally friendly and effective
158 treatments to synthetic chemicals pesticides, addressing both pathogen suppression and
159 sustainability concerns. Among the eight biorationals tested (ginger, onion, bougainvillea,
160 garlic, chili, turmeric, neem, and aloe vera).

161 Neem extract showed the highest inhibition at 10% and 15% concentrations, followed
162 by Ginger and chilli extracts, neem and chili extracts emerged as promising treatments for
163 further exploration and application in integrated disease management strategies for castor wilt
164 under field conditions also.

165 The current results are in conformity with Vahunia *et al*, 2017; Yelmane *et al*, 2010
166 and Elhelaly and Ammar, 2022 who reported that the extract of turmeric rhizome showed
167 inhibition of 42.22 per cent followed by marigold leaves extract (36.67%) and least by datura
168 (18.89%)

169 Singh *et al*. (2021) evaluated the botanicals in the management of *F. o. f. sp. ciceri* of
170 chickpea and found that neem leaves was most effective at all three concentrations (5, 10 and
171 15%) with per cent of inhibition of 86.61, 89.66 and 94.29 respectively.

172 Among combi fungicides tested at four different concentrations (250, 500, 1000, 1500
173 ppm). Carbendazim 12% + Mancozeb 63% recorded highest growth inhibitions of 66.39,
174 69.02, 69.88, 75.12 percent respectively. The least was recorded in Zineb 68% +
175 Hexaconazole 4% with the inhibition per cent of 37.07, 43.66, 45.23 and 48.16 per cent at
176 500, 1000, 1500 and 2000 ppm, respectively.

177 Vani *et al*. (2019) reported that Mancozeb 50% + Thiophanate-methyl 25% found as
178 best treatment at the concentration of 1500 ppm followed by Propineb (86.6%) at 2000 ppm
179 and least inhibition was observed in Azoxystrobin (55.8%) at 1000 ppm.

180 **4. CONCLUSION**

181 Evaluation of biorationals and new fungicide molecules against *F. oxysporum* f. sp.
182 *ricini* provided significant valuable insights into their potential for managing wilt disease in
183 castor. Among the biorationals, neem extract demonstrated the highest efficacy in inhibiting
184 mycelial growth at 10% and 15% concentrations, followed by chili extract at 20%. These
185 findings highlight the potential use of neem and chili extracts as promising biocontrol agents
186 for sustainable castor wilt disease under field conditions too. Picoxystrobin 6.78% +
187 Tricyclazole 20.33% and the two standard check fungicides, Carbendazim 50% and
188 Thiophanate-methyl 70% achieved cent per cent inhibition across all tested concentrations.
189 Fluopyram 17.7% + Tebuconazole 17.7% served as next best alternative in reducing the
190 mycelia growth of *F. oxysporum* f. sp. *ricini* and maximum inhibition percentage.

191 **5. FUTURE LINE OF WORK**

192 Future research should focus on field trials to validate the efficacy of the most
193 promising biorationals and fungicide combinations which were proved as effective under *in*
194 *vitro* conditions. Additionally, exploring the synergistic effects of combining biorationals
195 with fungicides could enhance overall disease management strategies. Investigations into the
196 modes of action, optimal application methods, and long-term environmental impacts of these
197 treatments will be crucial for developing integrated and sustainable approaches for managing
198 the *Fusarium* wilt in castor.

199

200

201

202 **ETHICAL STATEMENT**

203 All the experimental procedures involving only on plant species which were
204 conducted in accordance with the University of Agricultural Science, Bangalore institutional
205 guidelines. There are no human and animal subject/ trails conducted in this article and
206 informed consent is not applicable.

207

208 **REFERENCES**

- 209 1. Chakrabarty S, Islam AKMA, Yaakob, and Islam AKMM. Castor (*Ricinus*
210 *communis*): An underutilized oil crop in the South East Asia. *Agroecosystems-Very*
211 *Complex Environmental Systems*; IntechOpen: London., UK, 2021; 61.
- 212 2. Yamanura and Kumar RM. An overview of utility, status, retrospective and prospects
213 of castor: A Review. 2020.
- 214 3. Raof MA, Nageshwar Rao TG. Effect of soil solarization on castor wilt. *Indian*
215 *Journal Plant Protection*. 1996; 25: 154-159.
- 216 4. Chattopadhyay C. Seed borne nature of *Fusarium oxysporum* f. sp. *ricini* and
217 relationship of castor wilt incidence with seed yield. *Journal of mycology plant*
218 *pathology*. 2000; 30: 265.
- 219 5. Rieuf P. A note on wilt of castor plant. *Review of Pathology*. 1953; 32: 120-129.
- 220 6. Nanda S, Prasad N. Wilt of castor a new record. *Indian Journal of mycology and plant*
221 *Pathology*. 1974; 4(1): 103-105.
- 222 7. Cárdenas-Laverde D, Barbosa-Cornelio R, Coy-Barrera E. Antifungal activity against
223 *Fusarium oxysporum* of botanical end-products: an integration of chemical
224 composition and antifungal activity datasets to identify antifungal bioactives. *Plants*.
225 2021; 10(12): 2563.
- 226 8. Vincent JM. Distribution of fungal hyphae in the presence of certain inhibitors.
227 *Nature*. 1947; 159: 850.
- 228 9. Nene YL, Thapliyal PN. *Fungicide in plant diseases control* 2nd edn. 1973.
- 229 10. Vahunia B, Singh P, Vaja SJ. *Eco-Friendly Management of Castor (*Ricinus communis**
230 *L.) Wilt Caused by *Fusarium oxysporum* f. sp. *ricini* by Organic Manure/Cakes in*
231 *Vitro*. 2017.
- 232 11. Yelmame MG, Mehta BP, Deshmukh AJ, Patil VA. Evaluation of some organic
233 extracts in *in vitro* to control *Fusarium solani* causing chilli wilt. 2010.
- 234 12. Elhelaly SH, Ammar HA. Bio-efficacy of some Botanicals, Antagonistic fungi, and
235 Fungicides against *Fusarium oxysporum* f. sp. *phaseoli* causing *Fusarium* wilt on
236 common bean plants. *Egyptian Journal of Crop Protection*. 2022; 17(1): 1-14.

- 237 13. Vani MS, Kumar S, Gulya R. *In vitro* evaluation of fungicides and plant extracts
 238 against *Fusarium oxysporum* causing wilt of mungbean. The Pharma Innovation.
 239 2019; 8(8): 297-302.
- 240 14. Jadav AH, Akbari LF, Barad AJ, Bhaliya CM. Efficacy of different fungicides for the
 241 management of castor wilt (*Fusarium oxysporum* f. sp. *ricini*). The Pharma
 242 Innovation. 2022; 11(6): 2322-2325.

243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258

UNDER PEER REVIEW

259 **Table 1. Efficacy of different biorationals on mycelial growth of the test pathogen,**
 260 ***Fusarium oxysporum* f. sp. *ricini* in vitro**

Tr. No.	Biorational Extracts	Radial growth (mm)* of pathogen at different Biorational concentration (%)	Radial growth Mean**	Per cent inhibition of pathogen over control (mm) * Concentrations (%) of	Per cent Inhibition Mean**
---------	----------------------	--	----------------------	---	----------------------------

					(mm)	biorationals			(%)
		10	15	20		10	15	20	
T₁	Ginger	67.67	42.00	52.00	53.89	24.81 (29.87)	53.33 (46.89)	42.22 (40.51)	40.12 (39.29)
T₂	Onion	51.67	61.33	73.67	62.22	42.59 (40.72)	31.85 (34.35)	18.15 (25.20)	30.86 (33.74)
T₃	Bougainville	52.67	52.67	73.00	59.44	41.48 (40.08)	41.48 (40.08)	18.89 (25.75)	33.95 (35.62)
T₄	Garlic	52.67	53.00	54.67	53.44	41.48 (40.08)	41.11 (39.86)	39.26 (38.78)	40.62 (39.58)
T₅	Turmeric	47.33	46.33	52.67	48.78	47.41 (43.50)	48.52 (44.13)	41.48 (40.08)	45.80 (42.58)
T₆	Neem	41.33	41.67	62.00	48.33	54.07 (47.32)	53.70 (47.11)	31.11 (33.89)	46.30 (42.86)
T₇	Chilli	58.33	43.33	44.00	48.56	35.19 (36.37)	51.85 (46.04)	51.11 (45.62)	46.05 (42.72)
T₈	Aloe vera	61.33	57.67	75.33	64.78	31.85 (34.35)	35.93 (36.81)	16.30 (23.80)	28.02 (31.95)
	Mean	54.13	49.75	60.92		43.33 (41.15)	44.72 (41.95)	32.31 (34.63)	
	Source	SE(d)		C. D. (p 0.01)		SE(d)		C. D. (p 0.01)	
	Biorational (B)	0.56		1.51		0.39		1.05	
	Concentration (C)	0.34		0.93		0.24		0.65	
	B × C	0.97		2.62		0.68		1.83	

261

262 ** Mean of the three replications

Table 2: *In vitro* evaluation of new fungicide molecules against *Fusarium oxysporum* f. sp. ricini

Tr.No.	Fungicide	Radial growth (mm)*				Radial growth Mean** (mm)	Per cent inhibition over control *				Per cent Inhibition Mean** (%)
		Concentration (ppm)					Concentration (ppm)				
		500	1000	1500	2000		500	1000	1500	2000	
T ₁	Iprovalicarb 5.5% + Propineb 61.25% WP	61.33	76.66	62.33	52.66	63.25	31.85 (34.35)	14.81 (22.63)	30.74 (33.66)	41.48 (40.08)	29.72 (33.02)
T ₂	Fluopyram 17.7% + Tebuconazole 17.7% SC	12.00	11.667	11.33	11.66	11.67	86.67 (68.56)	87.04 (68.87)	87.41 (69.19)	87.04 (68.87)	87.04 (68.87)
T ₃	Picoxystrobin 6.78% + Tricyclozole 20.33% SC	0.00	0.00	0.00	0.00	0.00	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)
T ₄	Boscalid 25.2 % + Pyraclostrobin 12.8 % WG	27.33	25.00	17.33	14.66	21.08	69.63 (56.54)	72.22 (58.17)	80.74 (63.94)	83.70 (89.96)	76.57 (61.03)
T ₅	Thiophanate-methyl 70% WP	0.00	0.00	0.00	0.00	0.00	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)
T ₆	Carbendazim 50% WP	0.00	0.00	0.00	0.00	0.00	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)	100.00 (89.96)
Mean		16.78	18.89	15.17	13.17		81.36 (64.39)	79.01 (62.71)	83.15 (65.74)	85.37 (63.57)	
Source		SE(d)		C. D. (p 0.01)			SE(d)		C. D. (p 0.01)		
Fungicide (F)		0.39		1.06			0.30		0.82		
Concentration (C)		0.32		0.86			0.25		0.67		
F × C		0.79		2.11			0.61		1.63		

**Mean of the three replications




Figure 1: 1A. *In vitro* evaluation of biorationals showing radial growth (mm) and mycelial inhibition (%)

Figure 1B. *In vitro* evaluation of new fungicide molecules against *Fusarium oxysporum f. sp. ricini*




Plate 1A. *In vitro* evaluation of biorationals against *Fusarium oxysporum f. sp. ricini*

Plate 1B. *In vitro* evaluation of new fungicide molecules against *Fusarium oxysporum f. sp. ricini*