

Impact of Bioagents on Growth and Yield Parameters of Root-knot Nematode Infested Mulberry Plants

ABSTRACT

Mulberry (*Morus alba* L.) is a vital crop for sericulture and its productivity is crucial for the industry's sustainability. This study investigated the influence of bioagents viz., *Purpureocillium lilacinum*, *Trichoderma harzianum*, *Trichoderma viride*, *Pochonia clamydosporea* and *Pseudomonas fluorescens*, alongside control treatments including carbofuran 3G and neem cake on growth and yield parameters in severely root-knot nematode infested mulberry plants. The treatments involved are generally recommended to manage root-knot nematode infestation. The results revealed that *T. viride* @ 5 kg/ha significantly improved plant growth, leaf area and leaf yield leading to enhanced productivity. The findings of this study underscore the potential of bioagents in fostering environmentally sustainable and economically viable sericulture.

Aims: The study aims to assess the impact of bioagents on growth and yield parameters of root-knot nematode, *Meloidogyne incognita* infested mulberry plants.

Study design: The data collected from the experimental field were analyzed statistically by using one-way RCBD.

Place and Duration of Study: The experiment was carried out in the root-knot nematode infested mulberry garden in Kalyapura Village, Shidlaghatta Taluk, Chikkaballapura District, Eastern Dry Zone (Zone-5) of Karnataka, India between January to September 2022, covering three harvests.

Methodology: The treatments were incorporated near the root zone of mulberry plants within a week after pruning as per the recommendations. The observations were recorded once before the imposition of the treatments (initial) and during the first (60 DAT) and second (120 DAT) shoot harvests in mulberry after imposing the treatments.

Results: Among the bioagents *Trichoderma viride* showed better results for shoot height (120.00 cm), number of leaves per plant (511.75), leaf area (102.10 cm²) and leaf yield per plant (528.38 g) during the second harvest i.e., 120 DAT. The minimum leaf yield per plant was recorded in the untreated check (216.43 g).

Conclusion: Incorporating *T. viride* @ 5 kg/ha mixed with 5 tons FYM near the root zone of mulberry infested with RKN showed promising results in promoting plant growth and yield.

Keywords: Mulberry, Root-knot nematode, Bioagents, Growth, Yield

1. INTRODUCTION

The silkworm, *Bombyx mori* L. is a monophagous lepidopteran insect that feeds exclusively on mulberry foliage. The perennial plant mulberry (*Morus alba* L.) is highly adaptable to varied climatic conditions ranging from temperate to tropics and thrives well under different soils. Foliage is the major economic part of mulberry that ultimately decides the quality of raw silk

20 since the silkworm feeds on mulberry leaf alone to derive its nutrients for growth and
21 productivity.

22

23 Among several factors in obtaining a successful cocoon crop, mulberry leaf alone contributes
24 to around 38.20 per cent that is followed by microclimate in the rearing house (37.00 %),
25 silkworm rearing techniques (9.30 %) and the breed (4.20 %), which signifies the role of quality
26 foliage in cocoon production [1]. Apart from soil parameters, the biotic and abiotic stress
27 factors greatly affect the quality of mulberry leaves. The nutritive values get degraded due to
28 diverse biotic stresses viz., diseases and pests and the mulberry plants attract these pests
29 and diseases due to their perennial, fast-growing and lush green characteristics. Plant-
30 parasitic nematodes, particularly root-knot nematode (RKN), *Meloidogyne incognita* (Kofoid
31 and White), pose a significant threat to mulberry productivity, causing substantial reductions
32 in herbage yield and leaf quality, as well as curtailing plant longevity. As a major pest of
33 mulberry, *M. incognita* infestations result in quantitative and qualitative losses, underscoring
34 the need for effective management strategies to mitigate their impact on this economically
35 important crop [2]. Root gall formation, accompanied by symptoms of stunted growth, chlorotic
36 discolouration and diminished plant vigour, characterizes the manifestation of this disease. *M.*
37 *incognita* alone is known to cause 20-50 per cent loss in mulberry leaf yield apart from
38 deteriorated quality [3].

39

40 The applications of synthetic pesticides in controlling RKNs have been found to induce highly
41 toxic residual effects on the environment, particularly on non-target organisms. Some are
42 mobile, volatile in the air and soluble in water [4]. Alternately, bio-pesticides have intrinsic
43 application in the control of plant pests and diseases, due to their eco-friendly nature and
44 bioavailability as opposed to synthetic pesticides [5]. The bioagents include parasites,
45 predators, fungi, bacteria, viruses, plant products and others that are specific and safe to non-
46 target species, beneficial insects, higher animals and the man with the least effects on the
47 environment and ecosystem. The use of fungal and bacterial bioagents has great potential in
48 nematode management and has a great influence on growth and yield parameters. The
49 bioagents viz., *Purpureocillium lilacinum*, *Trichoderma harzianum*, *Trichoderma viride*,
50 *Pochonia clamydosporia* and *Pseudomonas fluorescens* recommended against RKN were
51 assessed for their influence on growth and yield parameters of mulberry in the present
52 investigation.

53

54 2. EXPERIMENTAL DETAILS

55

56 The experiment was carried out in the RKN infested mulberry garden in Kalyapura Village,
57 Shidlaghatta Taluk, Chikkaballapura District i.e., in the Eastern Dry Zone (Zone-5) of
58 Karnataka, India at 13°14'20"N latitude and 77°52'17"E longitude, at an altitude of 904m above
59 mean sea level. The mulberry plantation selected for the study had red sandy loam type of
60 soil with six years old V1 variety plants, planted at the spacing of 90x90cm; bottom pruning
61 (Kolar method) was followed; the field was irrigated in two-day intervals with drip irrigation
62 facility i.e., based on soil moisture conditions; organic manures and inorganic fertilizers were
63 applied following the recommended package of practice. Except for the management of RKN,
64 the selected field was well maintained.

65

66 The following bioagents were selected based on the reviews and were used for the
67 management of RKN in mulberry. The selected bioagents were,

68

- 69 a) *Purpureocillium lilacinum*
- 70 b) *Trichoderma harzianum*
- 71 c) *Trichoderma viride*
- 72 d) *Pochonia clamydosporia*

73 e) *Pseudomonas fluorescens*

74

75 Along with the five bioagents, Nemahari (bio-nematicide developed by CSRTI, Mysore), neem
76 cake, carbofuran 3G and control (untreated check) were included in the treatments for
77 comparison of the efficacy.

78

79 List 1: Treatment details

80

Treatments		Recommendation
T ₁	<i>Purpureocillium lilacinum</i>	5 kg/ha (with 5 tons FYM) [6]
T ₂	<i>Trichoderma harzianum</i>	
T ₃	<i>Trichoderma viride</i>	
T ₄	<i>Pochonia clamydosporea</i>	
T ₅	<i>Pseudomonas fluorescens</i>	
T ₆	Nemahari	40 kg/ha (with 400 kg FYM) [7]
T ₇	Neem cake	2000 kg/ha [8]
T ₈	Carbofuran 3G (Standard check)	40 kg/ha [8]
T ₉	Control (Untreated check)	-

81

82 The treatments were incorporated in four replications each near the root zone of mulberry
83 plants within a week after pruning as per the recommendations.

84

85 2.2 Observations

86

87 The growth and yield parameters viz., shoot height, number of shoots per plant, number of
88 leaves per plant, leaf area and leaf yield per plant were recorded once before the imposition
89 of the treatments (initial) and during the first (60 DAT) and second (120 DAT) shoot harvests
90 in mulberry after imposing the treatments.

91

92 Shoot height was recorded from the base of the shoot to the topmost fully opened leaf in
93 randomly selected five shoots of each replication of every treatment. The mean height of five
94 shoots was worked out to obtain the shoot height of the replication. The number of shoots in
95 each replication (plant) was counted and recorded as the number of shoots per plant. The
96 total number of leaves in selected five shoots of each replication was counted separately and
97 the mean was calculated, then multiplied by the number of shoots per plant. The fully opened
98 third, fifth and seventh leaves from the top were plucked separately from the selected five
99 shoots of each replication and their area was determined using the leaf area meter. The leaf
100 yield per plant was determined by multiplying the number of shoots per plant with the leaf
101 weight in grams per shoot.

102

103 2.3 Statistical analysis

104

105 The data collected from the experimental field were analyzed statistically by using one-way
106 RCBD for testing of significance by Fisher's method of analysis of variance [9]. The level of
107 significance used in the F-test was P = 0.05.

108

109 **3. RESULTS AND DISCUSSION**

110

111 The efficacy of different bioagents on various growth and yield attributes of mulberry viz., shoot
 112 height (cm), number of shoots per plant, number of leaves per plant, leaf area (cm²), leaf
 113 weight per shoot (g) and leaf yield per treatment (g) were recorded once before the imposition
 114 of treatments (initial), intermediary (i.e., first shoot harvest - 60 DAT) and final (i.e., second
 115 shoot harvest - 120 DAT).

116

117 **3.1 Shoot height**

118 The observations reveal that the shoot height was found to be unwavering before the
 119 imposition of treatments (initial). The observations among the bioagents reveal that on 60
 120 DAT, the lengthiest shoot was recorded in **Nemahari (153.50 cm) on par with *T. viride* (152.75**
 121 **cm)** followed by *P. lilacinum* (145.25 cm), *T. harzianum* (140.75 cm), *P. fluorescens* (139.50
 122 cm), neem cake (135.75 cm), *P. clamydosporia* (132.25 cm) and carbofuran 3G (125.50 cm).
 123 The shortest shoot was observed in the untreated check (108.00 cm).

124 The lengthiest shoot among the treatments on 120 DAT was recorded in Nemahari (193.00
 125 cm) on par with *T. viride* (192.00 cm) followed by *P. lilacinum* (181.25 cm), *T. harzianum*
 126 (176.25 cm), *P. fluorescens* (174.50 cm), *P. clamydosporia* (166.75 cm), neem cake (166.50
 127 cm) and carbofuran 3G (152.75 cm). The shortest height of the shoot was recorded in the
 128 untreated check (109.75 cm) (Table 1).

129

130 **3.2 Number of shoots per plant**

131 Among the treatments, except for the control the number of shoots per plant was found to be
 132 consistent before imposition of treatments (initial) and also on 60 DAT without any significant
 133 difference. The maximum number of shoots per plant among the treatments on 120 DAT was
 134 recorded in the untreated check (22.00) followed by Nemahari (20.50), *P. clamydosporia*
 135 (18.25), *P. lilacinum* (17.75), carbofuran 3G (17.75), *T. harzianum* (17.50), neem cake (17.00),
 136 *T. viride* (16.25) and *P. fluorescens* (15.75) (Table 1).

137 It was observed that the control showed stunted plant growth which might have resulted in
 138 yielding higher number of stunted shoots. The increase in the number of healthy shoots in the
 139 treatment plot involving Nemahari might be a result of its manurial composition.

140

141 **3.3 Number of leaves per plant**

142 The number of leaves per plant was found to be consistent in all treatments before the
 143 imposition of treatments (initial). The observations reveal that on 60 DAT, the maximum
 144 number of leaves per plant was recorded in **Nemahari (448.50) on par with *T. viride* (445.75)**
 145 followed by *P. lilacinum* (439.75), *T. harzianum* (431.00), neem cake (413.50), *P.*
 146 *clamydosporia* (409.25), *P. fluorescens* (380.50) and carbofuran 3G (327.75). The least was
 147 observed in the untreated check (236.00).

148 The maximum number of leaves per plant among the treatments on 120 DAT was recorded
 149 in *T. viride* (511.75) on par with Nemahari (515.50) followed by *P. lilacinum* (486.25), *T.*
 150 *harzianum* (463.50), neem cake (461.25), *P. clamydosporia* (452.25), *P. fluorescens* (450.00)
 151 and carbofuran 3G (401.75). The least number of leaves per plant was observed in the
 152 untreated check (265.25) (Table 1).

153

154 **Table 1. Effect of bioagents used against RKN on the growth parameters of mulberry**

155

Treatments	Shoot height (cm)			Number of shoots per plant			Number of leaves per plant		
	Initial	60 DAT	120 DAT	Initial	60 DAT	120 DAT	Initial	60 DAT	120 DAT
T ₁	109.25	145.25	181.25	18.50	18.75	17.75	235.75	439.75	486.25

T ₂	109.75	140.75	176.25	17.00	18.25	17.50	236.00	431.00	463.50
T ₃	109.00	152.75	192.00	17.25	17.75	16.25	235.75	445.75	511.75
T ₄	109.50	132.25	166.75	17.50	18.50	18.25	237.00	409.25	452.25
T ₅	109.75	139.50	174.50	17.00	17.25	15.75	237.75	380.50	450.00
T ₆	109.75	153.50	193.00	17.25	21.25	20.50	236.00	448.50	515.50
T ₇	109.00	135.75	166.50	18.25	18.25	17.00	238.75	413.50	461.25
T ₈	109.25	125.50	152.75	17.50	17.75	17.75	236.75	327.75	401.75
T ₉	109.25	108.00	109.75	18.25	21.50	22.00	238.25	236.00	265.25
F-test	NS	*	*	NS	NS	*	NS	*	*
SEm ±	-	6.09	5.88	-	-	0.74	-	44.96	31.49
CD_{0.05}	-	17.77	17.16	-	-	2.18	-	131.24	91.92

DAT – Days after treatment imposition; NS – Non-significant; * - Significant

156
157

3.4 Leaf area

158

159

The leaf area of mulberry was found to be consistent in all treatments before the imposition of treatments (initial). The observations reveal that on 60 DAT, the maximum leaf area was recorded in *T. viride* (78.72 cm²) on par with Nemahari (79.39 cm²) followed by *P. lilacinum* (63.52 cm²), *T. harzianum* (62.98 cm²), *P. fluorescens* (60.42 cm²), neem cake (58.38 cm²), *P. clamydosporia* (58.14 cm²) and carbofuran 3G (54.94 cm²). The minimum leaf area was observed in the untreated check (36.68 cm²).

165

166

167

168

The maximum leaf area among the treatments on 120 DAT was recorded in Nemahari (104.08 cm²) on par with *T. viride* (102.1 cm²) followed by *P. lilacinum* (96.43 cm²), *T. harzianum* (94.80 cm²), *P. fluorescens* (90.44 cm²), neem cake (89.39 cm²), *P. clamydosporia* (87.21 cm²) and carbofuran 3G (74.08 cm²). The minimum leaf area was observed in the untreated check (35.50 cm²) (Table 2).

169

170

3.5 Leaf yield per plant

171

172

173

174

175

176

177

The mulberry leaf yield per plant was found to be consistent before the imposition of treatments (initial). The observations reveal that on 60 DAT, the maximum leaf yield per plant was recorded in Nemahari (478.78 g) on par with *P. lilacinum* (471.97 g) followed by *T. viride* (454.12 g), *T. harzianum* (435.28 g), neem cake (410.25 g), *P. fluoroscens* (384.62 g), *P. clamydosporia* (380.16 g) and carbofuran 3G (339.71 g). The minimum leaf yield per plant was recorded in the untreated check (217.00 g).

178

179

180

181

182

The maximum leaf yield per plant among the treatments on 120 DAT was recorded in Nemahari (531.92 g) on par with *T. viride* (528.38 g) followed by *T. harzianum* (509.32 g), *P. lilacinum* (489.07 g), *P. fluoroscens* (467.28 g), neem cake (465.13 g), *P. clamydosporia* (456.27 g) and carbofuran 3G (397.96 g). The minimum leaf yield per plant was recorded in the untreated check (216.43 g) (Table 2).

183

184

185

186

187

188

189

190

191

192

The increased plant growth and yield parameters might be due to the suppression of RKN population at the infection site preventing the nematode entry and multiplication in the mulberry root [10]. The positive impact of *T. viride* on growth and yield parameters of mulberry might be due to its inhibitory effect and reduction of galls and egg masses [11] [12]. *T. viride* in combination with organic amendments was also known to produce growth hormones, which were observed to have an added response in boosting the plant vigour [13]. The observations on shoot length, root length, shoot weight and root weight in cowpeas treated with bioagents revealed that *T. harzianum* @ 10 g/kg seed recorded maximum compared to *T. viride* and *P. fluorescens* @ 5, 7.5 and 10 g/kg seed [14]. The evaluation of the combination of chemical

193 nematicide, bioagent, oilcake and trap crop in managing the RKN in mulberry revealed that
 194 carbofuran + *T. viride* performed better than all other combinations in terms of nematode
 195 population reduction and enhancement of growth and yield parameters of mulberry [15].
 196

197
 198

Table 2. Effect of bioagents used against RKN on yield parameters of mulberry

Treatments	Leaf area (cm ²)			Leaf yield per treatment (g)		
	Initial	60 DAT	120 DAT	Initial	60 DAT	120 DAT
T ₁	39.76	63.52	96.43	222.15	471.97	489.07
T ₂	38.52	62.98	94.8	220.94	435.28	509.32
T ₃	38.57	78.72	102.1	221.16	454.12	528.38
T ₄	38.24	58.14	87.21	218.85	380.16	456.27
T ₅	37.29	60.42	90.44	217.22	384.62	467.28
T ₆	38.9	79.39	104.08	219.38	478.78	531.92
T ₇	38.18	58.38	89.39	224.51	410.25	465.13
T ₈	37.06	54.94	74.08	219.9	339.71	397.96
T ₉	37.26	36.68	35.5	220.68	217	216.43
F-test	NS	*	*	NS	*	*
SEm ±	-	1.78	2.14	-	44.2	33.12
CD_{0.05}	-	5.21	6.25	-	129.02	96.68

DAT – Days after treatment imposition; NS – Non-significant; * - Significant

199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215




Fig. 1. Effect of bioagents on mulberry leaf yield per plant (g)

216
 217
 218
 219
 220
 221
 222
 223
 224

4. CONCLUSION

The study demonstrated the efficacy of bioagents used against RKNs in enhancing growth and yield parameters of mulberry. Among the bioagents, incorporating *T. viride* @ 5 kg/ha mixed with 5 tons FYM near the root zone showed promising results in promoting plant growth and yield. The integration of bioagents with organic amendments can further enhance their effectiveness. Future research should focus on exploring the potential of bioagents in

225 combination with other compatible bioagents, to develop sustainable management strategies
226 for RKNs in mulberry. Additionally, investigating the mechanisms of action of bioagents and
227 their impact on soil microbiome can provide valuable insights into their role in plant health and
228 ecosystem functioning. The findings of this study have significant implications for the
229 development of environmentally friendly and sustainable sericulture practices.

230

231 **CONFLICT OF INTEREST**

232

233 None.

234

235 **ACKNOWLEDGEMENTS**

236

237 The authors express their sincere gratitude to the Nematology Department, IIHR-Bangalore,
238 for providing the necessary bioagents on time. We also appreciate the farmer's willingness to
239 participate and his cooperation in facilitating the experiment on his farm, which was
240 instrumental in the success of this study.

241

242 **COMPETING INTERESTS**

243

244 Authors have declared that no competing interests exist.

245

246 **AUTHORS' CONTRIBUTIONS**

247

248 'Bharath KB' conducted the experiment, recorded observations, analyzed the data, performed
249 the statistical analysis, wrote the protocol and wrote the manuscript. 'Vinoda KS',
250 'Banuprakash KG' 'Shashidhar KR' were involved in experimental site selection and planning
251 the study design. 'Kavitha TR' managed the selection of treatment and provided insights on
252 RKN. 'Naveenchandra Reddy' helped to record field-level observations. All authors read and
253 approved the final manuscript.

254

255 Disclaimer (Artificial intelligence)

256 Option 1:

257 Author(s) hereby declare that NO generative AI technologies such as Large Language Models
258 (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or
259 editing of manuscripts.

260 Option 2:

261 Author(s) hereby declare that generative AI technologies such as Large Language Models,
262 etc have been used during writing or editing of manuscripts. This explanation will include the

263 name, version, model, and source of the generative AI technology and as well as all input
264 prompts provided to the generative AI technology

265 Details of the AI usage are given below:

266 1.

267 2.

268 3.

269

270 REFERENCES

271

272 1. Miyashita Y. A report on mulberry cultivation and training methods suitable to bivoltine
273 rearing in Karnataka. Central Silk Board, Bangalore, India. 1986:1-7.

274

275 2. Muthulakshmi M, Devrajan K, Jonathan EI. Biocontrol of root-knot nematode, *Meloidogyne*
276 *incognita* (Kofoid and White) Chitwood in mulberry (*Morus alba* L.). *J*
277 *Biopesticides*. 2010;**3**(2):479-482.

278

279 3. Arunakumar GS, Revanna S, Kumar V, Yadav VK, Sivaprasad V. Studies on scanning
280 electron microscopy and fungal association with root-knot nematode in major mulberry
281 growing areas of Southern Karnataka. *J Entomol Zool Stud*. 2018;**6**(4):511-518.

282

283 4. Gapasin RM, Vasquez EA, Rendon GA. Bioassay-guided identification of the nematocidal
284 secondary metabolites from *Paecilomyces lilacinum* for the control of root-knot nematode
285 (*Meloidogyne graminicola*, Golden and Birchfield). *Ann Trop Res*. 2011;**33**(2): 22-43.

286

287 5. Helen OI, Godwin PA, Sidney ON. The Effects of Some Bio-Nematicides on the Productivity
288 of *Capsicum annum*. *J Environ Treat Tech*. 2016;**4**(4):111-117.

289

290 6. Saxena AK, Chakdar H, Kumar M, Rajawat MVS, Dubey SC, Sharma TR. 2021, ICAR
291 Technologies: Biopesticides for Eco-friendly Pest Management. Indian Council of Agricultural
292 Research, New Delhi. 2021:23-27.

293

294 7. Nishita Naik V, Pratheeshkumar PM. Nemahari - A plant based product for management of
295 root-knot disease in mulberry. Central Sericultural Research and Training Institute, Mysore,
296 India. 2015;(1): (*Tech Bull*).

297

298 8. Dandin SB, Giridhar K, editors. Handbook of Sericulture Technologies. Central Silk Board,
299 Bangalore, India. 2014.

300

301 9. Snedecor GW, Cochran WG. Statistical Methods. Iowa State University Press, Ames, Iowa.
302 1979.

303

304 10. Kshirsagar RV. Studies on effectiveness of Nemahari against root-knot nematode disease
305 *Meloidogyne incognita* of mulberry to reduction of cocoon yield losses of bivoltine silkworm in
306 Pune region. *Dnyanamay J*. 2017;**3**(1):13-15.

307

308 11. Narasimhamurthy TN. Evaluation of indigenous and commercial bioagents for
309 management of root-knot nematode, *Meloidogyne incognita* (Kofoid and White) Chitwood
310 1949 in mulberry (*Morus alba* L.). University of Agricultural Sciences, Bangalore, India.
311 2010:1-122 (*PhD Thesis*).

312

313 12. Bharath KB, Vinoda KS, Banuprakash KG, Kavitha TR. Evaluation of bioagents for their
314 efficacy in management of root-knot nematode infesting mulberry. *Int J Environ Agric Res.*
315 2024;10(7):1-7.

316

317 13. Chang YC, Baker R, Kleifeld A, Chet I. Increased growth of plants in the presence of the
318 biological control agent, *Trichoderma harzianum*. *Plant Disease*, 1986;70:145-148.

319

320 14. Nama CP, Sharma HK. Bio-management of root-knot nematode, *Meloidogyne incognita*
321 on cowpea (*Vigna unguiculata* L.). *J Entomol Zool Stud.* 2017;5(3):50-52.

322

323 15. Zaheer Ahamed B. Studies on the root-knot nematode *Meloidogyne incognita* (Kofoid and
324 White) Chitwood on Mulberry (*Morus alba* L.) and its management. University of Agricultural
325 Sciences, Bangalore, India. 2001:122 (*PhD Thesis*).

326