

Evaluating the Response of Watermelon (*Citrullus lanatus* (L.) to Organic and Inorganic Fertilization in Makurdi, Southern Guinea Savannah, Nigeria

ABSTRACT

Watermelon consumption in Nigeria is on the increase due to increasing awareness of its nutritional value while its production is on the decrease due to inefficiency of resource management (inputs like fertilizers). This study aims to optimize watermelon (*Citrullus lanatus* L. production through the use of organic and inorganic fertilizers. The field experiment was conducted at the Teaching and Research Farm of Joseph Sarwuan Tarka University, Makurdi in the 2019 cropping season under rain-fed conditions. The site falls within the Latitude 7°4'1N and Longitude 8°3'7E at an elevation of about 97m above sea level in the Southern Guinea Savanna Agro-ecological Zone of Nigeria. The experiment treatments, namely and they were T1 (Control 0kg/ha), T2 (150 kg/ha Compost), T3 (200 kg/ha Compost), T4 (250 kg/ha Compost) and T5 (300 kg/ha NPK 20:10:10) laid out in Randomized Complete Block Design with three replications. The study revealed that in comparison with control treatments, 300 kg/ha NPK 20:10:10 produced (31.04t/ha) fruit yield, the application of 250 kg/ha compost enhanced growth and yield parameters of watermelon with fruit yield of (17.65t/ha). The research findings suggest that using compost manure at the rate of 250 kg/ha for watermelon production is viable since it is economical and environmentally friendly alternative to NPK fertilizer due to it cheap and easy to prepare methods, its beneficial effects on soil, groundwater and the environment. It is further recommended that, in Makurdi, smallholder farmers, should adopt the application of compost at the rate of 250 kg/ha for optimum yield

Keywords: Watermelon, organic, inorganic, growth and yield

1. INTRODUCTION

“Watermelon (*Citrullus lanatus* (L.), a member of the *Cucurbitaceae* family, is widely cultivated and the most consumed cucurbit in the world” (Huh *et al.*, 2008; Huh *et al.*, 2014; Rolnik and Olan, 2020). “In Nigeria, watermelon is mostly cultivated in the Northern part of the country because of its warm loving nature, which makes its production seasonal in other parts of the country” (Charles, 2005). “The crop is a very important fruit that contains nutrients and phytochemicals that have been reported to be of high benefit to human health” (Ijah *et al.*, 2015). “The fruit is said to be a good source of vitamins B, C, and E as well as minerals such as phosphorus, magnesium, calcium and iron” (Romdhane *et al.*, 2017). “Watermelon is mostly cultivated for its vegetative and fruit parts” (Eifediyi *et al.*, 2017) and accounts for 6.8% of the world’s vegetable production (Goreta *et al.*, 2005). According FAOSTAT (2022), total watermelon production was 99,958 million tons, this was 2.21% less than in the previous year and 3.77% more than 10 years ago. China, tops the 10 ranked countries and accounted for 60.4% of watermelon production in the world. Other top rank countries of production include Turkiye, India, Algeria, Brazil, Russia, Pakistan, USA, Senegal and Uzbekistan (FAOSTAT, 2022). “The crop is said to have its optimum production when planted on well-drained fertile sandy or sandy loamy soils” (Audi *et al.*, 2013) and on soils with high organic matter content (Lawal, 2000), “especially in this part of the country (the north central region), where soil is said to have low

fertility" (Abdel *et al.*, 2005). "In order to meet the high demands of watermelon, it is important to boost the nutrient status of the soil, and one of the ways of increasing the nutrient status is by boosting the soil nutrient content either with the use of organic materials such as poultry manure, animal waste, and compost or with inorganic fertilizers" (Dauda *et al.*, 2005). "The nutrient requirement for the whole production period of the crop is in the ratio of N: P₂O₅: K₂O of 3.28: 1: 4.33" (Audi, *et al.*, 2013). "The amount of N, P₂O₅ and K₂O as suggested by researchers, needed by the crop depends on the growth stage, as the seedling stage requires 0.0003, 0.0008, and 0.0022 g/plant, while the fruiting stage requires 0.44, 0.127, and 0.624 g/plant per day, respectively" (Audi *et al.*, 2013).

"The major constraint to crop production in Nigeria, especially in the North Central region is the low organic matter content in the soil, which is a result of constant bush burning and subsequent erosion" (Eifediyi *et al.*, 2017). "The farmers in this particular area, mostly peasant farmers, have resorted to the continual usage of inorganic fertilizers and this act causes soil acidity, thereby reducing fruit and vegetative proliferation" (John *et al.*, 2004). "Harnessing the energy from biological waste is important for crop production; when these wastes are recycled and used as manure for agriculture, it reduces the pollution of streams and rivers" (Balasubramanian and Palaniappan, 2003).

"Research has shown that mineral fertilizers alone cannot increase the production of the crop optimally, and there is a need for mineral fertilizers to be augmented with organic manure" (Roe, 1998). "Generally, high nitrogen fertilizer and high temperature promote the production of male flowers of the fruit and reduce the production of the female flowers, thereby resulting in less fruit set" (Chedha, 2006). "Soil amendments using both organic and inorganic fertilizers have been reported to improve of soil conditions and enhance plant growth by replenishing of soil nitrogen" (Audi *et al.*, 2013). "The combined effect of mineral fertilizers and organic matter at 40kg P₂O₅/ha, 60kg N/ha, and 2.7kg K₂O/ha increased millet grain yield by 87% and cowpea grain yield by 200% in a continuous cropping system" (Moutonnet, 2000). **Therefore, the objective of the present study is to evaluate the growth and yield response of watermelon to organic and inorganic fertilization in Makurdi.**

2. MATERIALS AND METHODS

2.1 Experimental Location

The field experiment was conducted at the Teaching and Research Farm of the Federal University of Agriculture, Makurdi, during the 2019 cropping season under rain-fed conditions. The site falls within the Latitude 7°4'1N and Longitude 8°3'7E at an elevation of about 97m above sea level in the Southern Guinea Savanna Agro Ecological Zone of Nigeria **during the period of August to November, 2019.**

2.2 Experimental Soils

The **experimental soils were** sandy loamy with a slightly acidic pH of 6.54 (Table 1). Before planting, the top soil (0-15cm) samples were collected at different points with a soil auger for physicochemical characterization following routine procedures.

2.3 Cultural Practices

Experimental ridges were constructed manually using hoe and the field was demarcated into plots measuring 20 m². Three seeds were sown per hole **on the 15th of August, 2019** and were later thinned to two seeds per stand at 3 **weeks after planting (WAP)**. **Weeding was carried out when necessary. Harvesting was done with the fruits attains market maturity at 3 months after planting (MAP).**

2.4 Compost Preparations

The crops residues of some cereals and legumes with potentials of being used as compost materials grown in the area were selected. The materials include; rice straws, cowpea haulms, groundnut haulms, millet stovers, sorghum stovers, maize straws and soybean haulms. The plant samples were collected, grounded and separately kept in a labeled polythene bags and used for analysis. All samples were analyzed in a laboratory using standard procedures.

2.5 Experimental Design

The experimental design was a Randomized Complete Block Design (RCBD) that was replicated three times. The experiment consists of five treatments, control inclusive, namely T1 (150 kg/ha compost), T2 (200 kg/ha compost), T3 (250 kg/ha compost), T4 (300 kg/ha NPK 20:10:10), and T5 (control) laid out in a randomized complete block design (RCBD) replicated three times. The growth parameters recorded were vine length and the number of leaves per plant at 4, 8, and 12 weeks after planting (WAP), while the number of fruits, fruit weight, and fruit yield were collected after harvest.

2.6 Statistical Analysis

All data collected were subjected to Analysis of Variance (ANOVA) using the GENSTAT statistical package (2015), and the means were separated using Fishers Least Significant Difference (F-SLD) at the 5% level of probability.

3. RESULTS AND DISCUSSION

3.1 Effect of organic and inorganic fertilizers on soil physico-chemical properties

The pre-planting soil data (Table 1) indicated that the soil was slightly acidic before planting. Table 2 shows the effects of organic and inorganic fertilizers on the soil properties. After various applications, the results showed that the lowest pH (6.41) was obtained from the control treatment, and the highest soil pH was obtained from the application of 250 kg/ha of compost. The soil organic matter increased after planting across all treatments, with the application of 250 kg/ha compost providing the highest soil organic matter content and the least soil organic matter content from the untreated treatment. Total Nitrogen in the soil increased with the application of fertilizers; the highest nitrogen concentration was obtained with the application of 250 kg/ha compost, giving a value of 0.25%, and the lowest recorded was with the application of 300 kg/ha NPK 20:10:10, giving a value of 0.22%. Application of 250 kg/ha compost gave the highest phosphorus concentration with a value of 1.94 mg/kg and application of 300 kg/ha NPK 20:10:10 gave the lowest concentration (1.75 mg/kg). The highest exchangeable cations (K, Ca, Mg, Na) were recorded from the application of 250 kg/ha compost, while the lowest values were recorded from the application of 300 kg/ha NPK 20:10:10 and control plots (Table 2). The low organic matter content of the experimental soil is characteristic of soils in the southern Guinea savannah of Nigeria, due to constant bush burning and high temperatures leading to a high rate of organic matter decomposition (Eifediyi *et al.* 2017). Akinrinade and Obigbesan (2000) reported an organic matter content of 3% as best for watermelon production in this area; thus, the low organic matter content invariably results in soil fragility in this environment. The above requires organic soil amendment in this area. Ahamefule *et al.* (2015) reported that organic matter improves soil microbial activities and soil-water holding capacity, among other roles. The significant improvement observed in the chemical properties, particularly in the organic matter and total nitrogen of the organically amended soil, against the decline in NPK-amended and control soils with time is indicative of structural enhancement following soil organic amendment.

Table 1: Soil Properties (0-15 cm) of the experimental site before Planting in Makurdi

Soil Parameters	Values
Sand (%)	73.90
Silt (%)	10.91
Clay (%)	15.19
Textural Class	Sandy Loam
pH (H ₂ O)	6.54
Organic matter (%)	1.68
Total Nitrogen (%)	0.24
Phosphorus (mg/kg)	2.75
Potassium (cmol/kg)	0.26
Calcium (cmol/kg)	0.29
Magnesium (cmol/kg)	2.53
Sodium (cmol/kg)	3.02
Exchangeable Acidity (cmol/kg)	1.12
Cation Exchange Capacity (CEC)	7.22

Table 2: Effects of organic and inorganic fertilizers on the soil properties after treatments during the 2019 cropping season in Makurdi

Treatment	pH (H ₂ O)	OM (%)	N (%)	P (mg/kg)	K (cmol/kg)	Ca (cmol/kg)	Mg (cmol/kg)	Na (cmol/kg)	EA (cmol/kg)	ECEC (cmol/kg)
T1	6.67	2.01	0.23	1.86	0.25	3.75	2.51	0.25	1.02	7.78
T2	6.69	2.04	0.25	1.92	0.27	3.77	2.53	0.27	1.10	7.94
T3	6.71	2.10	0.25	1.94	0.27	3.78	2.55	0.30	1.12	8.02
T4	6.60	1.98	0.22	1.71	0.20	3.70	2.50	0.18	1.12	7.70
T5	6.41	1.59	0.21	1.64	0.21	3.73	2.50	0.17	1.02	7.63

T1 = (150kg/ha compost),

T2 = (200kg/ha compost),

T3 = (250kg/ha compost),

T4 = (300kg/ha NPK 20:10:10)

T5 = (control)

3.2 Effect of organic and inorganic fertilization on the growth and yield of water melon

Table 3 shows the effect of organic and inorganic fertilizers on the number of leaves of watermelon. The highest number of leaves was obtained from plots treated with 250 kg/ha compost at 4 WAP, 300 kg/ha NPK 20:10:10 at 8 and 12 WAP. The difference was statistically significant across the various treatments; however, the number of leaves produced from the application of 250 kg/ha compost and 300 kg/ha NPK was at par, and no significant difference was observed. Generally, untreated plots (control) had the lowest number of watermelon after time of evaluation. The reason why the 250 kg/ha compost-treated plot outclassed the other rates may have been due to the comparatively faster decomposition rate and higher nutrient status. However, the number of leaves of watermelon in the study was similar to the findings of Aniekwe and Nwokwu, (2014); Eifediyi *et al.* (2017) but was fewer than that reported by Enujeke *et al.* (2013) using different rates of organic manure in Asaba, Nigeria.

Table 3: Effects of organic and inorganic fertilizers on the Number of Watermelon Leaves in Makurdi

Treatments	Number of Leaves		
	4 WAP	8 WAP	12 WAP
T1 = (150kg/ha compost),	12 b	18 c	19 c
T2 = (200kg/ha compost),	13 ab	19 b	21 b
T3 = (250kg/ha compost),	14 a	19 b	22 a
T4 = (300kg/ha NPK 20:10:10)	13 ab	20 a	22 a
T5 = (control)	11 c	15 d	14 d
F-LSD(P=0.05)	1.78	0.51	0.50

In a column, means with similar letter (s) are not significantly different by LSD at 5% level of significance
WAP- Weeks after planting

Table 4 shows effects of organic and inorganic fertilizers on plant length, which indicated that the longest primary lengths were obtained from plots treated with 300 kg/ha NPK 20:10:10 at 4, 8, and 12 WAP, and the difference was significant. The findings collaborate with those of Adeyeye *et al.* (2016) but are shorter in the response of watermelon to organic and inorganic fertilizers in Ogbomoso, Oyo State, Nigeria. The earlier reason adduced for the observation in the growth parameters may also have been responsible for that in the yield parameters; this is because the yield components followed a similar trend with the growth parameters.

Table 4: Effects of organic and inorganic fertilizers on the Plant length of watermelon in Makurdi

Treatments	Vine Length (cm)		
	4 WAP	8 WAP	12 WAP
T1 = (150kg/ha compost),	19.14 c	27.15 d	34.04 d
T2 = (200kg/ha compost),	19.61 c	28.07 c	36.67 c
T3 = (250kg/ha compost),	22.06 b	30.08 b	44.11 d
T4 = (300kg/ha NPK 20:10:10)	30.48 a	39.07 a	46.89 a
T5 = (control)	17.21 d	25.07 e	30.15 e
F-LSD(P=0.05)	0.58	0.14	1.10

In a column, means with similar letter (s) are not significantly different by LSD at 5% level of significance
WAP- Weeks after planting

Table 5 shows the effect of organic and inorganic fertilizers on yield parameters of watermelon, which indicated that the highest number of fruits, highest fruit weight, and highest fruit yield were attributed to the plots treated with the 300 kg/ha NPK 20:10:10 fertilizer, while the least were attributed to plots treated with control fertilizer (no application). The number of fruits increases with an increase in the rates of compost. The number of fruits per plant obtained in this study is lower than that reported by Eifediyi *et al.* (2017), who used organic and inorganic fertilizers in Illorin, north-central Nigeria. The yield (weight of fruits and yield (t/ha)) experienced in this experiment was also lower than that reported in a similar work by Enujoke, (2013) using different rates of poultry manure on watermelon and Eifediyi *et al.* (2017), who reported a yield range of 334-402 t/ha, 306-390 t/ha, and 38.25-59.20 t/ha for NPK, poultry manure, and control treatments, respectively. Similarly, several authors have reported higher yields of melon in fertilized plots either NPK or organic fertilizers as compared to control (untreated plots). (Aluko, 2020; Aluko *et al.*, 2020; Adeyeye *et al.*, 2016; Ojo *et al.*, 2014)

Table 5: Effects of organic and inorganic fertilizers on the yield parameters of Watermelon in Makurdi

Treatments	Number of Fruits	Fruit weight (kg/ha)	Fruit Yield (t/ha)
T1 = (150kg/ha compost),	8.0 d	2.28 a	17.47 b
T2 = (200kg/ha compost),	9.0 c	1.97 a	16.01 c
T3 = (250kg/ha compost),	10.0 b	2.28 a	17.65 b
T4 = (300kg/ha NPK 20:10:10)	11.0 a	2.74 a	31.04 a
T5 = (control)	8.0 d	1.64 b	13.79 d
F-LSD(P=0.05)	0.45	1.04	0.48

In a column, means with similar letter (s) are not significantly different by LSD at 5% level of significance

4. CONCLUSION

It can be concluded from this study that the use of compost manure at 250 kg/ha and 300 kg/ha of NPK fertilizer is appropriate for the cultivation of watermelon in Makurdi. However, NPK fertilizer is expensive and is generally beyond the reach of smallholder farmers. Therefore, the use of 250 kg/ha compost manure is advocated, bearing in mind its beneficial effects on the soil. **The combination of the organic compost and varied rates of NPK under different locations with the southern guinea savannahs and seasons (rainy and irrigated) is also recommended for further research.**

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

OPTION 1:

AUTHOR(S) HEREBY DECLARE THAT NO GENERATIVE AI TECHNOLOGIES SUCH AS LARGE LANGUAGE MODELS (CHATGPT, COPILOT, ETC) AND TEXT-TO-IMAGE GENERATORS HAVE BEEN USED DURING WRITING OR EDITING OF MANUSCRIPTS.

OPTION 2:

AUTHOR(S) HEREBY DECLARE THAT GENERATIVE AI TECHNOLOGIES SUCH AS LARGE LANGUAGE MODELS, ETC HAVE BEEN USED DURING WRITING OR EDITING OF MANUSCRIPTS. THIS EXPLANATION WILL INCLUDE THE NAME, VERSION, MODEL, AND SOURCE OF THE GENERATIVE AI TECHNOLOGY AND AS WELL AS ALL INPUT PROMPTS PROVIDED TO THE GENERATIVE AI TECHNOLOGY

DETAILS OF THE AI USAGE ARE GIVEN BELOW:

- 1.
- 2.
- 3.

REFERENCES

Adeyeye, A. S., Akanbi, W. Sobola, O.O., Lamidi, W. and Olalekan, K. K. (2016). Growth and fruit of watermelon (*Citrullus lanatus*) as influenced by compost and NpK fertilizers. FUW trends in Science and Technology Journal. Vol. 1(1): 80-83

- Abdel A. M., Eldesuki, R.M., Salman, S.R., Hussien, S.D (2005). Performance of some snap bean varieties as affected by different levels of mineral fertilizers. *Journal of Agronomy* Vol. 4 pp 242-247
- Ahamefule E. Henry, Nwokocha C. Chukwuma and Amana S. Mathew. (2015). Stability and hydrological modifications in a tilled soil under selected organic amendments in South-eastern Nigeria. *Albanian Journal of Agric. Sci.* 14:127-136.
- Akinrinade, E.A and Obigbesan, G.O. (2000). Evaluation of the fertility status of selected soils for crop production in five ecological zones of Nigeria. *Proceedings 25th Annual Conference of Soil Science Society of Nigeria, Ibadan*, Pp 279-288.
- Aluko, M. (2020). Sowing dates and fertilizers application on growth and yield of muskmelon (*Cucumis melo* L.) at Ado-Ekiti. *Asian Journal of Agricultural and Horticultural Research* Vol. 5(3): 11-21. <https://doi.org/10.9734/ajahr/2020/v5i330052>
- Aluko, M., Ayodele, O.J. and Salami, A. E. (2020). The response of muskmelon to varying rates of NPK 15-15-15 fertilizers application, planting dates and in Ado-Ekiti. *Academia Journal of Agricultural Research* 8(8): 211-220
- Aniekwe, G. and Nwokwu, L. (2014). Impact of Different Mulching Materials on the Growth and Yield of Watermelon (*Citrullus lanatus*) in *Abakaliki, Southeastern Nigeria Journal of Biology, Agriculture and Healthcare* 4: 22 – 30.
- Audi, W., Aguyoh, J.N. and Gao-Qiong, L (2013). Yield and Quality of Watermelon as affected by Organic and Inorganic Nitrogen Sources. *Asian Journal of Agriculture and Food Sciences*. Vol 1- Issue 4.
- Balasubramaniyan, P. and Palaniappan, S.P. (2003). Principles and practices of Agronomy. *Agrobios Jodhpur India*
- Charles, F. A. (2005). Watermelon breeder cucurbit breeding horticultural science. Retrieved jul 17.
- Chedha, K.L. (2006). Handbook of horticulture. Pp.226-229.
- Dauda, S.N., Aliyu, L., Chiezey, U.F. (2005). Effects of variety, seedling age and poultry manure on growth and yield of garden egg (*Solanum gilo* L.). *The Nigerian Academic Forum* 9(1): 88-95
- Eifediyi, K.E., Remison, S.U., Ahamefule, H.E., Azeez, K.O., Fesobi, P.O (2017). Performance of Watermelon (*Citrullus lanatus* (L.) in response to organic and NPK fertilizers. DOI: 10.1515/ausae-2017-0001
- Enujeke, E. C. (2013). Response of watermelon to five different rates of poultry manure in Asaba area of Delta State, Nigeria. *Journal of Agriculture and Veterinary Science*, Vol 5(2): 45-50.
- FAOSTAT (2022). Food and Agriculture Organization of the United Nations, Rome, Italy. <https://www.fao.org/faostat/en/#data>
- Huh, C. Y., Choi, H. S., Solmaz, I., Sari, N. and Kim, S. (2014). Morphological characterization of Korean and Turkish watermelon germplasm. *CNU Journal of Agricultural Science* 41: 309-314. <https://doi.org/10.7744/cnujas.2014.41.4.309>

Huh, C. Y., Solmaz, I. and Sari, N. (2008). Morphological characterization of Korean and Turkish watermelon germplasm. Pitrat M. (ed) Cucurbitaceae 2008, Proceedings of the 9 EUCARPIA meeting on genetics and breeding of Cucurbitaceae Avignon (France), May 21-24, 2008, pp 327-333

GENSTAT (2015). GENSTAT 17.0 Release 17.1DE, Discovery Edition 4, Lawes Agricultural Trust, Rothamsted Experimental Station.

Goreta, S., Perica, S., Dumicic, G., Bucan, L. and Zanic, K (2005). Growth and Yield of watermelon on Polyethylene Mulch with Different Spacing and Nitrogen Rates. *America Journal of Horticultural Science.*, Vol 40, no 2, pp. 366-369,

Ijah, U. J., Ayodele, H.S., Aransiola, S.A. (2015). Microbiological and some sensory attributed of Watermelon Juice and Watermelon Orange Juice. *Journals of Food Resources Science.* Vol 4, pp. 49-61. DOI: 10.3923/jfrs.2015.49.61.

Lawal, A.B. (2000). Response of cucumber to intercropping with maize and varying rates of farmyard manure and inorganic fertilizer, *Agriculture and Environment.* 2(1): 78-83.

Moutonnet, P (2000). Report of the second research co-coordinating meeting of the FAO/IAEA co-ordinated research project on managements of nutrients and water in rainfed and semi-arid areas for increasing crop production.

Nowaki, D., Hiyoshi, R., Filio, C.B., Arthur, De Faria, T., Rogerio, Wamser, Fernando A., Cortez, M., Waldir, J. (2017). Effect of Nitrogen Fertilization on Yield and Quality of Watermelon, CV. Mossoro, Vol. 30, no 1, pp. 161-171.

Ojo, J. A., Olowoake, A. A. and Obembe, A. (2014). Efficacy of organomineral fertilizer and un-amended compost on the growth and yield of watermelon (*Citrullus lanatus* Thamb) in Ilorin Southern Guinea Savanna zone of Nigeria. *Int. J. Recycl Org Waste Agricult*, 3:121-125. <https://doi.org/10.1007/s40093-014-0073-z>

Roe, N.E (1998). Compost utilization of vegetable and fruit crops. *Horticultural Science*, vol 33, no 6, pp 934-937

Romdhane, M.B., Haddar, A., Ghazala, I., Jeddou, K.B., Helbert, C.B., Ellouz-Chaabouni, S. (2017). Optimization of Polusaccharides Extraction from Watermelon Rinds: Structure, Functional and Biological Activities. *Food Chemisty* Vol. 216, pp. 355-364. DOI: 10.1016/j.foodchem.2016.08.056.