

APPLICATION OF LAPLACE TRANSFORM TO DERIVE EXPONENTIATED LOGISTIC AND INVERSE GAUSSIAN DISTRIBUTIONS.

Abstract

Raising a cumulative distribution function (cdf) or survival function to a power is a method of generalizing a distribution, known as exponentiated distribution. [1] summarized various generalizations of the logistic distribution. This work constructs the generalized exponentiated distributions for the logistic distribution using a beta generated distribution. Specifically, we introduce two new distributions: the Generalized Exponentiated Logistic Type I and Type II. The cdf and pdf of the standard logistic are shown as special cases of these exponentiated distributions. Additionally, we express these exponentiated distributions in terms of the Laplace transform. We derive the Laplace transform for the Generalized Inverse Gaussian (GIG), Inverse Gaussian (IG), and Gamma distributions, demonstrating that the reciprocal Inverse Gaussian is a special case of the GIG when $\lambda = \frac{1}{2}$. We also explore the behavior of the shapes of these new distributions with varying parameter values, highlighting their flexibility and applicability in modeling statistical data. Generalizing makes the logistic distribution flexible and tractable to be used in analysis of quantal response data and probit analysis. Since the generalized distributions have a wide range of skewness and kurtosis, they can be easily applied in studying robustness tests. [2] showed the application of generalized logistic distributions to modeling extreme movement of stock market data making it suitable for investors assessing high risk portfolio.

Key words:

Cumulative distribution function, Generalized Exponentiated Logistic, Laplace Transform.

1 Introduction

Adding one or more parameters to a probability distribution is called generalization. The purpose of generalization is to make a distribution more flexible and tractable. It is well known, in general, that a generalized model is more flexible than the ordinary model and it is preferred by many data analysts in analyzing statistical data. [3] proposed a new method of introducing an extra parameter to a family of distributions for more flexibility. [4] discusses various generalizations of the logistic distribution. One method of generalizing a distribution is to raise a cumulative distribution function (*cdf*) or a survival function to a power. The raised function is said to be exponentiated. [5] introduced a skewness parameter in generalization of logistic so that the generalized distribution can be used to model data exhibiting a unimodal density function. [6] introduced the Exponentiated Weibull (EW) distribution that allowed for a non-monotone hazard rate. [7] studied exponentiated exponential (EE) distribution which they named Generalized Exponential distribution. [8] summarizes 8 well known generators for distributions. The generalized distributions nest various sub-families of distributions.

The concept of **EXPONENTIATION** was applied by [9] when solving a differentiation equation in *cdf*. The *cdfs* obtained were raised to powers. Out of the 12 *cdfs* obtained, 9 were raised to powers. The three *cdfs* not raised to powers and the other raised ones are known as Burr I, Burr II, Burr III up-to Burr XII.

Exponentiated distributions can be obtained as special cases of a beta generated distribution introduced by [10]. [11] investigated the shapes of density and hazard rate functions for the exponentiated half logistic family of distributions. [12] applied the type I generalized logistic distribution to engineering data using the Bayesian approach. [13] modified the Weibull distribution and showed that its flexibility increased with increase in number of parameters. [14] modified the Laplace distribution using the exponentiation approach.

The objective of this study is to: derive generalized exponentiated logistic distributions from the beta generator and show explicitly the Laplace transform of the exponentiated standard logistic, generalized inverse Gaussian, inverse Gaussian and Gamma distributions respectively.

The paper is organized as follows; Section two shows how an exponentiated distribution is deduced from a beta generated distribution and applied to the logistic distribution. Section three shows how an exponentiated mixture is expressed in terms of Laplace transform of a mixing distributions. In section four the Laplace transforms of the generalized inverse Gaussian, inverse Gaussian and Gamma distributions are derived. The concluding remarks are in section five.

2 Generalized Exponentiated Distributions.

The derivation of Exponentiated generalized logistic distributions follows;

Let the *cdf* and **probability density function** (*pdf*) of the logistic distribution be defined as:

$$S(x) = \frac{e^{-x}}{1 + e^{-x}}, \quad -\infty < x < \infty \quad (2.1)$$

$$s(x) = \frac{e^{-x}}{(1 + e^{-x})^2}, \quad -\infty < x < \infty \quad (2.2)$$

[10] considered a *cdf* of the classical beta distribution as;

$$T[S(x)] = \int_0^x \frac{w^{a-1}(1-w)^{b-1}}{B(a,b)} dw \quad (2.3)$$

Since $0 \leq x \leq 1$, [10] replaced it by a *cdf* $S(x)$. Thus we have;

$$V(x) = T[S(x)] = \int_0^{S(x)} \frac{w^{a-1}(1-w)^{b-1}}{B(a,b)} dw, \quad 0 \leq x \leq 1 \quad (2.4)$$

Using equation (2.4), put $b = 1$ in (2.3) we obtain;

$$\begin{aligned} V_1(x) &= \int_0^{S(x)} aw^{a-1} dw = \frac{aw^a}{a} \Big|_0^{S(x)} \\ &= [S(x)]^a \end{aligned} \quad (2.5)$$

Differentiating (2.5) we obtain;

$$v_1(x) = s(x)a \left(S(x) \right)^{a-1}, \quad a > 0, \quad -\infty < x < \infty \quad (2.6)$$

when values of the parameter a are varied Equations (2.5) and (2.6) are exponentiated generators. They can be used to generate new distributions with $S(x)$ as the parent *cdf*. **Substituting equations (2.1) and (2.2) in equations (2.5) and (2.6) we obtain;**

$$V_1(x) = \left(1 + e^{-x} \right)^{-a}, \quad a > 0, \quad -\infty < x < \infty \quad (2.7)$$

$$v_1(x) = \frac{ae^{-x}(1 + e^{-x})^a}{(1 + e^{-x})^3}, \quad a > 0, \quad -\infty < x < \infty \quad (2.8)$$

which is the **GENERALIZED EXPONENTIATED TYPE I LOGISTIC** distribution. When $a = 1$ in (2.7) and (2.8) we obtain (2.1) and (2.2) as special cases. Using (2.3), when $a = 1$, we obtain;

$$V_2(x) = \int_0^{S(x)} b(1-w)^{b-1} dw \quad (2.9)$$

Put

$$y = 1 - w \implies dy = -dw$$

Therefore, we proceed as follows:

$$\begin{aligned} V_2(x) &= \int_1^{1-S(x)} by^{b-1}(-dw) \\ &= - \left[\frac{by^b}{b} \right]_{1-S(x)}^1 \\ &= y^b \Big|_{1-S(x)}^1 \\ &= 1 - [1 - S(x)]^b \end{aligned} \tag{2.10}$$

$$\begin{aligned} 1 - V_2(x) &= [1 - S(x)]^b \\ v_2(x) &= b \left(1 - S(x) \right)^{b-1} s(x) \end{aligned} \tag{2.11}$$

Using the (2.1) and (2.2) in (2.10) and (2.11), we obtain the **Generalized Exponentiated Type II Logistic** as shown below;

$$V_2(x) = 1 - \left[\frac{e^{-x}}{1 + e^{-x}} \right]^b, \quad b > 0, -\infty < x < \infty \tag{2.12}$$

$$v_2(x) = \frac{be^{-xb}}{(1 + e^{-x})^{b+1}}, \quad b > 0, -\infty < x < \infty \tag{2.13}$$

[15] called (2.12) the Generalized logistic type I. [16] in his *Phd thesis* summarized five different methods of constructing the Generalized logistic type I. When $b = 1$ in (2.12), (2.1) is obtained as a special case.

2.1 Graph of the Exponentiated Generator

When varying parameters of a and b are plotted against the density, the graph of the exponentiated generator as shown in equation (2.6) is monotonically increasing resulting in a j- shape as shown below;




Figure 1:

2.2 Graph of the cdf of the standard logistic distribution

When varying parameters of a and b from equation (2.7) are plotted against the cdf, the graph of the cdf of standard logistic distribution is almost symmetrical with sharp peak indicating a higher concentration of data values hence the data is leptokurtic as shown in figure (2);




Figure 2:

2.3 Graph of the pdf of the standard logistic distribution

When $a = 1$ from equation (2.8) is plotted against the density, the graph of the pdf of standard logistic distribution is symmetrical as expected and is shown in figure (3);




Figure 3:

3 Laplace Transform of Exponentiated Distributions.

Let $G(x)$ be a *cdf* of a random variable X . It's power is denoted by $[G(x)]^\theta$ where θ is varying parameter. Then a continuous mixture of the raised *cdf* is given by;

$$F(x) = \int_0^\infty [G(x)]^\theta g(\theta) d\theta \quad (3.1)$$

where $g(\theta)$ is a continuous mixing distribution. Equation (3.1) can be expressed as a Laplace transform of a mixing distribution as shown below;

$$\begin{aligned} F(x) &= \int_0^\infty [G(x)]^\theta g(\theta) d\theta \\ &= \int_0^\infty \exp(\ln[G(x)]^\theta) g(\theta) d\theta \\ &= \int_0^\infty \exp(\theta \ln(G(x))) g(\theta) d\theta \end{aligned} \quad (3.2)$$

$$\begin{aligned}
&= \int_0^{\infty} \exp[-(-\theta \ln(G(x)))]g(\theta)d\theta \\
&= \int_0^{\infty} e^{-[\theta(-\ln(G(x)))]}g(\theta)d\theta \\
&= \int_0^{\infty} e^{-\theta(-\ln(G(x)))}g(\theta)d\theta \\
&= E[e^{-\theta(-\ln G(x))}] \\
&= L_{\theta}[-\ln G(x)]
\end{aligned} \tag{3.3}$$

which is the Laplace transform of the mixing random variable θ evaluated at $-\ln(G(x))$. **Substituting (2.1) in (3.3) we obtain;**

$$F(x) = L_{\theta}[\ln(1 + e^{-x})] \tag{3.4}$$

which is the Laplace transform of the logistic with the mixing random variable θ .

4 Laplace Transform of The Generalized Inverse Gaussian, Inverse Gaussian and Gamma Distributions.

We now wish to consider Laplace transform of generalized inverse Gaussian distribution and Gamma distribution.

4.1 Generalized Inverse Gaussian distribution

Generalized inverse Gaussian distribution (GIG) distribution is based on modified Bessel function of the third kind with index λ evaluated at ω denoted by $K_{\lambda}(\omega)$ and defined as;

$$K_{\lambda}(\omega) = \frac{1}{2} \int_0^{\infty} x^{\lambda-1} e^{-\frac{\omega}{2}(\frac{1}{x}+x)} dx \tag{4.1}$$

For $-\infty < \lambda < \infty$ and $\omega > 0$ [17] used the following parametrization

$$\omega = \frac{\mu}{\beta} \tag{4.2}$$

Therefore,

$$K_{\lambda}\left(\frac{\mu}{\beta}\right) = \frac{1}{2} \int_0^{\infty} x^{\lambda-1} e^{-\frac{\mu}{2\beta}(\frac{1}{x}+x)} dx \tag{4.3}$$

And the transformation

$$x = \frac{z}{\mu} \tag{4.4}$$

Substituting(4.4) in (4.3) we obtain;

$$K_{\lambda}\left(\frac{\mu}{\beta}\right) = \frac{1}{2} \int_0^{\infty} \left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{\mu}{2\beta}\left(\frac{z}{\mu} + \frac{\mu}{z}\right)} dz \quad (4.5)$$

$$= \frac{1}{2} \int_0^{\infty} \left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)} dz \quad (4.6)$$

Therefore,

$$1 = \int_0^{\infty} \frac{\left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)}}{K_{\lambda}\left(\frac{\mu}{\beta}\right)} dz \quad (4.7)$$

Thus the pdf of the Generalized Inverse Gaussian distribution according to [17] and transform is given by;

$$g(z) = \frac{\left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)}}{2K_{\lambda}\left(\frac{\mu}{\beta}\right)} \quad (4.8)$$

For $z > 0$; $-\infty < \lambda < \lambda$, $\mu > 0$, $\beta > 0$

Substituting (4.5) for the denominator in (4.8) we obtain;

$$g(z) = \frac{\left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)}}{\int_0^{\infty} \left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)} dz} \quad (4.9)$$

$$= \frac{z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)}}{\int_0^{\infty} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)} dz} \quad (4.10)$$

The Laplace transform of GIG distribution is given by;

$$L_z(s) = E(e^{-sz}) \quad (4.11)$$

$$= \int_0^{\infty} e^{-sz} g(z) dz \quad (4.12)$$

$$= \frac{\int_0^{\infty} e^{-sz} \left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{z}{\beta} + \frac{\mu^2}{\beta} \frac{1}{z}\right)} dz}{2K_{\lambda}\left(\frac{\mu}{\beta}\right)}$$

$$= \frac{\frac{1}{2} \int_0^{\infty} \left(\frac{1}{\mu}\right)^{\lambda} z^{\lambda-1} e^{-\frac{1}{2}(2s + \frac{1}{\beta})z + \frac{\mu^2}{\beta} \frac{1}{z}} dz}{K_{\lambda}\left(\frac{\mu}{\beta}\right)}$$

$$\begin{aligned}
&= \frac{\frac{1}{2} \int_0^\infty \left(\frac{1}{\mu}\right)^\lambda Z^{\lambda-1} e^{-\frac{1}{2}\left(\frac{2\beta s+1}{\beta}\right)z + \frac{\mu^2}{\beta} \frac{1}{2}} dz}{K_\lambda\left(\frac{\mu}{\beta}\right)} \\
&= \frac{\left(\frac{1}{\mu}\right)^\lambda}{K_\lambda\left(\frac{\mu}{\beta}\right)} \frac{1}{2} \int_0^\infty z^{\lambda-1} e^{-\frac{1+2\beta s}{2\beta}\left(z + \frac{\mu^2}{1+2\beta s}\right) \frac{1}{z}} dz
\end{aligned} \tag{4.13}$$

Let

$$z = \frac{\mu}{\sqrt{1+2\beta s}} t \implies dz = \frac{\mu}{\sqrt{1+2\beta s}} dt$$

Therefore,

$$L_Z(s) = \frac{(\mu^{-1})^\lambda}{K_\lambda\left(\frac{\mu}{\beta}\right)} \left(\frac{\mu}{\sqrt{1+2\beta s}}\right)^\lambda \frac{1}{2} \int_0^\infty t^{\lambda-1} e^{-\frac{\mu}{2\beta} \sqrt{1+2\beta s} \left(t + \frac{1}{t}\right)} dt \tag{4.14}$$

$$\begin{aligned}
&= \frac{(\mu^{-1})^\lambda}{K_\lambda\left(\frac{\mu}{\beta}\right)} \left(\frac{\mu}{\sqrt{1+2\beta s}}\right)^\lambda K_\lambda\left(\frac{\mu}{\beta} \sqrt{1+2\beta s}\right) \\
&= \left(\sqrt{1+2\beta s}\right)^{-\lambda} \frac{K_\lambda\left(\frac{\mu}{\beta} \sqrt{1+2\beta s}\right)}{K_\lambda\left(\frac{\mu}{\beta}\right)} \\
&= (1+2\beta s)^{-\frac{\lambda}{2}} \frac{K_\lambda(\mu\beta^{-1}(1+2\beta s))^{\frac{1}{2}}}{K_\lambda(\mu\beta^{-1})}
\end{aligned} \tag{4.15}$$

which is the Laplace transform of the GIG.

4.2 Laplace Transform of the Inverse Gaussian Distribution

The Laplace transform of inverse Gaussian distribution is obtained by putting

$$\lambda = -\frac{1}{2}$$

Therefore,

$$L_Z(s) = \left(\sqrt{1+2\beta s}\right)^{\frac{1}{2}} \frac{K_{-\frac{1}{2}}(\mu\beta^{-1}(1+2\beta s))^{\frac{1}{2}}}{K_{-\frac{1}{2}}(\mu\beta^{-1})} \tag{4.16}$$

Therefore,

$$L_Z(s) = \frac{\left(\sqrt{1+2\beta s}\right)^{\frac{1}{2}} \left[\frac{\pi}{2\mu\beta^{-1}(1+2\beta s)^{\frac{1}{2}}}\right]^{\frac{1}{2}} e^{-\mu\beta^{-1}(1+2\beta s)^{\frac{1}{2}}}}{\left(\frac{\pi}{2\mu\beta^{-1}}\right)^{\frac{1}{2}} e^{-\mu\beta^{-1}}} \tag{4.17}$$

Therefore,

$$\begin{aligned}
L_z(s) &= e^{\mu\beta^{-1} - \mu\beta^{-1}(1+2\beta s)^{\frac{1}{2}}} \\
&= e^{\frac{\mu}{\beta} [1 - \sqrt{1+2\beta s}]} \\
&= e^{-\frac{\mu}{\beta} [\sqrt{1+2\beta s} - 1]}
\end{aligned} \tag{4.18}$$

4.3 Laplace transform of Gamma distribution

Using equation (4.15) i.e. the Laplace transform of GIG, we have

$$L_z(s) = \frac{\int_0^\infty \left(\frac{1}{\mu}\right)^\lambda z^{\lambda-1} e^{-\frac{1}{2}(2s+\frac{1}{\beta})z} dz}{\int_0^\infty \left(\frac{1}{\mu}\right)^\lambda z^{\lambda-1} e^{-\frac{z}{2\beta}} dz} \tag{4.19}$$

$$\begin{aligned}
&= \frac{\int_0^\infty z^{\lambda-1} e^{-\frac{(1+2\beta s)z}{2\beta}} dz}{\int_0^\infty z^{\lambda-1} e^{-\frac{z}{2\beta}} dz} \\
&= \frac{\Gamma(\lambda) \left(\frac{1}{2\beta}\right)^\lambda}{\left(\frac{1+2\beta s}{2\beta}\right)^\lambda \Gamma(\lambda)} \\
&= \left(\frac{1}{1+2\beta s}\right)^\lambda \\
&= \left(\frac{\frac{1}{2\beta}}{s + \frac{1}{2\beta}}\right)^\lambda
\end{aligned} \tag{4.20}$$

which is the Laplace transform of a gamma distribution with parameters λ and $\frac{1}{2\beta}$.

4.4 Laplace Transform of The Inverse Gaussian

The Laplace transform of reciprocal inverse Gaussian distribution is obtained by putting $\lambda = \frac{1}{2}$ in equation (4.15) as follows;

$$L_z(s) = (\sqrt{1+2\beta s})^{-\frac{1}{2}} \frac{K_{\frac{1}{2}}(\mu\beta^{-1}(1+2\beta s)^{\frac{1}{2}})}{K_{\frac{1}{2}}(\mu\beta^{-1})} \tag{4.21}$$

$$\begin{aligned}
&= \frac{(\sqrt{1+2\beta s})^{-\frac{1}{2}} \left[\frac{\pi}{2\mu\beta^{-1}(\sqrt{1+2\beta s})} \right]^{\frac{1}{2}} e^{-\mu\beta^{-1}\sqrt{1+2\beta s}}}{\left(\frac{\pi}{\mu\beta^{-1}}\right)^{\frac{1}{2}} e^{-\mu\beta^{-1}}} \\
&= \frac{1}{\sqrt{1+2\beta s}} e^{-\mu\beta^{-1}[\sqrt{1+2\beta s}-1]} \\
&= \left(\frac{1}{1+2\beta s}\right)^{\frac{1}{2}} e^{-\mu\beta^{-1}(\sqrt{1+2\beta s}-1)} \tag{4.22}
\end{aligned}$$

Thus the Laplace transform of reciprocal inverse Gaussian distribution is the product of a Laplace transform of a gamma distribution with parameters $\frac{1}{2}$ and $\frac{1}{2\beta}$ and the Laplace transform of inverse Gaussian distribution.

5 Conclusion

In this work, we explicitly derived the generalized Exponentiated logistic type I and II distributions. We also showed how the logistic distribution can be obtained as a special case from the generalized distribution. If the *cdf* of a distribution is known, its exponentiated distributions can be obtained. The generalized logistic type I of Johnson 1995 has been derived and shown as the Exponentiated logistic type II. The graphs of the exponentiated generators are symmetrical and have a sharp peak. Further work can be done on generalizations by control the location and scale parameters. The generalized distributions obtained can also be applied to data.

We have also shown that, the Laplace transform of reciprocal inverse Gaussian distribution is the product of a Laplace transform of a gamma distribution with parameters $\frac{1}{2}$ and $\frac{1}{2\beta}$ and the Laplace transform of inverse Gaussian distribution. Therefore the generalized inverse Gaussian distribution nests other distributions which are obtained as special cases.

References

- [1] H. Omukami , P. Weke and J Ottieno (2021). Generalized Logistic Distributions Based on Various Methods for Constructing the Standard Logistic Distribution. *Advances and Applications in Mathematical Sciences*, Vol. 20 (10), pp. 2371.
- [2] Gettinby Gareth D, Sinclair C Donald, Power David M and Brown Richard A (2004). An analysis of the distribution of extreme share returns in the UK from 1975 to 2000. *Journal of Business Finance & Accounting*, Vol. 21 (5-6), pp. 607-646.

- [3] Mahdavi Abbas and Kundu Debasis (2017). A new method for generating distributions with an application to exponential distribution'. *Communications in Statistics-Theory and Methods, Taylor & Francis*, Vol. 46 (13), pp. 6543–6557.
- [4] Nassar MM and Elmasry A (2012). A study of generalized logistic distributions. *Journal of the Egyptian Mathematical Society, Elsevier*, Vol. 20 (2), pp. 126-133.
- [5] Gupta Rameshwar D and Kundu Debasis (2010). Generalized logistic distributions. *Journal of Applied Statistical Science*, Vol. 18 (1), pp. 51.
- [6] Mudholkar Govind S, Srivastava Deo Kumar and Freimer Marshall (1995). The exponentiated Weibull family: A reanalysis of the bus-motor-failure data. *Technometrics, Taylor & Francis*, Vol. 37 (4), pp. 436-445.
- [7] Gupta Rameshwar D and Kundu Debasis (2007). Generalized exponential distribution: Existing results and some recent developments. *Journal of Statistical planning and inference, Elsevier*, Vol. 137 (11), pp. 3537-3547.
- [8] Oluyede Broderick, Chipepa Fastel and Wanduku Divine (2020). The Exponentiated Half Logistic-Power Generalized Weibull-G Family of Distributions: Model, Properties and Applications. *Eurasian Bulletin of Mathematics*, Vol. 3 (3), pp. 134-161.
- [9] Burr and Irving W (1942). Cumulative frequency functions. *The Annals of mathematical statistics*, Vol. 13 (2), pp. 215-232.
- [10] Eugene Nicholas, Lee Carl and Famoye Felix (2002). Cumulative frequency functions. *Communications in Statistics-Theory and methods, Taylor & Francis*, Vol. 31 (4), pp. 497-512.
- [11] Cordeiro Gauss M, Alizadeh Morad and Ortega Edwin MM(2014). The Exponentiated Half-Logistic Family of Distributions: Properties and Applications. *Journal of Probability and Statistics, Wiley online library*, Vol. 2014 (1), pp. 864396.
- [12] Lagos Alvarez Bernardo, Jerez-Lillo Nixon, Navarrete Jean P and Figueroa-Zu Niga Jorge and Leiva, Victor (2022). A type I generalized logistic distribution: Solving its estimation problems with a bayesian approach and numerical applications based on simulated and engineering data. *Symmetry*, Vol. 14 (4), pp. 655.

- [13] Almalki Saad J and Yuan Jingsong (2013). A new modified Weibull distribution. *Reliability Engineering & System Safety, Elsevier*, Vol. 111, pp. 164-170.
- [14] Agu FI and Onwukwe CE (2019). Modified Laplace distribution, its statistical properties and applications. *Asian Journal of Probability and Statistics*, Vol. 4 (1), pp. 1-14.
- [15] Johnson Norman L, Kotz Samuel and Balakrishnan Narayanaswamy (1995). Continuous univariate distributions. *John wiley & sons*, Vol. 2.
- [16] Omukami Howard A (2022). Generalizations of Logistic Distribution. *University of Nairobi*.
- [17] Willmot Gord (1986). Mixed compound Poisson distributions. *ASTIN Bulletin: The Journal of the IAA, Cambridge University Press*, Vol. 16 (S1), pp. S59-S79.