

Impact of Small Scale Mechanical Rice Transplanters on the Yield, Economics of Rice System in North Coastal Zone of Andhra Pradesh, India

ABSTRACT

Rice is a principal food crop for about half of the population in the world. In India the rice cultivation area has declined due to many reasons labor scarcity, climate change, inapt market price, increased cost of cultivation, yield stabilization. At present, to meet the increased demand, to the growing population it is apparent to increase the area under rice crop with low manpower is needed. In wet transplanting, Mechanised System Rice Intensification (MSRI) planting technology was demonstrated by the Krishi Vigyan Kendra, Amadalavalasa to popularize in the farmer fields. As part of the demonstrations, the production and economic parameters in Paddy transplanted with MSRI and manual planting methods were compared for two years from 2022 and 2023 during *kharif*. The results revealed that, the higher number of productive tillers 16 & 20 per hill and yield 6000 & 6975 kg ha⁻¹ were recorded with the MSRI planting when compared to the Manual planting method 11 & 13 productive tillers per hill and 5600 kg & 6252 kg ha⁻¹ respectively in both the years of the study. An economic point of view in MSRI planting recorded lower cost of cultivation (Rs. 52570 & Rs. 55875 ha⁻¹), higher net returns (Rs. 69500 & Rs. 92981 ha⁻¹) and B:C ratio (2.3:1 & 2.66:1) respectively, compared to the manual planting in both the years.

Key Words: MSRI planting, rice, labour scarcity, Manual Planting, Yield, Economics.

INTRODUCTION

Rice (*Oryza sativa* L.) is the most important staple food for more than half of the world's including populated regions as well the countries with rapid growth. India has the largest area among rice-growing countries and stands second in production with an area of 43.90 million hectares producing 114.45 million tons and productivity of 2.6 tones ha⁻¹. In Andhra Pradesh, an area of 2.25 million hectares is under rice cultivation with a production of 7.79 million tons and productivity of 3.4 t ha⁻¹ (Agricultural Statistics at a Glance 2022, Ministry of Agriculture & Farmers Welfare, Govt of India). Andhra Pradesh contributes to 4.84% of the total area and 5.98 % to total rice production in India. In Srikakulam district, rice cultivated in 1.6 lakh ha with an average productivity of 5.4 t ha⁻¹, the transplanting of rice crop requires nearly 25% of the additional labor for irrigated rice production (Awan *et. al.*, 2008)

Though rice is an important food crop to the world's population, for several reasons area under paddy cultivation decreasing at an increasing rate. The timely availability of manpower itself a major hurdle to the farmer. High cost of cultivation, conversion of cultivated lands into non-agricultural lands and converting for commercial purposes, climatic changes is some of the reasons for decreased rice area (Srinivas *et.al.*, 2020). In order to address these issues Krishi Vigya Kendra, Amadalavalasa has taken up demonstration on Mechanized System of Rice Cultivation (MSRI) in paddy cultivation.

Material and Methods:

Krishi Vigyan Kendra Amadalavalasa conducted Front Line Demonstrations (FLD) on Paddy mechanization with MSRI was conducted for two years (2022, 2023) during *kharif* in the farmers' fields in 5 clusters in 10 locations across the Srikakulam district. The soils of the selected fields were clay loam in texture, low in available nitrogen, medium in phosphorus, potassium and low in organic carbon.

Treatment Details:

T1: Paddy transplanting with MSRI Planter

T2 : Manual transplanting.

Each treatment was imposed in 0.4 ha in the farmers' fields and the yield attributes, yield and economics of each treatment.

The variety used for this demonstration was MTU 1061 (Indra) which is a long duration (155 days), non-lodging, suitable for over-aged seedling planting, tolerant to BPH, with stands under inundated conditions, performs well even slightly saline soils.

Nursery Raising:

For MSRI planting nursery raising is the important management aspect. The following precautions have been taken for successful nursery raising

- Nursery was raised in the trays which have the dimensions of 60cm x 30 cm x 2.5 cm.
- Each tray filled with 2-2.5 kg of finely pulverized soil as the first layer after that 120 g of overnight soaked seed placed with the help of a manual seeding machine. For one hectare 200 trays were filled, and utilized 30-35 kg of seed.
- The seeding tray with the seed was moistened with sparkling water
- Finally, 1.0 to 1.5 kg fine soil along with vermicompost was added as the top layer of the tray, to cover the seed as well. The trays are allowed to stake overnight in the shade. Next day, trays are shifted to the field and spread on the plain laser leveled field covered with paddy straw for three days.
- At seedling emergence, covered paddy straw removed from the trays and start regular watering.
- To support the seedling growth, 19:19:19 @ 3-5 g per lit was sprayed.
- 18 days old seedlings from nursery were used to transplanting with MSRI Planter.
- The entire process tray filling was done with the help of an automatic seeding machine (please add company and make) with the support of skilled staff.

MSRI Planter:

Four row and six row MSRI planters (Kubota Manufacturer company) of Krishi Vigyan Kendra Amadalavalasa were used by the farmers for wet transplanting of rice. The planter is with 18-22 HP capacity, operated by petrol. These planters can plant 2.4 hectares per man-day. The planter can plant 4 seedlings per hill with a spacing of 30 cm X 14-18 cm.

Main field Preparation:

Main fields used for demonstration were properly puddled and leveled. Initial puddling was done 5-6 days before planting and the second puddling was performed one day before planting.

Weed management:

As part of the weed management protocol, immediately after planting pre-emergence application of Pretilachlor @ 2 ml/lit in 500 lit water per ha was recommended. At 15 DAT, Bispyribac Sodium @ 250 ml/ha and Metsulfuron Methyl+

Chlorimuron Ethyl 20 g/ha were recommended as tank mix for control of Grasses, Sedges, and BLW control.

Nutrient Management:

All the demonstrated fields received uniform dose of nutrients as per the recommendation (80:60:50 kg NPKha⁻¹) for the North Coastal Andhra Pradesh by Acharya N.G. Ranga Agricultural University

Data Collection:

The observations were recorded from 5 randomly selected plants in each field. plant population per m⁻² was taken in the field randomly with the help of quadrant. Yield attributes viz., No. of productive tillers plant⁻¹ and m⁻² were also taken.

Grain yield was taken with help of a Crop Cutting (CC) experiment. In the CC experiment 5 m x 5 m plot was marked in the field and the crop was harvested, threshing was completed and grain yield was noted in kgs and converted into kgha⁻¹.

Results and Discussions:

The data on plant population m⁻², No of Productive tillers per hill and yield of rice depicted in Table 1. revealed that, in the manual planting method (without proper spacing), no of productive tillers per hill was less (11 and 13 respectively in 2022 and 2023) it was attributed due to high intra competition within the hill among the seedlings, deep placement of the seedlings in the muddy fields (Rao *et al.*, 2020). In manual planting, generally, the workers kept more number of seedlings (6 or more) per hill, also inadequate plant stand per m⁻² due to the deep placement of seedlings, tillers producing node of the rice plant kept very deep (>4-5 cm) in the mud reduces the tillering ability of the plant which in turn reduce the productive tillers too. While in the MSRI planting, no of seedlings per hill we can adjust as per our requirement. In machine transplanting it was observed 4 seedlings per hill were kept at uniform inter and intra row spacing (30 cm X 14 cm) and shallow depth placement of rice seedlings resulted more number of tillers per hill contributed increase productive tillers 16 and 20 respectively in both the years of the study. Pasha *et al.*, (2012) reported that, better tillering, number of loaded grains panicle⁻¹, and more panicle hills, with yangi eight-row transplanter than conventional transplanting in rice. Singh *et al.*, (2005) reported, quick establishment, and consistent development of rice seedlings was due to the consistent placement of seedlings at a particular depth and spacing with a comparable number of seedlings per hill under one square meter. Plant height, total panicles m⁻², effective panicles m⁻², grains per panicle, and 1000 grain weight were noticed in the self-propelled rice transplanter which in turn produces the highest grain yields compared to manual transplanting and direct seeding. Mechanical transplanting increased number of panicles hill⁻¹ and viable grains per panicle was reported by Sheeja *et al.*, (2012). Rao *et al.*, (2020) revealed that the

highest no of productive tillers m^{-2} (254 and 242) and grains per panicle (211 and 188) were noticed in MSRI planting when compared to the manual planting (203,206, 188 and 189) in 2018 and 2019 during *kharif* respectively.

Uniform wider spacing in the MSRI planting method inturn improves the solar radiation interception in all directions of the rice plant and CO_2 replacement may happen in the plant microclimate, which could be attributed to increased photosynthetic rate, photosynthate translocation also.This will increase number of filled grains per panicle which turn result in anincreased yield in the MSRI planting (6000 and 6975 $kg\ ha^{-1}$) when compared to manual planting (5600 and 6252 $kg\ ha^{-1}$). These findings are in conformity with Haung *et al.*, (2020). He identified early sowing can improved biomass translocation, and grain yield in machine-transplanted late-season rice under single-seedling sowing than conventional seeding.Similar results were reported with Singh *et.al.*, 2006. Mechanical transplanting significantly increased grain yield by about 23, 37 and 63 %, straw yield by about 17,14 and 22 % and biological yield by about 20, 24 and 39 % over manual transplanting, dry direct seeding and direct seeding of sprouted rice in puddled conditions, respectively.Rao *et.al.*, (2020) recordedthe higher grain yield in the demo plot (6975 $kg\ ha^{-1}$ & 6537 $kg\ ha^{-1}$) which was planted with the MSRI planter than in manual planting (5860 $kg\ ha^{-1}$ & 5337 $kg\ ha^{-1}$) in two consecutive years.

In the manual method of planting aged seedlings,(30 days or even more aged seedlings) were used to be planted in the main field but in MSRI planting, young aged (18 days) seedlings are used because of age advantage, theseyounger seedlings have more vigour potential to put forth high tillering ability in the main field.Mechanically transplanted rice had the most effective tillers m^{-2} because younger seedlings were transplanted at the proper distance and depth, minimizing transplanting tremors and allowing early establishment and growth factor exploitation (Manjunath *et al.* 2009). Srinivas *et.al.*,(2020) revealed in their study that, with the young seedlings (12 -14days), the minimum (two-three) seedlings per hill and maximum (20 cm) intra-row space adjustment to sufficient number matrix of hills per square meter is adjusted with the MSRI planter which can contribute to higher yields. Revathi *et al.* (2016) observed higher mean grain, straw yield and B:C ratio (average of two years) in machine transplanting (7.0, 16.02 $t\ ha^{-1}$ & 2.20) followed by farmers method (manual) of planting (6.30, 13.32 $t\ ha^{-1}$ & 2.00) and was found on par to SRI method (5.8,11.40 $t\ ha^{-1}$ & 1.60). The higher number of panicles m^{-2} (317), grains per panicle (277), grain yield (5345 $kg\ ha^{-1}$) and straw yield (6305 $kg\ ha^{-1}$) were recorded in machine transplanting over conventional, broadcasting and drum seeding establishment methods (Ramulu *et al.*, 2019) and (Srinivas *et.al.*,2020). According to Sreenivasulu and Reddy 2014,Mechanical transplanting improved productive tiller, panicle length, panicle⁻¹ grains, and rice test

weight over hand transplantation. Mechanically transplanted rice yielded 9-14% more than hand-transplanted rice was identified by Xing *et al.*, 2017. Mechanically transplanted paddy yielded 7.42 t ha⁻¹, while broadcasted paddy yielded 5.31 reported by Regmi *et al.*, 2020.

Table.1: Yield attributes and yield of paddy influenced by normal and MSRI planting methods.

Treatment	Plant population m ⁻²		No. of Productive tillers/hill		Yield kg ha ⁻¹	
	2022	2023	2022	2023	2022	2023
T1: Paddy transplanting with MSRI Planter	24	24	16	20	6000	6975
T2: Manual transplanting	28	27	11	13	5600	6252
% Increase			45.4	53.8	7.1	11.5

Economic parameters:

The economic parameters of both systems were depicted in Table 2 and compared in Table 3. The results revealed that, in normal planting cost of cultivation was high due to 1. Seed cost, 2. High labour involvement, 3. Nursery pulling and planting, 4. More pest and disease incidence and lower yield. In case of the MSRI planting savings on expenditure noticed on seed, labour and pesticides. In addition an increased yield also noticed in MSRI Planting. Higher cost of cultivation (Rs. 57850 and Rs. 66375 ha⁻¹) was recorded in manual planting in both the years respectively. In MSRI planting highest net returns (Rs. 69500 and Rs. 92981 ha⁻¹) were noticed in both the years. A similar trend was noticed in B: C ratio also. These findings are similar with the other researchers of Srinivas *et al.*, 2020 and Sreenivasulu and Reddy, 2014., On-farm trials indicated that the yield parameters and yield were higher in mechanized transplanting than in manual transplanting. Mechanized transplanting recorded more grain yield (6359 kg ha⁻¹) and net returns (Rs. 31870/- ha⁻¹) with less cost of cultivation (Rs. 29796/- ha⁻¹) compared to manual transplanting.

Table.2: Yield, Economic parameters of Paddy in Manual and MSRI planting methods.

Treatment	Yield (Kg ha ⁻¹)		Cost of Cultivation (Rs.per ha)		Net returns (Rs.per ha)		B:C ratio	
	2022	2023	2022	2023	2022	2023	2022	2023
T1: Paddy transplanting with MSRI Planter	6000	6975	52570	55875	69500	92981	2.3:1	2.66:1
T2: Manual transplanting	5600	6252	57850	66375	56250	66375	1.9:1	2:1
% Increase	7.1	11.5						

Table.3: Minimum Expenditure per ha on paddy under Manual and MSRI planting conditions

S.No.	Particulars	Manual Method (Rs)	MSRI (Rs)
1	Seed	3000	1500
2	Nursery raising (including ploughing)	2250	-
3	Field preparation	3750	3750
3	Trays filling @ 20 per tray (200 trays)	-	4000
4.	19:19:19 spray	-	300
5.	Nursery pulling and spreading of nursery	10000	-
6.	Transplanting	7500	6250
7	Fertilizers	6425	6425
8	Formation of Alley ways and one hand	3500	3000

	weeding		
8	weedicides	7450	5500
9	Pesticides	8500	5300
10	Harvesting	8750	8750
11	Labour charges for fertilizers and irrigation	4600	4600
	Total	65725	49375




Fig 1 : Nursery in Trays initial stage and ready to transplanting



fig 2 : Planting of the paddy with MSRI Planter



Fig 3 : After planting with MSRI planter



Fig 4 : Taking of Biometric Observations

References:

Awan, T.H.; Ali, I.; Safdar, M.E.; Ahmad, M.; Akhtar, M.S. Comparison of parachute, line, and traditional rice transplanting methods at farmer's field in rice growing area. *Pak. J. Agric. Sci.* 2008, 45, 432–438.

Agricultural Statistics at a Glance 2022, Ministry of Agriculture & Farmers Welfare, Govt of India.

Manjunatha M V, Reddy B G M, Shashidhar S D, Joshi V R. 2009. Studies on the performance of self-propelled rice transplanter and its effect on crop yield. *Karnataka Journal of Agricultural Sciences*. 22(2):385–387.

Min Huang, *, Shengliang Fang, Fangbo Cao, Jiana Chen, Shuangli Shana, Yu Liua, Tao Leia, Alin Tiana, Zui Taoa, Yingbin Zou. 2020. Early sowing increases grain yield of machine-transplanted late-season rice under single-seed sowing. *Field crops research*. 107832

Pasha L, Bhadrud, Krishna L, Naik R B M. 2012. Comparative performance of different rice planting methods under Nagarjuna sagar project left canal command area of nalgonda district. *Crop Research*. 44(1-2): 1–4.

Ramulu, C.H., Raghu Rami Reddy, P and Narsaiah, E. 2019. Effect of different establishment methods and nitrogen levels on yield attributes, yield, nutrient uptake and economics of rice. *Journal of Research PJTSAU*, Vol. 47, No. 3, 27-32 ref. 15 ref.

Regmi R C, Kharel R, Regmi R. 2020. Effect of planting methods on yield and yield components of spring rice in Bardiya, Nepal. *Acta Scientifica Malaysia (ASM)*, Zibeline International Publishing. 4(2): 61- 63. <http://doi.org/10.26480/asm.02.2020.61.63>

Revathi, P., Devi, K. B. S., Reddy, B. G., Rao, V. P., Padmaja, G and Sanker, A.S. 2016. Influence of planting methods and integrated nutrient management practices on yield attributes, yield and economics of rice. *Journal of Research PJTSAU*, Vol. 44, No. 4, 25-32 ref. 12

Sheeja K R, Mathew R, Jose N, Leenakumary S. 2012. Enhancing the productivity and profitability in rice cultivation by planting methods. *Madras Agriculture Journal*: 99 (10-12): 759-761

Singh R P, Singh C M and Singh A K. 2005. Effect of crop establishment methods, weed management and split nitrogen application on weeds and yield of rice. *Ind. J Agri Sci.* 75: 285-287.

Singh KK, Lohan SK, Jat AS and Rani T (2006). New technologies for planting rice for higher production. *Research on crops* 7(2): 369-371.

Srinivasa Rao, M.M.V., Roy, G. S. and Lakshmana, K. (2020). Mechanized system rice intensification (MSRI) is boon to farmers to save money and time in rice cultivation in Vizianagaram district of North Coastal Zone of Andhra Pradesh. *Agriculture update*: 15 (3): 162-166.

Sreenivasulu, S and Reddy, P. B. H. 2014. Effect of mechanized transplanting on yield, yield attributes and economics of rice (*Oryza sativa*). *Journal of Research ANGRAU*, Vol. 42, No. 2, 9-12 ref. 8

Rao, K T., Pradeep Kumar, P. B and Chandrayudu, E. 2020. Mechanized system rice intensification (MSRI) in rice cultivation at Visakhapatnam district of Andhra Pradesh, *International journal of plant Sciences* 15 (2): 135-138.

Xing Zhi-peng, Wu Pei, Zhu Ming, Qian Hai-jun, Hu Ya-jie, Guo Bao-wei, Wei Hai-yan, Xu Ke, Huo Zhong-yang, Dai Qi-gen, Zhang Hong-Cheng. 2017 Temperature and solar radiation utilization of rice for yield formation with different mechanized planting methods in the lower reaches of the Yangtze River, China. *Journal of Integrative Agriculture*. 16(9): 1923–1935.

UNDER PEER REVIEW

UNDER PEER REVIEW