

Response of chickpea (*Cicer arietinum* L.) to phosphate solubilizing microorganisms and phosphorus levels on nutrient availability, uptake and seed yield in inceptisol.

Commented [A1]: Kindly revise your title. Make it clear and it should portray your work

Abstract

The present Investigation was carried out for the “Response of chickpea (*Cicer arietinum* L.) to phosphate solubilizing microorganisms and phosphorus levels on -nutrient availability, uptake and seed yield in inceptisol.” The experiment was conducted in factorial randomized block design (FRBD) with three replications .during Rabi 2019-20 at department research farm of SSAC, College of Agriculture, Latur. The treatments comprises four main (absolute control, *Bacillus megaterium*, *Aspergillus niger* and *Aspergillus awamori* @ 10 ml kg⁻¹ seed treatment) and four sub treatments (0,45,60 and 75 P₂O₅ kg ha⁻¹) The results indicates that, the incorporation of phosphate solubilizing microorganisms viz.*Aspergillus awamori* @ 10 ml kg⁻¹ seed in combination with soil application of 75 P₂O₅ kg ha⁻¹ found to be significantly higher availability of nutrient, increased total N, P and K uptake of chickpea as compared to *Aspergillus niger* and *Bacillus megaterium* along with 60 P₂O₅ kg ha⁻¹ and over control .Further results revealed that seed yield was significantly improved with the seed treatment of *Aspergillus awamori* @ 10 ml kg⁻¹ seed in combination with application of 75 P₂O₅ kg ha⁻¹.

Commented [A2]: No need to state this one, rather, please make short introductory statements covering the background of your study and stating your objectives. Also, please be mindful of your spacing and grammar. Consider rephrasing the rest of the abstract.

Formatted: Highlight

Formatted: Highlight

Keywords: chickpea, microorganisms, nutrient, phosphorus, uptake, yield

1. Introduction

Pulses, edible seeds from the legume family, are an important group of food crops that can play an important role to address national food and nutritional security and also tackle environmental challenges. Among the pulses, India is one of the major chickpea producing country in the world and ranked first in area and production with 34 per-cent and 26 per-cent respectively, as compare to total world production. Chickpea is the rabi pulse crop grown in country supplementing protein (17-25 per cent), amino acid, Vit. A, Vit. C, Vit. B, Vit. K, source of folic acid demand of vegetarian diet. Chick pea is not only important human diet but also it plays a

Commented [A3]: Please consider your reader as others are not also familiar with rabi. In writing your paper kindly consider a wide range of readers as they may not be familiar with.

Commented [A4]: You mean in your country? Please be very clear with your phrases or sentences.

significant role in improving soil fertility by fixing the atmospheric nitrogen. Chickpea meet about 80 per cent of nitrogen requirement from symbiotic nitrogen fixation from air. It leaves substantial amount of residual nitrogen for subsequent crops and add plenty of organic matter to maintain and improve soil health and fertility. (Prajapati *et al.*, 2017) ^[18].

Formatted: Highlight

Formatted: Highlight

Phosphorus can be termed as 'life mineral' because of its crucial role in metabolic and energy transfer reactions in plant. Phosphorus is an essential element in DNA and RNA that contain the genetic code of the plant to produce protein and other compounds essential for plant structure, seed yield and genetic transfer. It is also associated with increased root growth, chlorophyll content, straw strength and crop maturity in cereals and N₂-fixation in legumes. Thus, phosphorus is essential for vigorous growth and development of reproductive parts in crops. The phosphorus deficiency leads to stunted root and shoot growth, bluish green coloration of leaf, delayed maturity and poor grain development in cereals. Thus, phosphorus has become a major constrain in agricultural production mainly because of its fixation in soils involving both adsorption and precipitation reactions. The rate and magnitude of phosphate adsorption depends upon the properties of soils and phosphorus resources Barros *et al.*, 2005, Boparai^[6]. Use of phosphorous in soils phosphorous solubilizing microorganisms (PSMs) are capable to convert insoluble phosphorous to soluble forms can function as biofertilizers to increase the native phosphorous in soil (Narsian and Patel, 2000) ^[14]. Low fertility, particularly phosphorus deficiency is one of the major constraints to increase the chickpea productivity (Srinivasrao *et al.*, 2003)^[21]. Use of biofertilizers is low-cost renewable source of plant nutrients, which supplement chemical fertilizer. Biofertilizers solubilize plant nutrients like N and P through their activities in the soil or rhizosphere and make available to plant in gradual manner. PSB solubilize insoluble phosphorus compounds by exerting organic acids, which is the primary mechanism of solubility of insoluble inorganic phosphates. Besides organic acids, production of chelating substances, mineral acids and proton extrusion also involved (Rooge, *et al.*, 1998) ^[19]. Phosphate solubilizing microorganisms, such as *Aspergillus* and *Bacillus*, play a crucial role in solubilizing soil phosphorus through mechanisms like pH reduction and organic acid secretion Bharadkar *et al.*, (2021). ^[4] Mittal *et al.*, (2008) ^[12] found that seed inoculation of chickpea with *Aspergillus awamori* increased shoot height by 7-12 per-cent, a nearly 3 fold increase in seed

Formatted: Highlight

Formatted: Highlight

Commented [A5]: Kindly cite your reference.

Commented [A6]: Vague statement. Please consider rephrasing.

Commented [A7]: Kindly use punctuation.

weight as compared to uninoculated control. Seed inoculation with *Aspergillus awamori* increased the growth total P content and biomass of mungbean (Jain *et al.*, 2012) ^[08].

2. Materials and methods

The field experiment was conducted during *Rabi* 2019-20 at research farm departmental farm of SSAC, College of Agriculture, Latur using chickpea crop (Var.BDNG-797). In order to evaluate the response of chickpea (*Cicer arietinum* L.) to phosphate solubilizing microorganisms and phosphorus levels on nutrient availability, uptake and seed yield in inceptisol. The experimental soil pH was found to be 7.8, Electrical conductivity (0.28 dSm^{-1}), organic carbon 4.6 g per kg, and CaCO_3 5.6 per cent. Available N, P_2O_5 and K_2O were 143.25, 6.28 and $317.19 \text{ kg ha}^{-1}$, respectively. The experimental soil was clayey in texture, moderately alkaline in reaction, low in available nitrogen and phosphorus and high in available potassium. Soil pH was determined by 1:2.5 soil water suspension ratio using a digital pH meter (Jackson, 1973) ^[07]. Electrical conductivity of soil was determined by 1:2.5 soil: water suspension ratio using the Conductivity Bridge (Jackson, 1973) ^[07]. Soil organic carbon was determined by modified method of Walkley-Black methods (1934) ^[26]. The calcium carbonate content in soil was determined by Rapid titration method (Jackson, 1973) ^[07]. Available nitrogen was determined by alkaline potassium permanganate method (Subbiah and Asija, 1967). Available phosphorus was extracted from the soil with 0.5 M Sodium bicarbonate by Olsen's method (Jackson, 1973) ^[07]. Available potassium was determined with neutral normal ammonium acetate and potassium in the extract was determined on Flame Photometer (Piper, 1966) ^[66].

After completion of preparatory tillage operations, the experiment was laid out in factorial randomized block design (FRBD) with sixteen treatments replicated thrice. Organic manures i.e. FYM was applied at the rate of 5 t ha^{-1} prior to 15 days of sowing of chickpea crop and all the plots were fertilized with recommended dose of fertilizer NPK (25:50:00 kg ha^{-1}) was applied as a basal dose through urea, SSP treatment wise at the time of sowing. The treatments comprises as a seed treatment T_0 : Control, T_1 : *Bacillus megaterium* @ 10 ml kg^{-1} seed, T_2 : *Aspergillus niger* @ 10 ml kg^{-1} seed, T_3 : *Aspergillus awamori* @ 10 ml kg^{-1} seed as a main treatments and application P_0 : 0 P kg ha^{-1} , P_1 :45P kg ha^{-1} , P_2 :60P kg ha^{-1} , P_3 :75P

Commented [A8]: Please clearly state the objectives of your study. Kindly, elaborate also the impact of your research.

Commented [A9]: Is this a result? Please state materials and methods in way of narrating or describing your steps or protocols. How did you conduct your experiment? Also, observe sectioning of your materials and methods to set things in order. Like, A. Preparation and Analysis of Experimental Soil, B. The Experimental Set-up, so on and so forth..

kg ha⁻¹ as a sub main treatments. Seed was inoculated with *Aspergillus awamori*, *Bacillus megaterium* *Aspergillus niger* @ 10 ml / kg seed.

Chickpea seed was sown on 09 October 2019 by dibbling method as per randomly replicated plot having size 3× 2 m² maintained row to row spacing 30 cm and plant to plant 10 cm and using a seed rate of 80 kg ha⁻¹. The crop was harvested at maturity stage on 22 January 2020. The observation recorded the seed yield, fodder yield were recorded at harvest stage. The data collected from the above observation were analysed statistically by the procedure prescribed by Panse and Sukhatme (1967)^[15]

For the determination of nutrient contents in plant samples fresh plant samples and grain sample were collected at harvest stage from each net plot and processed with following standard procedure of washing, sun drying and then oven drying and grinding. The grind plant materials were stored in the paper bags and used for further chemical analysis. One gram of fine powered plant and grain sample, 15 ml of tri-acid mixture (HNO₃:H₂SO₄ HClO₄ in a ratio of 10:1:4) was added and digested plant and grain sample as described by Piper (1966)^[17]. It was kept in digestion chamber till complete digestion of the sample. The residue was dissolved in double-distilled water and after filtration; final volume was made to 50 ml. Nutrient content in plant and grain sample were analyzed for the total nitrogen content determined by Micro-kjeldahl's method A.O.A.C., (1965)^[01], total phosphorus vanadomolybdo phosphoric acid yellow colour method spectrophotometrically as described by (Jackson, 1973), total potash estimated by di-acid extract (Piper 1966)^[17] by using flame photometer. Nutrient uptake (kg ha⁻¹) by using the following formula - {Nutrient uptake (kg ha⁻¹) = Nutrient content (%) X yield (kg ha⁻¹) / 100}.

Results and discussion

3.1 Effect of phosphate solubilizing microorganisms and phosphorus levels on available nutrients.

3.1.1 Available nitrogen:

The available nitrogen was influenced significantly due to seed inoculation of phosphate solubilizing microorganism (Table 1). The data indicated that the maximum available of N content in soil was found with seed inoculation of *Aspergillus awamori* (166.68 kg ha⁻¹) followed by *Aspergillus niger* (165.81 kg ha⁻¹) and *Bacillus megaterium* (158.44 kg ha⁻¹). The lower value of (154.89 N kg ha⁻¹)

Commented [A10]: Please clearly state your experimental set-up. Observe proper punctuation and spacing to avoid confusing sentences.

Commented [A11]: What is your reference for this formula?

Commented [A12]:

Commented [A13]: When you say available nitrogen, it pertains to the N present in the experimental soil? Or present on the plant and seed samples?

available N was recorded in control. The application of different phosphorus levels significantly increased the available N kg ha⁻¹. The highest available N was recorded with the application of phosphorus @ 75 kg P₂O₅ ha⁻¹ (163.52 kg ha⁻¹). As the levels of phosphorus decreases the available N content was also decreases. While minimum available N (158.01 kg ha⁻¹) was recorded in control. The data showed that the interaction effect of phosphate solubilizing microorganism and phosphorus levels on available nitrogen showed non-significant. Inoculation of phosphate solubilizing microorganism and application of phosphorus levels resulted in reduction in denitrification and leaching losses of N and that result in improve the soil health. Increased available nitrogen observed due to phosphorus play an important role in nodule initiation and root proliferation. Nodule formation increases the nitrogen fixation and ultimately increases soil available nitrogen (Jahangir et al., 2017) ^[09]. Similar results were reported by Kumar et al., (2019) ^[10] studied that the application of 60 kg P₂O₅ and inoculation of biofertilizers significantly increased available N (230.5 kg ha⁻¹) and (224.7 kg ha⁻¹) respectively in the soil. N status increased with increase in the levels of P and biofertilizers (PSB and AM). This might be attributed to the application of P and biofertilizers which enhanced and established better root system. Nutrients possibly stimulate the modulating bacteria for more fixation of atmospheric N₂ resulting in increase of its contents in the soil over control.

Table 1: Available N as influenced by phosphate solubilizing microorganism and phosphorus levels in chickpea.

PSM	Phosphorus levels (P ₂ O ₅ kg ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Available Nitrogen (kg ha ⁻¹)				
T0	149.73	155.86	156.96	157.02	154.89
T1	155.15	158.88	159.13	160.61	158.44
T2	163.12	165.84	166.60	167.66	165.81
T3	164.05	166.23	167.63	168.79	166.68
Mean	158.01	161.70	162.58	163.52	
	T		P		TXP
SE	0.29		0.29		0.58
C.D at 5%	0.83		0.83		NS

Commented [A14]: Are your set up replicated? I can see that it is a two factor analysis, how did you perform statistical analysis of your data? Please elaborate that on your methodology. In the table, no need to present raw data, results should be analyzed and presented as mean values ± standard deviation.

3.1.2 Available phosphorus

The data on effect of phosphate solubilizing microorganisms on available phosphorus in chickpea after harvest was reported in (table 2). The data indicated that seed inoculation with phosphate solubilizing microorganisms influenced significantly.

Commented [A15]:

Among the phosphate solubilizing microorganisms seed inoculation with *Aspergillus awamori* recorded higher value of available phosphorus (9.68 kg ha⁻¹) followed by *Aspergillus niger* (9.25 kg ha⁻¹) and *Bacillus megaterium* (8.67 kg ha⁻¹). While minimum available phosphorus (7.89 kg ha⁻¹) was recorded in control after harvest of chickpea.

The application of different phosphorus levels significantly increased the available phosphorus in soil. The maximum available phosphorus was recorded with application of phosphorus @ 75 kg P₂O₅ ha⁻¹ (9.42 kg ha⁻¹) followed by application of phosphorus @ 60 kg P₂O₅ ha⁻¹ (9.12 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (8.73 kg ha⁻¹). The lowest value of available phosphorus was recorded in control after harvest of crop. Thus, it was observed that the available phosphorus in soil was decreased significantly as decremented in the levels of phosphorus.

Further data shows that the interaction effect of phosphate solubilizing microorganisms and phosphorus levels on available phosphorus found significant. The interaction *Aspergillus awamori* and application of 75 kg P₂O₅ ha⁻¹ found maximum availability than rest of the interactions. The increase in the available phosphorus might be due to the fact that increase in the solubilization of inorganic P by secreting more organic acids in the soil which helps to increase the P availability Yadav *et al.*, (2017)^[24] reported that, the application of *Aspergillus awamori* + *Pseudomonas striata* to mungbean increased P content in soil as compared to control. *Pseudomonas striata* stimulate the microbial activity in soil and after decaying of their bodies in soil increases the P content in soil. Similar findings were also reported by Laharia *et al.*, (2019)^[11].

Table 2: Available P as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea

PSM	Phosphorus levels (kg P ₂ O ₅ ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Available phosphorus (P ₂ O ₅ kg ha ⁻¹)				
T0	6.43	7.50		8.33	7.57
T1	7.83	8.53	8.90	9.40	8.67
T2	8.53	9.23	9.50	9.73	9.25
T3	8.77	9.67	10.03	10.23	9.68
Mean	7.89	8.73	9.12	9.42	
	T		P		TXP
SE	0.18		0.18		0.37
C.D at 5%	0.25		0.25		0.50

3.1.3. Available potassium:

The results obtained in respect to phosphate solubilizing microorganism and phosphorus levels on available potassium after harvest of crop reported in (table 3). The inoculation of microorganism *Aspergillus awamori* recorded maximum available potassium (784.25 kg ha⁻¹), followed by *Aspergillus niger* (283.71 kg ha⁻¹) and *Bacillus megaterium* (280.69 kg ha⁻¹) while minimum available potassium recorded in control (278.96 kg ha⁻¹) after harvest of chickpea. However, available potassium could not reach to level of significance due to use of different phosphate solubilizing microorganisms but increase in potassium status in soil. The effect of application of different phosphorus levels were shows non-significant differences in available K. The maximum available K was recorded with application of phosphorus @ 75 kg P₂O₅ ha⁻¹ (283.35 kg ha⁻¹), followed by 60 kg P₂O₅ ha⁻¹ (295.82 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (280.89 kg ha⁻¹). The minimum available K content was recorded in control after harvest of crop. However increase in potassium status noticed as the levels of phosphorus increased. The interaction effect of phosphate solubilizing microorganism and phosphorus levels on available potassium was found non-significant. The availability of potassium was decreased as compared to initial available K (317.19 kg ha⁻¹) due to the unavailability of K content in fertilizer that resulted in decrease the available K content in soil. Laharia *et al.*, (2019) ^[11] reported that application of 125% RDP + PSB recorded available potassium in soil after harvest of chickpea (381.07 kg ha⁻¹). Available potassium increased in soil with increasing phosphorus level with PSB. Similar findings were also reported by Kumar *et al.*, (2019) ^[10] and Jahangir *et al.*, (2017) ^[09].

Table 3: Available K as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea

PSM	Phosphorus levels (P ₂ O ₅ kg ha ⁻¹)				
	P0	P45	P60	P75	Mean
	Available potassium (K ₂ O kg ha ⁻¹)				
T0	276.44	279.32	281.15	278.15	278.96
T1	281.21	279.94	281.98	279.64	280.69
T2	280.93	287.61	282.50	283.81	283.71
T3	284.96	276.34	284.70	291.00	284.25
Mean	280.89	280.20	282.58	283.35	
	T		P		TXP
SE	1.57		1.57		3.14

C.D at 5%	NS		NS		NS
-----------	----	--	----	--	----

3.2. Effect of phosphates solubilizing microorganism and phosphorus levels on nutrient uptake by chickpea.

3.2.1 N uptake.

The data pertaining to effect of phosphate solubilizing microorganism on N uptake by chickpea was presented in (table 4). The significantly maximum N uptake (58.65 kg ha⁻¹) was recorded with seed inoculation with *Aspergillus awamori* as compare to *Aspergillus niger* (55.81 kg ha⁻¹) and *Bacillus megaterium* (55.21 kg ha⁻¹). While minimum total N uptake was noticed in control. The data on effect of different phosphorus levels on chickpea shows that application of 75 kg P₂O₅ ha⁻¹ recorded significantly higher N uptake (60.90 kg ha⁻¹) against the application of 60 kg P₂O₅ ha⁻¹ (56.41 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (54.97 kg ha⁻¹). Whereas, minimum N uptake was recorded in control (51.99 kg ha⁻¹).

The combined effect of different phosphate solubilizing microorganism and phosphorus levels shows significant effect on N uptake. The interactive effect of seed inoculation with *Aspergillus awamori* and application of phosphorus @ 75 kg P₂O₅ ha⁻¹ found significant on uptake of N as compare to rest of interactions. This may be attributed to synergistic effect of microorganisms and phosphorus levels on increase in the availability of N in soil which helps to enhance vegetative growth of plants thereby increase in N uptake by chickpea. Similarly increase in N uptake owing to significant increase in seed and straw yield. Laharia *et al.*, (2019) ^[11] stated that N uptake by chickpea was significantly influenced with increasing levels of P along with PSB over control.

Table 4: N uptake as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea

PSM	Phosphorus levels (kg P ₂ O ₅ ha ⁻¹)				
	P0	P45	P60	P75	Mean
	Total N uptake (kg ha⁻¹)				
T0	50.10	52.73	54.00	58.28	53.77
T1	51.72	54.42	55.32	58.55	55.21
T2	51.99	55.57	56.13	59.55	55.81
T3	54.17	57.15	59.32	64.00	58.65
Mean	51.99	54.97	56.41	60.90	
	T		P		TXP
SE	0.21		0.21		0.42
C.D at 5%	0.62		0.62		1.24

Formatted: Highlight

3.2.2 P Uptake

The data pertaining to effect of phosphate solubilizing microorganism on P uptake by chickpea presented in (table 5). The significantly maximum P uptake (26.23 kg ha⁻¹) was recorded with seed inoculation with *Aspergillus awamori* as compare to *Aspergillus niger* (23.17 kg ha⁻¹) and *Bacillus megaterium* (18.86 kg ha⁻¹) while low total P uptake was found in control. Application of different phosphorus levels influenced significantly on P uptake by chickpea. Among the different P levels application of phosphorus @75 kg P₂O₅ ha⁻¹ recorded significantly higher P uptake (22.21 kg ha⁻¹) as compare to application of phosphorus @ 60 kg P₂O₅ ha⁻¹ (21.06 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (19.87 kg ha⁻¹). Whereas minimum value of P uptake was recorded in control (19.23 kg ha⁻¹).

The combined effect of different phosphate solubilizing microorganism and phosphorus levels shows significant effect on P uptake. The effect of seed inoculation of *Aspergillus awamori* and application of phosphorus @ 75 kg P₂O₅ ha⁻¹ found significant on P uptake. This relates to synergistic effect between P and phosphate solubilizing microorganisms thereby higher biomass production as well due to more plant accessible by phosphate solubilizing microorganism from native and applied phosphorus. (Vidhyashree *et al.*, 2017) [25] reported that *Aspergillus awamori* and *Aspergillus niger* increase the growth parameters of plant due to higher phosphatase activity in the rhizosphere and production of organic acid might have solubilized the insoluble and native phosphate and brought into soluble form. This type of results was also reported by Paratey and Wani (2005) [16].

Table 5: P uptake as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea.

PSM	Phosphorus levels (P ₂ O ₅ kg ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Total P uptake (kg ha ⁻¹)				
T0	12.35	13.44	14.30	16.35	14.11
T1	17.43	17.99	19.50	20.53	18.86
T2	21.81	22.40	23.79	24.70	23.17
T3	25.32	25.68	26.67	27.28	26.23
Mean	19.23	19.88	21.06	22.21	
	T		P		TXP
SE	0.12		0.12		0.24
C.D at 5%	0.34		0.34		0.69

3.2.3 K uptake

The data presented in table 6 indicated that the total K uptake of chickpea increased due to application of different phosphate solubilizing microorganisms. The maximum K uptake (60.24 kg ha⁻¹) was recorded with seed inoculation with *Aspergillus awamori* as compare to *Aspergillus niger* (59.20 kg ha⁻¹) and *Bacillus megaterium* (55.94 kg ha⁻¹) while low total K uptake was found in control. It is found that the application of different phosphate solubilizing microorganisms not reached upto the mark of significance but recorded maximum value of K uptake as compared to control. The data revealed that among the different P levels application of phosphorus @ 75 kg P₂O₅ ha⁻¹ recorded higher K uptake (59.63 kg ha⁻¹) as compare to application of phosphorus @ 60 kg P₂O₅ ha⁻¹ (58.63 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (57.96 kg ha⁻¹). While, low uptake of P was recorded in control (57.11 kg ha⁻¹).

The combined effect of different phosphate solubilizing microorganism and phosphorus levels shows non-significant effect on K uptake. The higher value of K uptake due to application of phosphate solubilizing microorganism and phosphorus levels might be due to fact that more microbial activity in the rhizosphere and addition of FYM at the time of sowing of chickpea increase the available K in soil ultimately increase the uptake of nutrients (Laharia *et al.*, 2019) [11]. P favors to better root proliferation in soil which helps to absorb more nutrient from soil and transported to plant there by increase in uptake of K (Singh *et al.*, 2014) [20]. Similar findings also reported by Biswas *et al.*, (2015) [05].

Table 6: K uptake as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea

PSM	Phosphorus levels (P ₂ O ₅ kg ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Total Uptake K (kg ha⁻¹)				
T0	54.37		55.81	57.50	55.94
T1	56.93	58.51	58.03	58.47	57.98
T2	59.68	59.26	58.56	59.33	59.20
T3	57.44	57.96	62.35	63.23	60.24
Mean	57.11	57.96	58.69	59.63	
	T		P		TXP
SE	0.66		0.66		1.32
C.D at 5%	NS		NS		NS

3.3 Effect of phosphate solubilizing microorganisms and phosphorus levels on yield and yield attributes of chickpea

3.3.1 Seed yield:

The seed yield of chickpea as influenced by phosphate solubilizing microorganism narrated in table 7. The seed inoculation with *Aspergillus awamori* produced higher seed yield (1502.68 kg ha⁻¹) as compared to application of *Aspergillus niger* (1406.6 kg ha⁻¹) and *Bacillus megaterium* (1344.53 kg ha⁻¹). Significant improvement in yield was noticed with the application different phosphorus levels. Among the different phosphorus levels application of phosphorus @ 75 kg P₂O₅ ha⁻¹ recorded higher seed yield (1451.0 kg ha⁻¹) as compare to application of phosphorus @ 60 kg P₂O₅ ha⁻¹ (1403.25 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (1346.18 kg ha⁻¹). While, minimum seed yield was recorded in control (1322.77 kg ha⁻¹). Interaction between phosphate solubilizing microorganisms and phosphorus levels was found to be significant. The seed inoculation with *Aspergillus awamori* and application of phosphorus @ 75 kg P₂O₅ ha⁻¹ recorded highest value of seed yield than rest of the interactions. This may attributed *Aspergillus awamori* increase more values of growth parameters at almost all growth stages and helped in reducing P fixation by its chelating effect and also solubilized the unavailable form of P leading to more uptake of nutrients resulted in better growth of the plant (Das *et al.*, 2013) ^[03]. The increase in seed yield due to increase in P level may be attributed to increase in the availability of P in soil. Similar findings are noted by Nawange *et al.* (2011) ^[13].

Table 7: Seed yield of chickpea as influenced by phosphate solubilizing microorganisms and phosphorus levels.

PSM	Phosphorus levels (P ₂ O ₅ kg ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Seed Yield kg ha ⁻¹				
T0	1198.85	1232.67	1303.35	1342.52	1269.35
T1	1344.64	1308.21	1354.48	1370.79	1344.53
T2	1345.15	1401.31	1416.74	1463.33	1406.63
T3	1402.44	1442.52	1538.42	1627.34	1502.68
Mean	1322.77	1346.18	1403.25	1451.00	
	T		P		TXP
SE	12.59		12.59		25.18
C.D at 5%	36.35		36.35		72.71

3.3.2 Straw Yield:

It is evidenced from the data presented in table 7. The straw yield of chickpea as significantly influenced by phosphate solubilizing microorganism, The seed inoculation with *Aspergillus awamori* produced maximum straw yield of chickpea (1803.2 kg ha⁻¹) as compared to application of *Aspergillus niger* (1688.0 kg ha⁻¹) and *Bacillus megaterium* (1613.4 kg ha⁻¹). Among the different phosphorus levels application of phosphorus @ 75 kg P₂O₅ ha⁻¹ recorded higher straw yield (1741.19 kg ha⁻¹) as compare to application of phosphorus @ 60 kg P₂O₅ ha⁻¹ (1683.90 kg ha⁻¹) and 45 kg P₂O₅ ha⁻¹ (1615.41 kg ha⁻¹). However, the minimum straw yield was recorded in control (1587.32 kg ha⁻¹). Interaction between phosphate solubilizing microorganisms and different phosphorus levels on straw yield was found to be significant. The seed inoculation with *Aspergillus awamori* and application of phosphorus @ 75 kg P₂O₅ ha⁻¹ recorded higher value of straw yield than rest of the interactions. This was mainly due to the fact that *Aspergillus awamori* and application of phosphorus @ 75 kg P₂O₅ ha⁻¹ increase in the availability of N and P caused better root development, better growth and development of plants and better diversion of photosynthates towards sink Tagore *et al.*, (2013) [23]. Kumar *et al.*, (2019) [10] reported that the straw yield of chickpea increased due to increase in phosphorus levels might be because of increase in the microbial activity in the root environment which accelerates cell division and formation of meristem.

Commented [A16]: It should be table 8. Kindly double check.

Table 8: Straw yield as influenced by phosphate solubilizing microorganisms and phosphorus levels in chickpea.

PSM	Phosphorus levels (kg P ₂ O ₅ ha ⁻¹)				Mean
	P0	P45	P60	P75	
	Straw Yield kg ha⁻¹				
T0	1438.6	1479.2	1564.0	1611.0	1523.2
T1	1613.6	1569.8	1625.4	1644.9	1613.4
T2	1614.2	1681.6	1700.1	1756.0	1688.0
T3	1682.9	1731.0	1846.1	1952.8	1803.2
Mean	1587.32	1615.41	1683.90	1741.19	
	T		P		TXP
SE	15.11		15.11		30.22
C.D at 5%	43.63		43.63		87.26

1. Conclusion

It can be inferred and concluded that, incorporation of phosphate solubilizing microorganisms viz. *Aspergillus awamori* in combination with application of

phosphorus @ 75 kg P₂O₅ ha⁻¹ improved N,P and K availability, uptake and seed yield of chickpea.

Commented [A17]: You can also state several future perspectives

References

1. A.O.A.C. (1965) Official methods of analysis of the chemicals, 10th Edn. Washington, D.C
2. Barros M.H, (2005). The *Saccharomyces cerevisiae* COQ10 gene encodes a START domain protein required for function of coenzyme Q in respiration. *J. Biol. Chem.* 280 (52): 42627-35.
3. Das, S., Pareek, B.L., Kumavat, A. and Dhikwal, S.R. (2013) Effect of phosphorus and biofertilizers on production of chickpea (*Cicer arietinum* L.) in north western Rajasthan, *Ind. Legume. Res.* 36 (6): 511- 514.
4. Bharadkar KS.(2021). Interactive effect of phosphate solubilizing microorganism and phosphorous levels on growth, soil nutrient dynamics, yield and quality of cowpea (*Vigna unguiculata* L.) in Inceptisols. M.Sc. (Agriculture) Thesis Vasantnao Naik Marathwada Krishi Vidyapeeth, Parbhani;
5. Biswas, P.K., Bhowmick, M.K., Kundu, M.C. Mondal, S. And Ghosh, G.K. (2015). Conjoint application of biofertilizer and phosphorous levels on growth, nodulation, nutrient uptake and productivity of lentil [*Lens culinaris* Medikus] in red and lateritic soils of West Bengal. *J. Crop. Weed.* 11(1):29-321.
6. Boparai, H.K., and Sharma, K.N. (2006). Phosphorus absorption and desorption characteristics of some soil as affected by clay and available phosphorus content. *J. Ind. Soci. Soil. Sci.* 54 (1):111-114.
7. Jackson ML.(1973). Soil Chemical Analysis. Prentice Hall of India, Pvt. Ltd. New Delhi;
8. Jain R., Saxena J., Sharma V. (2010). The evaluation of free and encapsulated *Aspergillus awamori* for phosphate solubilization in fermentation and soil-plan system. *App. Soil. Eco.* 46, 90-94.
9. Jahangir C.K.,singh.D.P.Meena R.H. and mahandra yadhav M (2017) Effect of fertility levels and biofertilizers on physical and chemical properties on soil under black gram (*Vigna mungo* L.)*Int.J.Curr.Microbio.App.Sci.*6(3):233-228

10. Kumar, J. Kumar, S. and Prakash V. (2019) effect of biofertilizers and phosphorus levels on soil fertility yield and nodulation in chickpea (*Cicer arietinum* L.). *J. Ind. Soci. Soil. Sci.* 67 (2):199-203.
11. Laharia G.S., Apotikar, V. Age A.B., Gite V. and Deshmukh D.P. (2019) Effect of phosphorus levels with PSB on yield nutrient use efficiency and uptake of nutrients by chickpea. *J. Pharma Phytochem.* 8(3):3128-3185
12. Mittal, V., Singh, O., Nayyar, H., Kaur, J. and Tewari, R. (2008). Stimulatory effect of phosphate-solubilizing fungal strains (*Aspergillus awamori* and *Penicillium citrinum*) on the yield of chickpea (*Cicer arietinum* L. cv. GPF2). *Soil. Bio. Bioche.* (40) 718-727.
13. Nawange, D.D., Yadav, A.S. and Singh, R.V. (2011). Effect of phosphorus and sulphur application on growth, yield attributes and yield of chickpea (*Cicer arietinum* L.). *Legume. Res.* 34(1):48-50.
14. Narsihan, V., Patel, H.H. (2000). *Aspergillus aculeatus* as a rock phosphate solubilizer. *Soil. Bio. Bioche.* (32):559-565.
15. Panse VG, Sukhatme PK. (1985) Statistical methods for agriculture workers, (IV Edn.) ICAR, New Delhi. 1985, 145-156.
16. Paratey, P.R. And Wani, P.V. (2005). Response of soybean to phosphate solubilizing biofertilizers. *Legume. Res.* 28(4): 268-271.
17. Piper, C.S. (1966) Soil and Plant Analysis, Asian Reprint Hans publishers, Bombay. 368 p.
18. Prajapati, B.J., Gudadhe, N. Gamit, V.R. and Chhaganiya, H.J. (2018). Effect of integrated phosphorus management on growth, yield attributes and yield of chickpea. *Fmg. & Mngmt.* 2(1): 36-40.
19. Rooge, R. B., Patil, V. C. and Ravikishan, P. (1998). Effect of phosphorus application with phosphate solubilizing organisms on the yield, quality and phosphorus uptake of soybean. *Legume. Res.* 21: 85-90.
20. Singh, R., Pratap, T., Singh, D., Singh, G. and Singh, A.K. (2014). Effect of phosphorus, sulphur and biofertilizers on growth attributes and yield of chickpea (*Cicer arietinum* L.). *J. Pharma. Phytochem.* 7(2): 3871-3875.
21. Srinivasarao, C.H. Ganeshamurthy, A.N. Ali, M. and Venkateswarlu, B. (2003). Phosphorus and micronutrient nutrition of chickpea genotypes in a multi-nutrient deficient typic ustochrept. *J. Plant. Nutr.* 29: 747-763.
22. Subbiah BV, Asija GL. Rapid procedure for the estimation of available

nitrogen in soil. *Current Science*. 1956; 25: 259-260.

23. Tagore, G.S., Namdeo, S.L., Sharma, S.K. and Kumar, N. (2013). Effect of *Rhizobium* and Phosphate Solubilizing Bacterial Inoculants on Symbiotic Traits, Nodule Leghemoglobin, and Yield of Chickpea Genotypes. *Int. J. Agron.* 8:581627.
24. Yadhav K.R., Manohar R.S, Kumwaty S.R. and yadhav v.k.(2017)Effect of phosphorus source and phosphorus solubilizing microorganisms on growth and yield of mungbean [*Vigna radiate*(L.)Wilczek].*Chem.SciRev.Lett*,6 (22):1152-1155
25. Vidhyashree,V., Naga, S.R., Yadav, B.L., Koli D.K. and Rao I.J. (2017). Effect of phosphorus and biofertilizers on growth and yield of mungbean [*Vigna radiate* (L.)Wilczek]. *Int. J. Curr. Micro. App. Sci.* 6(7): 3992- 3997.
26. Walkley A, Black CA. An estimation of method of determining soil organic matter & proposed modification of the chromic acid titration method. *Indian Journal of Agronomy*. 1934;66:544.

UNDER PEER REVIEW