

Integrated Effect of Organic Manures and Organic Liquid Formulations on Growth, Yield and Quality of Garlic cv. Agrifound Parvati in Bharsar Condition of Uttarakhand

ABSTRACT

The field study was carried out to VCSG, UHF, Pauri Garhwal in 2022-2023 to investigate the integrated effect of organic manures and organic liquid formulations on the growth and yield of garlic. Among fourteen different treatment combinations that consist of various combinations of organic manures (FYM and Vermicompost) and organic liquid formulations (Panchgavya and Jeevamrit) treatment T₃ i.e., FYM (10t/ha) + Panchgavya (2%) emerged as the best treatment concerning growth and yield of garlic. The findings showed that the maximum plant height (69.080 cm), number of leaves (8.400), leaf length (47.880 cm) and breadth (3.207 cm), and leaf area index (1.023) at all the 30 days intervals after sowing of garlic has been observed with the application of Treatment T₃. The fundamental reason for the increased growth might be due to growth-promoting substances like auxins and gibberellins in panchgavya and high nitrogen and phosphorus in FYM. Concerning yield and quality attributes, the bulb diameter (5.657 cm), fresh weight of bulb (64.643 cm), length of the bulb (5.853 cm), number of cloves per bulb (12.200), bulb yield (120.833q), total chlorophyll content (2.127 mg/g) and ascorbic acid content (15.333 mg/100g) was obtained maximum with the application of same treatment T₃ i.e., FYM (10t/ha) + Panchgavya (2%). However, the harvest index percent was found to be maximum (89.243 %) in treatment T₁₃ i.e., Vermicompost (5t/ha) + Jeevamrit (6%). Therefore, the application of FYM (10t/ha) along with a foliar spray of panchgavya is considered a better proposition for enhancing the growth, and yield of garlic.

Keywords: *Allium sativum*, FYM, Panchgavya, Jeevamrit, Growth Parameters

INTRODUCTION

Garlic (*Allium sativum* L.), after onions, is the second most important crop grown from bulbs which is used as a spice or condiment. It is often referred to as Lahsun in India and is an element of the Amaryllidaceae family. Some scholars believe that the wild ancestor of garlic is *Allium longicuspis* Regael. With 75% of the world's production, China is by far the greatest producer of garlic worldwide. India ranks second in terms of both area and production, but its productivity—5.22 tons/hectare—is extremely poor (Gupta *et*

al., 2012). It is cultivated on 3.21 lakh hectares on average, with an average yield of 5.27 t/ha, producing 1693 thousand metric tonnes. (Anonymous, 2017).

In the Financial Year 2022, more than two million metric tonnes of garlic were produced. Madhya Pradesh dominated all other Indian states in this regard (Statista, 2022). Garlic production in Uttarakhand is 11,270.19 metric tonnes, while its area is 1924.23 hectares (State Horticulture Mission, Horticulture production data, 2021).

In the tradition of Ayurveda, garlic possesses healing and medicinal properties; it is usually more in demand in the winter. Moreover, it has high levels of protein (6.30%), phosphorus (0.30%), magnesium (71 mg/100 g), and ascorbic acid content (13 mg/100 g) in addition to carbohydrates (29%).

It is a perennial crop that is hardy to frost and has narrow leaves. When the bulb reaches maturity, it needs a fairly dry period during growth and a chilly, humid temperature throughout maturity. Due to environmental concerns and greater awareness of global health, there is a growing need for organic food products. The irresponsible application of chemicals created pollution, a decline in soil fertility and productivity, and other negative consequences on the environment, water, and soil. Without a doubt, the application of organic fertilizer is crucial for modern agriculture as a means to achieve both better yields per unit area and crops of high quality.

The vast majority of small and marginal farmers in the entire country cannot afford the highly expensive chemical fertilizers. Its long-term negligent use has reduced microbial activity, harmed the environment, degraded soil fertility and health, and limited the availability of vital nutrients. As a result, organic liquid formulations and manures are inexpensive,

environmentally friendly, as well as financially beneficial for farmers. The principal organic manures include neem cake, FYM, and vermicompost. FYM is considered to be an ideal source of nutrients for soil microbiology and plant growth.

Vermicompost, which originates using a variety of worm species through the decomposition process—most often red wigglers, white worms, and other earthworms—improves soil fertility, aggregation, and structure. Organic liquid formulations, such as Panchgavya, Jeevamrit, Sanjivak, Bijamrita, Amritpani, and Vermiwash, are products derived from the fermentation or decomposition of organic matter. Panchgavya provides resistance to the plant and can stimulate growth. Its name suggests that it is composed of five ingredients: cow dung, urine, milk, curd, and ghee. Yet another natural liquid fertilizer that is extremely high in biomass and natural carbon and has every nutrient that's needed is known as Jeevamrit. Therefore, organic manures and organic liquid formulations are eco-friendly, cheap sources of nutrients, and are also cost-effective for farmers. Along with enhancing soil microbial population, slow nutrient release, fertilizer usage efficiency, and soil phosphate availability, they additionally increase cation exchange capacity, water retention capacity, and soil

phosphate availability.(Tadesse *et al.*, 2014).

Because inorganic fertilizers are used improperly, the texture and structure of the soil degrade, microbial activity declines, groundwater is contaminated, and eventually soil fertility and production are reduced. Whereas using organic manures improves output and reduces environmental risks by improving the texture, structure, humus, color, aeration, water-holding capacity, microbial activity, and nutrient-use efficiency of the soil (Pare

et al., 2000). The main cause of India's decreasing garlic output is the scanty and improper use of agronomic methods. Nowadays, the export market prefers and demands vegetables grown with very little or no inorganic fertilizer use to fetch higher prices. Garlic needs to be made more profitable by using additional nutrient sources in addition to ensuring good soil health for sustainable agriculture. This is crucial since garlic is one of the crops that has the potential to be eaten and marketed.

MATERIALS AND METHODS

A field experiment was carried out on garlic cv. Agrifound Parvati during the winter (rabi) season of 2022 at Organic Research Block, College of Horticulture, VCSG Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri Garhwal, Uttarakhand. The soil type of the experimental field was classified as clay-loam soil with a pH of 6.06, with organic carbon content of 1.47 % indicating sufficient organic matter. The available Nitrogen, P₂O₅, and K₂O in the soil were 524.96 kg/ha, 39.66 kg/ha, and 45.25 kg/ha respectively. The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications.

Each replication consists of fourteen treatments, *viz.* organic manures *viz.*, FYM, and Vermicompost as basal, Organic liquid formulations *viz.* Panchgavya and Jeevamrit at different concentrations were applied as foliar application at 30-day intervals after sowing. The sowing process involves sowing cloves at a spacing of 15cm x 10cm in a plot size of 0.8 m² each. The panchgavya that was utilized consisted mainly of five products *viz.* Cow dung, Cow urine, Cow milk, Curd, and Cow ghee. Other components like jaggery, ripened bananas, tender coconut, and water

were also used to boost the fermentation process (Sreenivasa *et al.* 2011). Jeevamrit, a wonderful source of natural carbon, nitrogen, phosphorous, potassium, and many other micronutrients required by

the crops, was another organic formulation used. It consists of a mixture of cow dung, cow urine, jaggery, pulse flour, and a handful of fertile soil in water.

RESULTS AND DISCUSSION

Growth Parameters

Application of organic liquid formulations and manures had a substantial impact on plant height, leaf area index, number of leaves, average length of leaves, average breadth of leaves, and average length of leaves. After sowing, data was collected every 30 days. Table 1 and 2 presents a summary of this. The maximum plant height (69.080 cm) was recorded with the application of treatment T₃ *i.e.*, FYM (10t/ha) + Panchgavya (2%) at 180 days after sowing of garlic respectively, Auxins and gibberellin, the two growth boosters found in Panchgavya, is considerably the reason for this. Conversely, the minimum was obtained under control. Similar results have been reported by Kumaravelu and Kadamban (2009) in green gram, Mathewset *al.* (2017) in cowpea, tomato, lady finger, chilli and brinjal, Sornalatha *et al.* (2018) in *Luffa acutangula*, Bua *et al.* (2017) in onion, Ranpariya *et al.* (2020) in garlic and Patle *et al.* (2021) in garlic. The maximum number of leaves at 180 days

after sowing of garlic was again obtained with the application of FYM (10t/ha) + Panchgavya (2%) which is due to the presence of appreciable amounts of major nutrients in Panchgavya. High nitrogen and phosphorus content in FYM causes more photosynthesis and, therefore greater number of leaves. Similar results have also been reported by Lal *et al.*, (2002) in onion, Yassen and Khalid (2009) in onion, Mathews *et al.*, (2017) in cowpea, tomato, lady finger, chilli and brinjal and Bua *et al.*, (2017) in onion.

Concerning the average length of leaves, the maximum value (47.880 cm) was obtained with the application of treatment T₃ at 180 days after sowing. Moreover, a higher dose of FYM provides more amount of nutrients to the plant, and the presence of magnesium in it helped in the synthesis of chlorophyll, which in turn increased the rate of photosynthesis resulting in higher leaf length. Similar outcomes were put out by Kumaravelu and

Kadamban (2009) in Greengram, Premsekhar and Rajshree (2009) in Okra, Maheshwari and Rajkumar (2020) in Cluster Onion and Pallavi and Anuja (2019) in Moringa. Applying the same treatment T₃ again showed the highest breadth of *i.e.*, 3.207 cm at 180 days after sowing. This may be caused by the gibberellin and auxins present in

panchgavya in addition to the micronutrients that FYM had supplied. With the application of treatment T₃, the leaf area index was also observed at its maximum *viz.*, 1.023 at 180 days after sowing. An increase in LAI may have been relied on by the bigger leaf surface area. The respective parameter's lowest readings were found under control.

UNDER PEER REVIEW

Table 1: Integrated effect of organic manures and organic liquid formulations on plant height (cm), Number of Leaves per plant, and Average Length of Leaves of Garlic (*Allium sativum* L.) at 180 DAS

	Treatments	Plant Height (cm)	Number of Leaves per Plant	Average Length of Leaves (cm)
T₁	Control	55.127	7.133	38.347
T₂	FYM (10t/ha) + Vermicompost (5t/ha)	61.153*	7.467	40.893
T₃	FYM (10t/ha) + Panchgavya (2%)	69.080*	8.400*	47.880*
T₄	FYM (10t/ha) + Panchgavya (4%)	60.767*	7.800*	43.160*
T₅	FYM (10t/ha) + Panchgavya (6%)	65.780*	8.000*	43.433*
T₆	FYM (10t/ha) + Jeevamrit (4%)	56.640	8.200*	39.993
T₇	FYM (10t/ha) + Jeevamrit (6%)	67.000*	8.133*	43.400*
T₈	FYM (10t/ha) + Jeevamrit (8%)	67.467*	7.933*	43.600*
T₉	Vermicompost (5t/ha) + Panchgavya (2%)	56.487	8.267*	38.860
T₁₀	Vermicompost (5t/ha) + Panchgavya (4%)	58.067	7.867*	40.213
T₁₁	Vermicompost (5t/ha) + Panchgavya (6%)	59.057	8.067*	41.747
T₁₂	Vermicompost (5t/ha) + Jeevamrit (4%)	59.713*	7.667	39.027
T₁₃	Vermicompost (5t/ha) + Jeevamrit (6%)	59.553	7.933*	40.147
T₁₄	Vermicompost (5t/ha) + Jeevamrit (8%)	59.003	7.867*	38.420
	S.E (d)	2.175	0.319	1.832
	C.D (0.05)	4.495	0.659	3.787

*Significant at 5% level of significance as compared with T₁ (Control)

	Treatments	Average Breadth of Leaves (cm)	Leaf Area Index
T ₁	Control	2.427	0.630
T ₂	FYM (10t/ha) + Vermicompost (5t/ha)	2.833*	0.773*
T ₃	FYM (10t/ha) + Panchgavya (2%)	3.207*	1.023*
T ₄	FYM (10t/ha) + Panchgavya (4%)	2.793*	0.807*
T ₅	FYM (10t/ha) + Panchgavya (6%)	2.820*	0.820*
T ₆	FYM (10t/ha) + Jeevamrit (4%)	2.633	0.703

Table 2: Integrated effect of organic manures and organic liquid formulations on Average Breadth of Leaves (cm) and Leaf Area Index of Garlic (*Allium sativum* L.) at 180 DAS

	T₇	FYM (10t/ha) + Jeevamrit (6%)		2.800*		0.810*		
	T₈	FYM (10t/ha) + Jeevamrit (8%)		2.607		0.757*		
	T₉	Vermicompost (5t/ha) + Panchgavya (2%)		2.653*		0.683		
		Treatment details	Bulb diameter (cm)	Fresh weight (g)	Length of bulb (cm)	No. of cloves	Bulb yield (q/ha)	Harvest Index (%)
	T₁₀	Vermicompost (5t/ha) + Panchgavya (4%)		2.620		0.667		
	T₁₁	Vermicompost (5t/ha) + Panchgavya (6%)		2.673*		0.743*		
	T₁₂	Vermicompost (5t/ha) + Jeevamrit (4%)		2.440		0.650		
T₁	Control		4.563	23.600	4.273	9.733	43.750	73.297
	T₁₃	Vermicompost (5t/ha) + Jeevamrit (6%)		2.780*		0.743*		
	T₁₄	Vermicompost (5t/ha) + Jeevamrit (8%)		2.613		0.670		
		S.E (d)		0.100		0.052		
		C.D_(0.05)		0.207		0.107		

*Significant at 5% level of significance as compared with T₁ (Control)

Table 3: Integrated effect of organic manures and organic liquid formulations on yield attributes of garlic (*Allium sativum* L.)

T₂	FYM (10t/ha) + Vermicompost (5t/ha)	5.020	52.643*	5.013*	11.200*	98.327*	79.183
T₃	FYM (10t/ha) + Panchgavya (2%)	5.657*	64.643*	5.853*	12.200*	120.833*	88.117*
T₄	FYM (10t/ha) + Panchgavya (4%)	5.117	49.867*	5.347*	10.533	92.917*	84.967*
T₅	FYM (10t/ha) + Panchgavya (6%)	5.297*	58.490*	5.160*	10.800	109.167*	75.417
T₆	FYM (10t/ha) + Jeevamrit (4%)	5.110	53.053*	5.453*	11.067*	98.790*	84.017*
T₇	FYM (10t/ha) + Jeevamrit (6%)	5.560*	60.547*	5.440*	11.267*	112.917*	80.773
T₈	FYM (10t/ha) + Jeevamrit (8%)	5.380*	56.747*	5.413*	11.533*	105.833*	76.400
T₉	Vermicompost (5t/ha) + Panchgavya (2%)	4.650	44.830*	5.253*	10.533	82.917*	83.183*
T₁₀	Vermicompost (5t/ha) + Panchgavya (4%)	5.017	47.067*	5.040*	10.933	87.917*	84.580*
T₁₁	Vermicompost (5t/ha) + Panchgavya (6%)	5.247*	50.303*	5.133*	10.733	93.753*	78.930
T₁₂	Vermicompost (5t/ha) + Jeevamrit (4%)	4.947	45.480*	5.227*	10.400	84.580*	85.337*
T₁₃	Vermicompost (5t/ha) + Jeevamrit (6%)	4.690	47.193*	5.080*	9.933	88.750*	89.243*
T₁₄	Vermicompost (5t/ha) + Jeevamrit (8%)	4.903	45.513*	5.067*	10.667	85.003*	83.053*
S.E. (d)		0.299	3.760	0.285	0.594	5.586	4.605
C.D_(0.05)		0.618	7.772	0.590	1.228	11.546	9.518

*Significant at a 5% level of significance as compared with T₁ (Control)

Yield Parameters

The application of treatment T₃, or [FYM (10t/ha) + Panchgavya (2%)], yielded the

largest bulb diameter (5.657 cm), fresh weight of the bulb (64.643 g), length of the bulb (5.853 cm), number of cloves (12.200), and bulb yield (120.833 q/ha) because nutrients and growth stimulants could be supplied easily, resulting in enhanced bulb diameter and length. The findings are in close conformity with the findings of Anbarasi and Haripriya (2020) in onion and Ranpariya *et al.*, (2020) in garlic. The application of treatment T₁₃ [Vermicompost (5t/ha) + Jeevamrit (6%)] produced the maximum harvest index (89.243%), which was followed by T₃ (88.117%). These results are closely agreed with the results obtained by Yadav *et al.*, (2017) in garlic, and Kenea and Gedamu (2019) in garlic. With a net return of 3,66,494.0 ₹/ha, treatment T₃ [FYM (10t/ha) + Panchgavya (2%)] was the most cost-effective. The highest C: B

ratio, however, was attained with treatment T₇, which is FYM (10t/ha) + Jeevamrit (6%).

Quality Parameters

The highest total chlorophyll content (2.127 mg/ g) and ascorbic acid content (15.333 mg/100 g) were observed with the application of treatment T₃ [FYM (10t/ha) + Panchgavya (2%)]. The primary reason is the increasing leaf area of the plant, which causes more photosynthetic activity, resulting in higher production of photosynthetic products and their accumulation in the plant. The augment of ascorbic acid might be due to the good growth of plants resulting from higher assimilation of micronutrients which are made available due to well-decomposed organic matter.

Table 4: Integrated effect of organic manures and organic liquid formulations on qualitative analysis of Garlic (*Allium sativum* L.)

Treatment code	Treatment Details	Total chlorophyll (mg/g)	Ascorbic acid content (mg/ 100 g)
T₁	Control	0.883	6.000
T₂	FYM (10t/ha) + Vermicompost (5t/ha)	1.057*	6.667
T₃	FYM (10t/ha) + Panchgavya (2%)	2.127*	15.333*
T₄	FYM (10t/ha) + Panchgavya (4%)	1.930*	14.667*
T₅	FYM (10t/ha) + Panchgavya (6%)	1.777*	11.333*
T₆	FYM (10t/ha) + Jeevamrit (4%)	1.583*	9.333*
T₇	FYM (10t/ha) + Jeevamrit (6%)	1.443*	8.000
T₈	FYM (10t/ha) + Jeevamrit (8%)	1.570*	8.667*
T₉	Vermicompost (5t/ha) + Panchgavya (2%)	1.707*	10.667*
T₁₀	Vermicompost (5t/ha) + Panchgavya (4%)	1.873*	13.333*
T₁₁	Vermicompost (5t/ha) + Panchgavya (6%)	1.227*	7.333
T₁₂	Vermicompost (5t/ha) + Jeevamrit (4%)	1.637*	10.000*
T₁₃	Vermicompost (5t/ha) + Jeevamrit (6%)	1.817*	12.000*
T₁₄	Vermicompost (5t/ha) + Jeevamrit (8%)	1.837*	12.667*
S.E. (d)		0.076	1.133
C.D. (0.05)		0.156	2.343

*Significant at a 5% level of significance as compared with T₁ (Control)

CONCLUSION

After the investigation, it was revealed that treatment T₃ [FYM (10t/ha) + Panchgavya (2%)] recorded significantly highest plant height, number of leaves, average length and breadth of leaves, leaf area index, bulb diameter, fresh weight of the bulb, length of the bulb, number of cloves and bulb yield. The maximum C: B ratio was obtained under treatment T₇ *i.e.*, FYM (10t/ha) + Jeevamrit (6%).

REFERENCES

- Anbarasi, D. and HariPriya, K. 2020. Response of aggregatum onion (*Allium cepa* L. var. *aggregatum* Don.) to organic inputs, biofertilizers and biostimulants. *Plant*

Archives. 20: 759-762.

Anonymous 2017. Third advance estimates of year 2016, Indian Horticulture Database. National Horticulture Board Gurgaon. pp. 7.

Bua, B., Owiny, R. and Ocwa, A. 2017. Response of Onion to Different Organic Amendments in Central Uganda. *Journal of Agricultural Science and Technology*. 79-85.

Gupta, R. P., Bhonde, S. R. and Gupta, H. P. 2012. National Horticultural Research and Development Foundation. Vision 2030.

Kenea, F. T. and Gedamu, F. 2019. Effect of vermicompost on growth, quality and economic return of garlic (*Allium sativum* L.). *African Journal of Agricultural Research*. 14(35): pp. 2159-2167.

Kumaravelu, G. and Kadamban, D. 2009. Panchagavya and its effect on the growth of the greengram cultivar K-851. *International Research Journal of Plant Science*. 4(2): 409-414.

Lal, S., Yadav, A. C., Mangal, J. L., Singh, A. and Batra, V. K. 2002. Effect of FYM and irrigation levels on growth and yield of onion cv. Hisar – 2, Haryana. *Journal of Horticultural Sciences*. 31 (3-4): 256-258.

Maheshwari, T. U. and Rajkumar, M. 2020. Effect of Biosolarization on the growth attributes of cluster onion (*Allium cepa* L. var. *Aggregatum* Don). *The Journal of Research ANGRAU*. 48(1): 87-91.

Mathews, N. M. S., Bharani, A. and Nandhini, D. U. 2017. Influence of organic inputs on growth parameters of vegetable crops under terrace farming. *International Journal of Chemical Studies*. 5(5): 763-765.

Pallavi, N. and Anuja, S. 2019. Effect of organic nutrients on growth parameters of moringa (*Moringa oleifera* lam.) for leaf production. *Plant Archives*. 19(2): 2439-2442.

Pare, T., Dinel, H. and Schnitzer, M. 2000. Carbon and nitrogen mineralization in soil amended with non-tabletized and tabletized poultry manure. *Canadian Journal of Soil Science*. 80(2): 271-282.

- Patle, A. K., Singh, S. S., Jadia, M. and Verma, K. S. 2021. Effect of organic, inorganic sources of nutrients on growth of garlic (*Allium sativum* L.). *Journal of Pharmacognosy and Phytochemistry*. 10(1): 2214-2217.
- Premsekhar, M. and Rajashree, V. 2009. Influence of Organic Manures on Growth, Yield and Quality of Okra. *American-Eurasian Journal of Sustainable Agriculture*, 6-8.
- Ranpariya, V. S., Parmar, K. B., Vekariya, L. C., Chovatia, P. K. and Jadeja, A. S. 2020. Effect of nutrient management through organic sources on yield and quality of Garlic (*Allium sativum* L.) under organic farming. *International Journal of Chemical Studies*. 8(4): 3663- 3668.
- Sornalatha, S., Tamilarasi, M. and Esakkiammal, B. 2018. Effect of liquid organic fertilizer of panchgavya on growth and development of *Luffa acutangula*. *Scire Science Multidisciplinary Journal*. 2(1):72-81.
- Sreenivasa, M. N., Naik, N. and Bhatt. 2011. Nutrient status and microbial load of different organic liquid manures. *Karnataka Journal of Agricultural Science*. 24: 583-584.
- State Horticulture Mission, 2021. Horticulture Production Data 2020- 2021. Available at: <https://shm.uk.gov.in/> (accessed date- 22 March 2023).
- Statista. 2022. Statista research department. Available at <https://www.statista.com/statistics/744339/india-garlic-production-by-state/> (accessed date- 6 October 2022).
- Tadesse, T., Dechassa, N., Bayu, W. and Gebeyehu, S. 2014. Effects of farm yard manure and inorganic fertilizer application on soil physico-chemical properties and nutrient balance in rainfed lowland rice ecosystem. *Am. Journal of Plant Science*. 4: 309-316.
- Yadav, R. N., Bairwa, H. L. and Gurjar, M. K. 2017. Response of Garlic (*Allium sativum* L.) to Organic Manures and Fertilizers. *International Journal of Current Microbiology and Applied Sciences*. 6(10): 4860-4867.
- Yaseen, A. A. A. and Khalid, K. A. 2009. Influence of organic fertilizers on the yield, essential oil and mineral content of onion. *International Agrophysics*. 23: 183-188.

UNDER PEER REVIEW