

CASE REPORT

Autosomal recessive spastic paraplegia type 51 caused by homozygous mutation of the AP4E1 gene: A Case Report of a 17-years-old boy

Abstract

Autosomal recessive spastic paraplegia-51 (SPG51) is a rare neurodevelopmental disorder caused by a homozygous mutation in the AP4E1 gene, located on chromosome 15q21. Spastic paraplegia-51 (SPG51) is an extremely rare autosomal recessive neurodevelopmental disorder characterized by neonatal hypotonia that progresses to hypertonia and spasticity. Mutations in the AP-4E1 gene on chromosome 15q21.2, which is part of the AP-4 complex, can lead to autosomal recessive spastic paraplegia 51 (SPG51). Different mutations in this gene have been identified in affected individuals, leading to disruption of the AP4 complex and vesicular trafficking processes. The specific characteristics of SPG51 are, due to only few cases, not well understood due to limited reports of affected families. Affected individuals also have global developmental delay with impaired intellectual development and poor or absent speech. Studies linked to an additive symptom of persistent stuttering. Affected individuals typically present with hypotonia in the neonatal period, which progresses to muscular hypertonia, especially in the lower limbs. They may also exhibit contractures, talipes equinovarus, decreased muscle mass, short stature, and microcephaly. Severe mental retardation, absent speech, and dysmorphic facial features are common. Some patients may experience seizures. Neurological examination often reveals spastic paraplegia of the lower limbs, and brain imaging may show atrophy of the cerebellar vermis and cortical atrophy. We present an extremely rare case of a 17 years-old boy with autosomal recessive spastic paraplegia type 51.

Keywords: autosomal recessive, SPG51, mutation, hypertonia

Introduction

"Hereditary spastic paraplegia (HSP) refers to a group of genetically and clinically heterogeneous neurodegenerative disorders characterized primarily by spasticity and weakness in the lower limbs. It includes four genetic inheritance forms: autosomal dominant inheritance (AD), autosomal recessive inheritance (AR), X-linked inheritance, and mitochondrial inheritance. HSP patients may have either pure or complicated HSP, differing based on symptoms. Patients with pure HSP simply develop spasticity and weakness of the lower limbs, while patients with complicated HSP are often accompanied by other symptoms, such as early cognitive impairment, ataxia, visual disturbance, macular degeneration, dysarthria, and callosal agenesis" [18].

Case Report

We present a case of a 17-years-old German boy with an autosomal recessive spastic paraplegia type 51 (SPG51). A compound heterozygous mutation in the AP4E1 gene was found in genetic analysis. An unclassifiable epilepsy most likely with complex focal seizures is mostly present in SPG51. Perampanel was initiated as an anticonvulsant therapy after previous treatments with Sultiam, Lacosamide, Lamotrigine, Oxcarbazepine, Levetiracetam, and Zonisamide. SPG51 is a combined developmental disorder and a severe expressive language development disorder. Past medical history showed an epileptological situation quite stable. No major seizure events were observed anymore, occasionally episodes of smacking occur, which were interpreted differently. These episodes were hardly noticeable and very brief with a duration of less than a minute. The current medication consisted of Baclofen 10-0-20 mg and Perampanel 3.5 mg. The main problem affecting both school and especially the home environment were extreme episodes of restlessness and aggression, provoked by unfamiliar people and environments. They also occur in the home environment. Examination findings included the patient in a fairly good general condition, quite communicative, but mostly with non-verbal communication. Muscle tone increasingly spastic distally and leg-dominant, there were no side differences, no clearly recognizable cranial nerve abnormalities. The EEG showed a polymorphic mixed wave rhythm, emphasized in the upper theta frequency spectrum. The amplitudes barely exceeded 70 volts. An on/off effect was hardly discernible. Plenty of muscle activity overlaid, but no specific pathological patterns, especially no seizure predisposition, was found. The assessment and recommendations included the epileptological situation was not the main focus, rather the behavioral disorder was hardly bearable.

Discussion

"Most of all hereditary spastic paraplegias (HSP) are classified due to the genetic loci that are classified by numbers according to their discovery (1-17). Four types are described in HSP and have been related to disturbances of different adaptor protein complex subunits" (4). "The functional loss of the AP-4 protein complex is the molecular mechanism in all AP-4 HSP disorders (4,10). AP-4-associated hereditary spastic paraplegia (HSP) is a group of progressive neurodegenerative disorders that typically present in infancy or early childhood. Spastic paraplegia type 51 is a genetic disorder characterized by the degeneration and dysfunction of certain nerve tracts (1-17). The AP-4E1 subunit is involved in recognizing and binding sorting signals within cells, which is crucial for proper protein localization in neurons. It is involved in sorting proteins to specific cell membranes and establishing protein localization in neurons" (4,8). The AP4E1 gene encodes a member of the adaptor complexes large subunit protein family, which are essential components of adaptor protein complexes involved in vesicle formation and sorting of integral membrane proteins in the secretory and endocytic pathways. The encoded protein is a large subunit of adaptor protein complex-4, associated with both clathrin- and nonclathrin-coated vesicles. Mutations in this gene may be linked to cerebral palsy. Multiple isoforms of this gene have been identified due to alternative splicing. This gene is expressed ubiquitously, including in lymph nodes. AP4 complex-mediated trafficking is essential for brain development, with the AP4E1 gene encoding the AP-4 adaptor protein complex subunit $\epsilon - 1$. Knock-out mice with mutations in this gene exhibit abnormal white blood counts, enlarged lateral ventricles, a smaller corpus callosum, and hypoferrremia. The clinical phenotype observed in this study aligns with previous reports of SPG51. Further mutational analysis and evaluation of AP4E1 in individuals with spastic paraplegia and their families are warranted to assess the disorder's frequency.

This condition is often mistaken for cerebral palsy, but differs in its association with parental age and birth order (5,9,14). Symptoms include hypotonia progressing to spasticity, leading to difficulty walking or standing. Upper extremities may also be affected, resulting in spastic tetraplegia. "Other complications include bladder and bowel dysfunction, dysphagia, foot deformities, and contractures" (3,9,12,14). "Seizures occur in about half of patients, and the condition is associated with microcephaly and developmental delays. Speech development is delayed or absent, stuttering could be rarely found" (6). One case report describes the genetic defect with a mycobacterial disease (15).

Conclusion

SPG51 is a rare and often underdiagnosed condition in childhood, frequently presenting with clinical inconsistencies that can lead to misdiagnosis. This case highlights the importance of genetic testing and careful clinical evaluation in suspected cases of HSP.

Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

Consent

As per international standards, parental written consent has been collected and preserved by the author(s).

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

References:

- (1) AbouJamra R, Philippe O, Raas-Rothschild A, Eck SH, Graf E, Buchert R, Borck G, Ekici A, Brockschmidt FF, Nöthen MM, Munnich A, Strom TM, Reis A, Colleaux L. Adaptor protein complex 4 deficiency causes severe autosomal-recessive intellectual disability, progressive spastic paraplegia, shy character, and short stature. *Am J Hum Genet.* 2011 Jun 10;88(6):788-795. doi: 10.1016/j.ajhg.2011.04.019. Epub 2011 May 27. PMID: 21620353; PMCID: PMC3113253.
- (2) Kong XF, Bousfiha A, Rouissi A, Itan Y, Abhyankar A, Bryant V, Okada S, Ailal F, Bustamante J, Casanova JL, Hirst J, Boisson-Dupuis S. A novel homozygous p.R1105X mutation of the AP4E1 gene in twins with hereditary spastic paraplegia and mycobacterial disease. *PLoS One.* 2013;8(3):e58286. doi: 10.1371/journal.pone.0058286. Epub 2013 Mar 5. PMID: 23472171; PMCID: PMC3589270.
- (3) Moreno-De-Luca A, Helmers SL, Mao H, Burns TG, Melton AM, Schmidt KR, Fernhoff PM, Ledbetter DH, Martin CL. Adaptor protein complex-4 (AP-4) deficiency causes a novel autosomal recessive cerebral palsy syndrome with microcephaly and intellectual disability. *J Med Genet.* 2011 Feb;48(2):141-4. doi: 10.1136/jmg.2010.082263. Epub 2010 Oct 23. PMID: 20972249; PMCID: PMC3150730.
- (4) Manoochehri J, Goodarzi HR, Tabei SMB. Spastic paraplegia 51: phenotypic spectrum related to novel homozygous *AP4E1* mutation. *J Genet.* 2022;101:40. PMID: 36226339.
- (5) Abdollahpour H, Alawi M, Kortüm F, Beckstette M, Seemanova E, Komárek V, Rosenberger G, Kutsche K. An AP4B1 frameshift mutation in siblings with intellectual disability and spastic tetraplegia further delineates the AP-4 deficiency syndrome. *Eur J Hum Genet.* 2015 Feb;23(2):256-9. doi: 10.1038/ejhg.2014.73. Epub 2014 Apr 30. PMID: 24781758; PMCID: PMC4297901.
- (6) Chow HM, Li H, Liu S, Frigerio-Domingues C, Drayna D. Neuroanatomical anomalies associated with rare *AP4E1* mutations in people who stutter. *Brain Commun.* 2021 Nov 13;3(4):fcab266. doi: 10.1093/braincomms/fcab266. PMID: 34859215; PMCID: PMC8633735.
- (7) Winkler I, Miotła P, Lejman M, Pietrzyk A, Kacprzak M, Kubiak M, Sobczyńska-Tomaszewska A, Skrzypczak M, Jaszczuk I. A new family with spastic paraplegia type 51 and novel mutations in AP4E1. *BMC Med Genomics.* 2021 May 18;14(1):131. doi: 10.1186/s12920-021-00980-5. PMID: 34006278; PMCID: PMC8130362.
- (8) Sager G, Turkyilmaz A, Ates EA, Kutlubay B. HACE1, GLRX5, and ELP2 gene variant cause spastic paraplegias. *Acta Neurol Belg.* 2022 Apr;122(2):391-399. doi: 10.1007/s13760-021-01649-7. Epub 2021 Apr 3. Erratum in: *Acta Neurol Belg.* 2023 Oct;123(5):2047. doi: 10.1007/s13760-021-01685-3. PMID: 33813722.
- (9) Rudenskaya GE, Guseva DM, Mironovich OL, Kadnikova VA, Dadali EL, Komar'kov IF, Novoselova OG, Ryzhkova OP. AP4-svyazannye nasledstvennyespasticheskieparaplegii [AP4-associated hereditary spastic paraplegias]. *ZhNeвроlPsikhiatrIm S SKorsakova.* 2021;121(2):71-78. Russian. doi: 10.17116/jnevro202112102171. PMID: 33728854.
- (10) Ebrahimi-Fakhari D, Teinert J, Behne R, Wimmer M, D'Amore A, Eberhardt K, Brechmann B, Ziegler M, Jensen DM, Nagabhyrava P, Geisel G, Carmody E, Shamshad U, Dies KA, Yuskaitis CJ, Salussolia CL, Ebrahimi-Fakhari D, Pearson TS, Saffari A, Ziegler A, Kölker S, Volkmann J, Wiesener A, Bearden DR, Lakhani S, Segal D, Udwadia-Hegde A, Martinuzzi A, Hirst J, Perlman S, Takiyama Y, Xiromerisiou G, Vill K, Walker WO, Shukla A, Dubey Gupta R, Dahl N, Aksoy A, Verhelst H, Delgado MR, Kremlikova Pourova R, Sadek AA, Elkhateeb NM, Blumkin L, Brea-Fernández AJ, Dacruz-Álvarez D, Smol T, Ghoumid J, Miguel D, Heine C, Schlump JU, Langen H, Baets J, Bulk S, Darvish H, Bakhtiari

S, Krueger MC, Lim-Melia E, Aydinli N, Alanay Y, El-Rashidy O, Nampoothiri S, Patel C, Beetz C, Bauer P, Yoon G, Guillot M, Miller SP, Bourinaris T, Houlden H, Robelin L, Anheim M, Alamri AS, Mahmoud AAH, Inaloo S, Habibzadeh P, Faghihi MA, Jansen AC, Brock S, Roubertie A, Darras BT, Agrawal PB, Santorelli FM, Gleeson J, Zaki MS, Sheikh SI, Bennett JT, Sahin M. Defining the clinical, molecular and imaging spectrum of adaptor protein complex 4-associated hereditary spastic paraplegia. *Brain*. 2020 Oct 1;143(10):2929-2944. doi: 10.1093/brain/awz307. Erratum in: *Brain*. 2021 Apr 12;144(3):e33. doi: 10.1093/brain/awaa424. PMID: 32979048; PMCID: PMC7780481.

(11) Ebrahimi-Fakhari D, Behne R, Davies AK, Hirst J. AP-4-Associated Hereditary Spastic Paraplegia. 2018 Dec 13. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2024. PMID: 30543385.

(12) Hatsugai N, Nakatsuji A, Unten O, Ogasawara K, Kondo M, Nishimura M, Shimada T, Katagiri F, Hara-Nishimura I. Involvement of Adapter Protein Complex 4 in Hypersensitive Cell Death Induced by Avirulent Bacteria. *Plant Physiol*. 2018 Feb;176(2):1824-1834. doi: 10.1104/pp.17.01610. Epub 2017 Dec 14. PMID: 29242374; PMCID: PMC5813582.

(13) Tüysüz B, Bilguvar K, Koçer N, Yalçınkaya C, Çağlayan O, Gül E, Sahin S, Çomu S, Günel M. Autosomal recessive spastic tetraplegia caused by AP4M1 and AP4B1 gene mutation: expansion of the facial and neuroimaging features. *Am J Med Genet A*. 2014 Jul;164A(7):1677-85. doi: 10.1002/ajmg.a.36514. Epub 2014 Apr 3. PMID: 24700674.

(14) Wang H, Xu Y, Chen M, Shang Q, Sun Y, Zhu D, Wang L, Huang Z, Ma C, Li T, He L, Xing Q, Zhu C. Genetic association study of adaptor protein complex 4 with cerebral palsy in a Han Chinese population. *Mol Biol Rep*. 2013 Nov;40(11):6459-67. doi: 10.1007/s11033-013-2761-6. Epub 2013 Sep 25. PMID: 24065543.

(15) Kong XF, Bousfiha A, Rouissi A, Itan Y, Abhyankar A, Bryant V, Okada S, Ailal F, Bustamante J, Casanova JL, Hirst J, Boisson-Dupuis S. A novel homozygous p.R1105X mutation of the AP4E1 gene in twins with hereditary spastic paraplegia and mycobacterial disease. *PLoS One*. 2013;8(3):e58286. doi: 10.1371/journal.pone.0058286. Epub 2013 Mar 5. PMID: 23472171; PMCID: PMC3589270.

(16) Castro P, Martí M, Oliván-Blázquez B, Boñar N, García V, Gascón-Santos S, Panzano A, Vela S, Tajadura S, Peña A, Tris-Ara MJ. Benefits of robotic gait assistance with ATLAS 2030 in children with cerebral palsy. *Front Pediatr*. 2024 Jul 29;12:1398044. doi: 10.3389/fped.2024.1398044. PMID: 39135857; PMCID: PMC11318455.

(17) İpek R, Kömür M, Direk MÇ, Bozdoğan ST, Okuyaz Ç. Two Sibling Cases of Spastic Paraplegia-45 with a Novel Pathogenic Variant in *NT5C2* Gene: Concomitant *RYR1* Gene in One Sibling. *Mol Syndromol*. 2024 Aug;15(4):297-302. doi: 10.1159/000536183. Epub 2024 Feb 23. PMID: 39119448; PMCID: PMC11305657.

(18) Lai ZH, Liu XY, Song YY, Zhou HY, Zeng LL. Case report: Hereditary spastic paraplegia with a novel homozygous mutation in ZFYVE26. *Frontiers in Neurology*. 2023 Aug 23;14:1160110.