

## Original Research Article

# Influence of Varying Levels of Organic Manure, Phosphorus and Bio-inoculants on Biological Properties of Soil Under Mungbean [*Vigna radiata* (L.) Wilczek]

### ABSTRACT

**Aims:** To varying levels of organic manure, phosphorus and bio-inoculants on biological properties of soil under Mungbean.

**Place and Duration of Study:** Instructional Farm, College of Agriculture, Jodhpur during both the years 2019 and 2020.

**Methodology:** A field experiment was laid out in factorial randomized block design (FRBD) with three replications. The experiment comprised of two levels of FYM (0 and 5 t/ha), four levels of phosphorus (0, 50, 75 and 100 % P<sub>2</sub>O<sub>5</sub> /ha) and three levels of bio-inoculants (No inoculation, *Enterobacter cloacae* and *Bacillus amyloliquefaciens*).

**Results:** Higher soil microbial biomass carbon (217.92 mg/kg), soil microbial biomass phosphorus (15.47 mg/kg), acid phosphatase (1.0919 µg PNP/g soil/h), alkaline phosphatase (7.13µg PNP/g soil/h) and dehydrogenase activity (6.10 µg TPF/g soil/h) recorded under application of FYM 5 t/ha at 40 DAS significantly higher as compared to without FYM application. Among the phosphorus levels, application of 100% P<sub>2</sub>O<sub>5</sub> /ha gave significantly the highest soil microbial biomass carbon (219.13mg/kg), soil microbial biomass phosphorus (15.71 mg/kg), acid phosphatase (1.010 µg PNP/g/soil/h), alkaline phosphatase (7.21µg PNP/g soil/h) and dehydrogenase activity (5.91µg TPF/g soil/h) over rest of P levels, however, it was found at par with 75 % P<sub>2</sub>O<sub>5</sub> /ha. Similarly, the seed inoculation with *B. amyloliquefaciens* recorded significantly higher values of all above biological properties as compared to un-inoculated control but it was found at par with seed inoculation with *E. cloacae*.

**Conclusion:** Application of FYM 5.0 t/ha , phosphorus 100% P<sub>2</sub>O<sub>5</sub> /ha and seed inoculation with biofertilizers (either with *E. cloacae* or *B. amyloliquefaciens* had significant effect in improving the biological properties of soil at 40 DAS of mungbean.

Keywords:FYM; Phosphorus, Bio-inoculants; Biological properties; Mungbean

## 1. INTRODUCTION

Mungbean (*Vigna radiata* (L.) Wilczek) is a leguminous pulse crop cultivated in arid and semi-arid regions of India. At present, the intensive land use in relation with continuous use of high analytical inorganic fertilizers has found to realize enhanced crop production but only at the cost of soil health. Thus a vital research priority is to sustain soil productivity and to supervision changes in physico-chemical and biological soil properties as influenced by management [1]. It is well established fact that application of biofertilizers and organic ameliorates to the soil can definitely influence the soil productivity [2&3] through rising soil properties involving microbial as well as enzymatic activities.

The soils of the Western region of Rajasthan have multiple challenges like desertic nature, coarse texture, slight alkalinity, and poor fertility status[4]. Addressing nutrient deficiency is crucial, necessitating attention to nutrient management strategies involving organic manures, inorganic fertilizers, and biofertilizers[5]. Among essential nutrients, phosphorus holds particular significance for legumes, being regarded as the 'energy currency element' [6]. However, applying phosphorus in problematic soils poses challenges due to its fixation, resulting in limited availability to plants. To enhance phosphorus availability, phosphorus solubilizing bacteria (PSB) like *Pseudomonas* sp. and *Bacillus* sp. play a crucial role in converting insoluble phosphorus into a readily available form in the soil. Seed inoculation with suitable PSB strains is a cost-effective approach that enhances yield by solubilizing unavailable phosphorus into an accessible form, reducing the dependency on costly inorganic phosphorus fertilizers [7]. Evaluating the feasibility and efficiency of organic

manures and biofertilizers is essential not only for improving soil fertility but also for increasing phosphatic fertilizer efficiency. The management of chemical fertilizers, organic manures, and biofertilizers has shown great promise in sustaining crop production stability, soil health, and productivity compared to using each component alone [8]. While chemical fertilizers have raised concerns regarding their impact on microbial activity and soil productivity, the application of organic manures and biofertilizers has been found to boost nutritional status, microbial activity, and soil productive potential. Soil microorganisms, being crucial sources and sinks of mineral nutrients, significantly contribute to soil fertility [8]. Soil organic matter, including living microorganisms, acts as a transitory nutrient reservoir, releasing nutrients from organic matter for plant use. Understanding the microbial processes is essential for managing farming systems, especially those relying on organic nutrient inputs [9].

The current study aimed to assess the effects of various levels of organic manure, phosphorus, and bio-inoculants on biological properties of soil for two years in mungbean cultivation in arid and semi-arid climatic condition. By exploring these aspects, we strive to contribute to the knowledge and understanding of sustainable agricultural practices that promote soil health and crop productivity while considering environmental concerns in the long term.

## 2. MATERIALS AND METHODS

A field experiment was carried out during the *kharif* 2019 and 2020 at Instructional Farm, College of Agriculture, Agriculture University, Jodhpur, Rajasthan. The initial soil properties of experimental field were loamy sand in texture, soil reaction 8.32, having EC 0.89 dS/mat 25 °C and low in organic carbon (1.3 g/kg). The experiment was laid out in factorial randomized block design (FRBD) in three replications. Treatments comprising of two levels of FYM (0 and 5 t/ha), four levels of recommended dose of phosphorus (0, 50, 75 and 100 %  $P_2O_5$  /ha) and three levels of bio-inoculants (no inoculation, *Enterobacter cloacae* and *Bacillus amyloliquefaciens*). Recommended dose of nitrogen and zinc were uniformly applied to all the plots as basal application through urea and zinc sulphate, respectively. The whole quantities of FYM and phosphorus (through single super phosphate) were applied as per treatment prior to sowing and incorporated manually in top 15 cm of soil. As per treatments, seeds of mungbean variety GM-4 were inoculated with PSB strains *i.e.* *Enterobacter cloacae* (PSB-17) and *Bacillus amyloliquefaciens* (PSB-41) before sowing as per the standard method and dried in shade and used for sowing. The fresh soil samples were collected at 0-15 cm depth from experimental field at 40 DAS and stored in refrigerator and were sieved through 2.0 mm sieve. The soil microbial biomass carbon was estimated by adopted fumigation extraction method [10], soil microbial biomass phosphorus was determined by chloroform fumigation-extraction method [11]. The soil enzymatic activities like acid and alkaline phosphatases ( $\mu\text{g PNP/g soil/h}$ ) were analyzed by measuring the amount of p-nitrophenol released as described by [12] and dehydrogenase activity ( $\mu\text{g TPF/g soil/h}$ ) in soil was determined by using the reduction of 2,3,5- triphenyleterazolium chloride [13].

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of FYM on biological properties

#### 3.1.1 Soil microbial biomass

Incorporation of FYM had significant influence on soil microbial biomass carbon (SMBC) and soil microbial biomass phosphorus (SMBP) content in soil at 40 DAS of mungbean (Table 1). Significantly higher SMBC (217.92 mg/kg) and SMBP (15.47 mg/kg) were recorded with the incorporation of FYM @ 5 t/ha over control (198.10 and 14.21 mg/kg) respectively. It might be due to the fact that FYM is a steady source of organic carbon to encourage the soil microbial community [14]. Moreover, build-up of soil microbial biomass is largely due to the contained of microbial biomass in the organic manures, resulting addition of substrate carbon, which encourage the native soil micro-biota [15, 16, 17 & 18].

#### 3.1.2 Soil enzymatic activity

The enzymatic activities were increased significantly with incorporation of FYM. The maximum acid phosphatase activity (1.0919  $\mu\text{g PNP/g soil/h}$ ), alkaline phosphatase activity (7.13  $\mu\text{g PNP/g soil/h}$ ) and dehydrogenase activity (6.10  $\mu\text{g TPF/g soil/h}$ ) at 40 DAS of soil were recorded under the treatment of FYM @ 5 t/ha. Application of FYM @ 5 t/ha increases the

APA, AIPA and DHA of soil to the tune of 6.81, 2.15 and 12.34 per cent, respectively over control. A significant increase in acid phosphatase activity under FYM manured treatments might be due to better growth and number of fine roots of lesser diameter and thus producing exudation of intracellular acid phosphate [19]. This might be due to the fact that FYM is a better source of carbon resultant enhanced microbial activity which secreted more alkaline phosphatase enzymes [19]. Comparable consequences were also reported by Bargaz [20]. Significant improvement in dehydrogenase enzyme activity under FYM treated plots might be due to increase in microbial growth with the addition of carbon substrate [21].

### 3.2 Effect of phosphorus on soil biological properties

#### 3.2.1 Soil microbial biomass

Significantly higher SMBC (219.13 mg/kg) and SMBP (15.71 mg/kg) content was observed under application of super optimal phosphorus level (100 % P<sub>2</sub>O<sub>5</sub> /ha) over control and 50% P<sub>2</sub>O<sub>5</sub>, while it was found statistically at par with 75% P<sub>2</sub>O<sub>5</sub>. Soil microbial biomass carbon augmented by increasing levels of inorganic fertilizers, might be due primarily increase in microbial population [22] besides secondary formation of root exudates, mucigel sloughed off cells and underground roots of previous cut crops which besides play a significant role in increasing biomass carbon [23]. Soil microbial biomass phosphorus increase with increases graded levels of phosphorus. It provided substrates indispensable for microbial growth and their activity, which in turn accountable for increase soil microbial biomass phosphorus [17].

#### 3.2.2 Soil enzymatic activity

The highest activities of all enzymes in soil at 40 DAS were observed under the crop fertilized with 100% P<sub>2</sub>O<sub>5</sub> followed by 75% and 50% P<sub>2</sub>O<sub>5</sub> while, minimum activity was noticed in control. The maximum values of APA, AIPA and DHA were 1.010, 7.21 µg PNP/g soil/h and 5.91 µg TPF/g soil/h under the 100% P<sub>2</sub>O<sub>5</sub> application, respectively on pooled mean basis. However, 75 and 100% P<sub>2</sub>O<sub>5</sub> were being at par with each other. The increase in the level of inorganic P fertilizers has led to an increase in the activity of dehydrogenase as well as phosphatase. It might be ascribed to the fact that inorganic sources of nutrient encouraged the enzyme activity of soil microorganisms which utilize the native pool of organic carbon as a source of carbon [18]. Comparable findings were also reported by [17].

### 3.3 Effect of biofertilizers on biological properties

#### 3.3.1 Soil microbial biomass

Significantly higher SMBC (218.87 mg/kg) and SMBP (15.78 mg/kg) content was observed in *B. amyloliquefaciens* inoculated plots as compared to un-inoculated plots and it was found at par with seed inoculation by *E. cloacae*. Increase in biomass carbon as a result of the secretion of cellulolytic or lignolytic enzymes which in turn might have improved the SMBC. Higher microbial activities obviously make the microbes to transform small fraction of available P into microbial biomass phosphorus [24].

**Table 1. Effect of varying levels of FYM, phosphorus and bio- inoculants on soil microbial biomass at 40 DAS**

Treatments	Soil microbial biomass carbon (mg/kg)			Soil microbial biomass phosphorus (mg/kg)		
	2019	2020	Pooled	2019	2020	Pooled
<b>FYM Levels</b>						
0 t/ha	197.61	198.59	198.10	14.19	14.22	14.21
5 t/ha	214.35	221.50	217.92	15.16	15.78	15.47
SEm±	1.30	1.39	0.90	0.11	0.12	0.08
CD (P=.05)	3.70	3.94	2.53	0.30	0.34	0.22
<b>Phosphorus levels (Recommended dose of P<sub>2</sub>O<sub>5</sub>)</b>						
0% P <sub>2</sub> O <sub>5</sub> /ha	188.24	189.17	188.70	13.17	13.31	13.24
50% P <sub>2</sub> O <sub>5</sub> /ha	205.45	208.84	207.14	14.63	14.92	14.78
75% P <sub>2</sub> O <sub>5</sub> /ha	214.34	219.81	217.07	15.40	15.86	15.63
100% P <sub>2</sub> O <sub>5</sub> /ha	215.88	222.38	219.13	15.50	15.91	15.71
SEm±	1.84	1.96	1.27	0.15	0.17	0.11
CD (P=.05)	5.23	5.58	3.58	0.43	0.48	0.30
<b>Biofertilizers</b>						

Un-inoculated control	185.98	189.03	187.50	13.04	13.15	13.09
<i>E. cloacae</i>	215.08	220.25	217.66	15.38	15.90	15.64
<i>B. amyloliquefaciens</i>	216.87	220.86	218.87	15.61	15.95	15.78
SEm±	1.59	1.70	1.10	0.13	0.15	0.09
CD ( $P=.05$ )	4.53	4.83	3.10	0.37	0.42	0.26


UNDER PEER REVIEW

### 3.3.2 Enzymatic activity

It is evident from data presented (Table 2& Fig. 1) indicated that that acid phosphatase activity (APA), alkaline phosphatase activity (AIPA) and dehydrogenase activity (DHA) of soil at 40 DAS were significantly influenced with biofertilizers seed inoculation. Significantly higher acid phosphatase activity (1.041 µg PNP /gsoil /h), alkaline phosphatase activity (7.44 µg PNP/gsoil/h) and dehydrogenase activity (6.11 µg TPF /g soil /h) were observed in *B. amyloliquefaciens* inoculated plots as compared to un-inoculated control, however it was found statistically at par with *E. cloacae* inoculated plot. Acid phosphatase activity increase with seed inoculation with PSB might due to secretion of acid phosphatase from legumes roots into rhizosphere to hydrolyze organic- P [25&26]. Secondly microorganisms, in addition to roots, are another vital source of acid phosphatase activity in rhizosphere [27&28]. Alkaline phosphatase activity increase with seed inoculation with PSB, which might due to producing of alkaline phosphatase phosphate solubilizing bacteria [29]. Alkaline phosphatase activity was found to be better than acid phosphatase activity in the rhizosphere soil due to the slightly alkaline reaction of experimental soil. Significantly increase in dehydrogenase activity through seed inoculation by biofertilizers. It might be due to increases microbial and root activity in the rhizosphere may largely account for higher activity including phosphatase [30]. Similar results were also reported by [24&31].

**Table 2. Effect of varying levels of FYM, phosphorus and bio- inoculants on enzymatic activity of soil**

Treatments	Enzymatic activity of soil at 40 DAS								
	Acid phosphatase activity			Alkaline phosphatase activity			Dehydrogenase activity		
	(µg PNP/g soil/h)			(µg PNP/g soil/h)			(µg TPF/g soil/h)		
	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
<b>FYM Levels</b>									
0 t/ha	0.950	0.958	0.954	6.95	7.01	6.98	5.40	5.45	5.43
5 t/ha	1.000	1.038	1.019	7.00	7.26	7.13	5.98	6.22	6.10
SEm±	0.006	0.007	0.004	0.04	0.05	0.03	0.038	0.040	0.026
CD (P=.05)	0.017	0.019	0.011	NS	0.13	0.08	0.10	0.12	0.07
<b>Phosphorus levels (Recommended dose of P<sub>2</sub>O<sub>5</sub>)</b>									
0% P <sub>2</sub> O <sub>5</sub> /ha	0.944	0.958	0.951	6.79	6.89	6.84	5.49	5.58	5.54
50% P <sub>2</sub> O <sub>5</sub> /ha	0.971	0.990	0.980	6.94	7.08	7.01	5.66	5.78	5.72
75% P <sub>2</sub> O <sub>5</sub> /ha	0.990	1.019	1.005	7.07	7.25	7.16	5.80	5.97	5.88
100% P <sub>2</sub> O <sub>5</sub> /ha	0.994	1.026	1.010	7.10	7.33	7.21	5.81	6.01	5.91
SEm±	0.008	0.009	0.006	0.06	0.07	0.04	0.054	0.057	0.036
CD at 5%(P=.05)	0.024	0.027	0.016	0.17	0.19	0.12	0.15	0.16	0.10
<b>Biofertilizers</b>									
Un-inoculated control	0.882	0.895	0.888	6.31	6.40	6.35	5.10	5.18	5.14
<i>E. cloacae</i>	1.017	1.044	1.030	7.28	7.46	7.37	5.96	6.13	6.04
<i>B. amyloliquefaciens</i>	1.026	1.056	1.041	7.34	7.55	7.44	6.01	6.20	6.11
SEm±	0.007	0.008	0.005	0.05	0.06	0.04	0.047	0.049	0.031
CD (P=.05)	0.021	0.023	0.014	0.15	0.16	0.10	0.13	0.14	0.09



[F<sub>0</sub>: 0 t/ha, F<sub>1</sub>: 5 t/ha; P<sub>0</sub>: 0%P<sub>205</sub>, P<sub>1</sub>: 50%P<sub>205</sub>, P<sub>2</sub>: 75%P<sub>205</sub>, P<sub>3</sub>: 100%P<sub>205</sub>; B<sub>0</sub>: Control, B<sub>1</sub>: *E. cloacae*, B<sub>2</sub>: *B. amyloliquefaciens*]

Fig. 1. Effect of different levels of FYM, phosphorus and bio-inoculants on enzymatic activity of soil at 40 DAS (Pooled basis)

#### 4. CONCLUSIONS

On the basis of above results, it is concluded that application of FYM @ 5.0 t/ha, 75 percent recommended dose of phosphorus and seed inoculation with biofertilizers (either with *E. cloacae* or *B. amyloliquefaciens*) had significantly improved the biological properties of soil at 40 DAS under mungbean.

**DISCLAIMER (ARTIFICIAL INTELLIGENCE)** Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### REFERENCES

1. Larson WE, Walsh LM, Stewart BA, Boelter DH, editors. Soil and water resources: research priorities for the nation. 1981.
2. McInroy JA, Kloepper JW. Studies on indigenous endophytic bacteria of sweet corn and cotton. Molecular ecology of rhizosphere microorganisms: biotechnology and the release of GMOs. 1994 Aug; 25:19-28.
3. Glick BR. The enhancement of plant growth by free-living bacteria. Canadian journal of microbiology. 1995 Feb 1; 41(2):109-17.
4. Kumar M, Kar A, Raina P, Singh SK, Moharana PC, Chauhan JS. . Assessment and Mapping of Available Soil Nutrients using GIS for Nutrient Management in Hot Arid regions of North-Western India. *Journal of the Indian Society of Soil Science*.2021; 69(2):119-132.
5. Mandiwal M, Shukla UN, Yadav VL, Sarita BH. Effect of phosphorus and biofertilizers on plant height and yield of mungbean [*Vigna radiata* (L.) Wilczek]. *International Journal of Chemical Studies*. 2019;7(4):63-65.
6. Tisdale SL, Nelson WL, Beaton JD, Havline, JL. Soil fertility and fertilizers. 5<sup>th</sup> (Eds.); Prentice Hall of India Private Limited, New Delhi, India.1997
7. Parveen S, Sagir KM, and Zaidi A. Effect of rhizospheric microorganisms on growth and yield of green gram. *Indian Journal of Agricultural Science*. 2002;72(7):421-423.
8. Nambiar KKM, Abrol JP. Long-term fertilizer experiments in India-An overview. *Fertilizer News*. 1992; 34:11-26.
9. Smith JL, Paul EA. The Significance of Soil Microbial Biomass Estimations. In: Bollag, J. and Stotzky, J.D.G., Eds., *SOIL Biochemistry*, Dekker, New York. 1990; 357-396.
10. Vance ED, Brookes PC, Jenkinson DS. An extraction method for measuring soil microbial biomass C. *Soil biology and Biochemistry*. 1987;19(6):703-707.
11. Brookes PC, Powlson DS, Jenkinson DS. Measurement of microbial biomass phosphorus in soil. *Soil biology and biochemistry*. 1982;14(4):319-329.
12. Tabatabai MA, Bremner JM. Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil biology and biochemistry*. 1969;1(4):301-307.
13. Casida JRLE, Klein DA, Santoro T. Soil dehydrogenase activity. *Soil science*. 1964; 98(6):371-376.
14. Bhattacharyya P, Chakrabarti K, Chakraborty A. Microbial biomass and enzyme activities in submerged rice soil amended with municipal solid waste compost and decomposed cow manure. *Chemosphere*. 2005; 60(3):310-318.
15. Verma G, Mathur AK. Effect of integrated nutrient management on active pools of soil organic matter under maize-wheat system of a Typic Haplustept. *Journal of the Indian Society of Soil Science*. 2009; 57(3):317-322.
16. Nath DJ, Ozah B, Baruah R, Barooah RC, Borah DK, Gupta M. Soil enzymes and microbial biomass carbon under rice-toria sequence as influenced by nutrient management. *Journal of the Indian Society of Soil Science*. 2012; 60 (1):20-24.

17. Mohammad I, Yadav BL, Ahamad A. Effect of phosphorus and bio-organics on yield and soil fertility status of mungbean [*Vigna radiata* (L.) Wilczek] under semi-arid condition of Rajasthan, India. *International Journal of Current Microbiology and Applied Sciences*.2017; 6(3), 1545-1553.
18. Yadav B, Naga SR, Yadav BL, Sharma S, Meena SK. Effect of different fertility levels and organic manures on biological properties of soil in Pearl millet. *Journal of Pharmacognosy and Phytochemistry*.2020;9 (1): 210-213
19. Jadhav AB, Suradkar R, Taggu B, Tamboli BD, Priyanka B. Effect of Phytase and FYM on Soil Enzyme Activities, Microbial Population and Nutrient Availability of Non-Calcareous Soil. *Journal of the Indian Society of Soil Science*. 2017; 65(2):222-229.
20. Bargaz A, Faghire M, Abdi N, Farissi M, Sifi B, Drevon JJ, Ghoulam C. Low soil phosphorus availability increases acid phosphatases activities and affects P partitioning in nodules, seeds and rhizosphere of *Phaseolus vulgaris*. *Agriculture*. 2012; 2(2): 139-153.
21. Manna MC, Kundu S, Singh M, Takkar PN. Influence of FYM on dynamics of microbial biomass and its turnover and activity of enzyme under a soybean-wheat system on a Typic Haplusterts. *Journal of the Indian Society of Soil Scienc*.1996; 44: 409-412.
22. Hasebe A, Kanaza WS, Takai Y. Microbial Biomass in Paddy Soil: II. "Microbial Biomass Carbon" Measured by Jenkinson's Fumigation Method. *Soil science and plant nutrition*. 1985;31(3):349-359.
23. Goyal S, Mishra MM, Hoda IS, Singh R, Beri V, Choudhary MR, Sandhu PS, Pasricha NS, Bajwa MS. Build-up of microbial biomass with continuous use of inorganic fertilizers and organic amendents. *Proceeding of the International Symposium on Nutrient Management for Sustained Productivity*. 1992;2: 149-151.
24. Sreelakshmi MM, Aparna, B. Effect of phosphorus solubilizers on enzymatic activity and microbial parameters in the soil. *Int. J. Curr. Microbiol. Appl. Sci*. 2019; 8(8):2647-2649.
25. Yadav R, Tarafdar J. Influence of organic and inorganic phosphorus supply on the maximum secretion of acid phosphatase by plants. *Biology and Fertility of soils*. 2001; 34:140-143.
26. George TS, Gregory PJ, Robinson JS, Buresh RJ. Changes in phosphorus concentrations and pH in the rhizosphere of some agroforestry and crop species. *Plant and Soil*. 2002 Sep; 246:65-73.
27. Marschner H, editor. *Marschner's mineral nutrition of higher plants*. Academic press; 2011 Aug 8.
28. Wei X, Ge T, Zhu Z, Hu Y, Liu S, Li Y, et al. Expansion of rice enzymatic rhizosphere: temporal dynamics in response to phosphorus and cellulose application. *Plant Soil*. 2018;445: 169-181.
29. Chen Q, Liu S. Identification and characterization of the phosphate-solubilizing bacterium *Pantoea* sp. S32 in reclamation soil in Shanxi, China. *Frontiers in microbiology*. 2019 Sep 19; 10:2171.
30. Tarafdar JC, Chhonkar PK. Status of phosphatases in the root-soil interface of leguminous and non-leguminous crops. *Zeitschrift für Pflanzenernährung und Bodenkunde*.1978; 141(3), 347-351.
31. Sahay R, Patra DD. Identification and performance of sodicity tolerant phosphate solubilizing bacterial isolates on *Ocimum basilicum* in sodic soil. *Ecological Engineering*. 2014; 71: 639-643.