

Original Research Article

COMPARISON OF THE MICROBIAL AND SENSORY ATTRIBUTES OF YOGHURT PREPARED FROM TIGER NUT (*Cyperus esculentus L.*) AND DIFFERENT FORMS OF COW MILK

Abstract

Yoghurt made from cow milk is popularly consumed because of its appealing taste and nutritious benefits. However, concerns about animal-based proteins call for action. This study evaluates the microbial and sensory attributes of yoghurt produced from tiger nut (*Cyperus esculentus L.*) milk and different forms of cow milk. Yoghurt was produced from tigernut milk and different forms of cow milk namely: full cream, fat-filled, and skim milk. By fermenting the milk samples at 42 °C for 8 h using a starter culture. A brand of commercial yoghurt was used as the control. The samples were evaluated for physicochemical, microbial using morphological and biochemical techniques and sensory attributes using multistage selection with 9-point hedonic scale. The pH value significantly ranged from 3.97-4.65 for tiger nut and skim milk yoghurts, respectively. While a reverse occurred with titratable acid ranging from 0.088-0.095 %. For both total plate and lactic acid bacteria count, control sample had the significantly lowest values of 1.361×10^5 and 1.063×10^5 CFU/ml, while full cream milk had the highest values of 2.123×10^5 and 1.853×10^5 CFU/ml, respectively. The isolated microorganisms were *Lactobacillus acidophilus*, *Lactobacillus bulgaricus* and *Streptococcus thermophilus*. Overall acceptability ranged from 6.88-7.63 for skim and full-cream yoghurt, respectively. While tiger nut yoghurt scored 7.23.

Therefore, tiger nut milk is a suitable plant-based alternative to cow milk for sustainable yoghurt production.

Keywords: Yoghurt, plant-based, milk, probiotics, microbial, sensory.

Introduction

The growing global population interested in healthy food options have led to a significant push to enhance food variety and nutrition. Meeting these interests, through animal sources alone is far-fetched, raising the need for sustainable plant-based alternatives. Also, plant-based diets have fewer calories required to address the challenges associated with high blood pressure, obesity, lactose intolerance and ethics of vegetarian diets [1,2].

Yoghurt is a fermented milk beverage listed as one of the healthiest food products, resulting mainly from its nutritional composition and probiotic quality [3,4]. Probiotics are living microorganisms that when consumed have positive effects on the host such as improved digestibility, enhancement of microflora, and immune stability [5,3].

Cow milk is known for its high protein content including a complex mix of fats, minerals and vitamins [6,7]. Fullcream milk still retains its natural fat content. Skim milk is produced by completely removing milkfat from fullcream milk [8]. While, fat-filled milk is formulated by blending skim milk with non-dairy fats such as vegetable oils [9]. Numerous industries presently utilize fat-filled milk in yoghurt production as a cost-effective alternative instead of fullcream. However, the affordability of plant-based milk tends to be more promising [10].

Tiger nut (*Cyperus esculentus* L.) an underutilized nut, is chosen for its high levels of essential nutrients, health benefits, affordability and availability [2,11,4]. Its milk content is high. However,

not many studies have compared tiger nut milk, especially with fat-filled milk for yoghurt. Therefore, this research aimed to compare the microbial and sensory attributes of yoghurt produced from tiger nut and different forms of cow milk.

Materials and methods

The samples used for this research were Dano milk powder (full cream, skim and fat-filled), fresh tiger nut tubers, freeze-dried starter culture (yogourmet) and a commercially known yoghurt to serve as the control. All were obtained from a local market in Enugu state and transported to the laboratory for analysis.

Sample preparation

Preparation of tiger nut milk

The method employed had slight modification [11]. Fresh tiger nuts were sorted, washed and drained. Water (2 L) was added to 500g of tiger nut, placed in a water bath at 45 °C for 12 hours, to enhance softening, milling and milk extraction. The hydrated nuts were drained and milled using a blender at 1800 rpm for 5 min. After that, distilled water (1L) was added to form tiger nut mush. Then, strain through a cheesecloth, and the filtrate (tiger nut milk) was pasteurised for 15 min at 75 °C [2]. The milk was cooled to 45 °C for inoculation and fermentation.

Preparation of cow milk

Full cream powdered cow milk (600 g) was dissolved in 1.5 L of distilled water and stirred properly. It was pasteurized at 85 °C for 5 min and cooled to 45 °C for inoculation, followed by fermentation. This same procedure was used for skim and fat-filled milk powders [12].

Preparation of yoghurt with the different forms of cow milk

Each sample (1 L) of tiger nut, full cream, skim and fat-filled milk was immediately inoculated with 5g of a freeze-dried starter culture(yogourmet), in separate beakers. After inoculation, samples were tightly covered and placed in an incubator at 42 °C for 8 h to ferment. The fermented samples (Yoghurt) were refrigerated at 4 °C for further analysis [13].

Physiochemical attributes of the yoghurt samples

Determination of pH values

The pH values of each sample were measured using a pH meter. The pH meter was standardized by testing the buffer solutions of known pH. This was aimed at testing the level of acidity of the yoghurt samples [14].

Determination of Titratable Acidity (TA)

Each yoghurt sample (1g) was mixed with 10 ml of hot distilled water (90 °C) and titrated to a faint colour with 0.1N NaOH comprising 0.5% phenolphthalein as an indicator. The percentage of lactic acid produced by fermentation in the sample was determined as follows [14].

$$\text{Titre value} \times 0.09 \times 100 \%$$

(Where the Titre value is the Volume of yoghurt sample solution used; 0.09 is a conversion factor).

Microbial and biochemical determination

Determination of microbial count

Homogenized yoghurt samples (1 ml) each were aseptically transferred into a corresponding sterile test tube containing 9 ml of distilled water up to a four-fold serial dilution. Using the pour plate method, dilutions 2 and 4 were cultured on MRS and Nutrient Agar, the plates were incubated at 37°C for 24h and the colonies were counted for bacteria load. All counts were expressed as CFU/ml. Pure cultures were obtained by sub-culturing on sterile fresh MRS and nutrient agar plates for Lactic acid bacteria count and Total coliform count, respectively. The plates were incubated at 37°C for 24h. The obtained pure culture was stored on agar slants and refrigerated at 4°C. Isolates were identified using morphological, biochemical and gram-staining tests [15,12].

Catalase test

The catalase test is used to distinguish microorganisms that can produce catalase enzymes. A clear, grease-free slide was treated with 3% hydrogen peroxide (H_2O_2), and a small amount of each bacterial isolate was placed on the glass slide using a sterile inoculating loop, allowing for the isolate's bubbles to develop. Bubbles show catalase positive, while the absence means catalase negative [16].

Oxidase test

The ability of organisms to produce cytochrome oxidase enzymes is employed in oxidase tests. The oxidase reagent was freshly prepared into a 1% solution, and filter paper strips were immersed. The culture was scratched with the inoculating wire loop. Positive reactions are indicated by a vivid, deep-purple hue that appears within 5–10 seconds, while adverse reactions are indicated by a lack of colour [17].

Coagulase test

The coagulase test is utilized for differentiating pathogenic from non-pathogenic test organisms through their ability to coagulate blood plasma. Two distinct grease-free slides each received a few drops of saline, and a loop of the bacterial isolates was emulsified on slides to create two suspensions. A sterile Pasteur pipette was used to collect a drop of human plasma, which was then gently mixed on the slides. The glass was checked for clumping after 5–10 minutes. The presence of clumping indicated coagulase-positive, while the absence showed coagulase-negative[14].

Gram staining

This test was employed to test the organisms microscopically and differentiate Gram-negative from Gram-positive using coloured stains. Smear was created, placed on a spotless glass slide, and stained for 30 seconds with crystal violet. The smear was then cleaned with distilled water. Gram's iodine was applied for 10 seconds, after which the smear was washed with tap water, decoloured with 95% acetone alcohol, and dyed with safranin for 30 seconds. The smear was then rinsed with tap water, dried by air, and examined with a 100X oil immersion objective [14].

Determination of Sensory attributes of the yoghurt samples

The yoghurt samples were coded and evaluated for sensory attributes by 100 individuals. Multistage selection was applied, such that various age groups, genders, occupations, and social strata were randomly chosen among students and staff of the university. A glass of water was given to each panellist so they could rinse their mouths after tasting each sample. They were also

given questionnaires to score the yoghurt samples for appearance, colour, aroma, taste, consistency and overall acceptability, using a 9-point hedonic scale ranging from 0 (extremely dislike) to 9 (extremely like) [18].

Statistical Analysis

All analyses were performed in triplicate and presented as mean \pm standard deviation. Statistical Analysis of Variance (ANOVA) using SPSS version 28 (SPSS, Inc., USA) was applied for means variation while, the Duncan Multiple Range Test (DMRT) at an acceptable level of $p \leq 0.05$ was utilised for the separation of means [19].

Results and Discussion

Microbial evaluation

The physiochemical properties and microbiological loads of the yoghurt samples are presented in Table 1.

Physicochemical attributes measured in this study are pH and titratable acid values, required to denote the acidity and possible shelf stability of the yoghurt.

Table 1: Physiochemical and microbial count of yoghurt samples made from tiger nut milk and different forms of cow milk.

SAMPLE	pH	TITRATABLE ACID (%)	TOTAL COUNT (CFU/ml)	PLATE LAB COUNT (CFU/ml)
FC	4.31 \pm 0.076 ^b	0.090 \pm 0.002 ^a	2.123 \times 10 ⁵ \pm 0.035 ^d	1.853 \times 10 ⁵ \pm 0.025 ^d
SK	4.65 \pm 0.035 ^d	0.088 \pm 0.001 ^a	1.960 \times 10 ⁵ \pm 0.010 ^c	1.570 \times 10 ⁵ \pm 0.026 ^b
FL	4.36 \pm 0.030 ^b	0.092 \pm 0.003 ^a	1.747 \times 10 ⁵ \pm 0.015 ^b	1.092 \times 10 ⁵ \pm 0.028 ^a
TN	3.97 \pm 0.017 ^a	0.095 \pm 0.010 ^b	1.750 \times 10 ⁵ \pm 0.030 ^b	1.630 \times 10 ⁵ \pm 0.026 ^c

CN	4.53±0.035 ^c	0.091±0.002 ^a	1.361 x10 ⁵ ±0.029 ^a	1.063 x10 ⁵ ±0.047 ^a
----	-------------------------	--------------------------	--	--

Means in the same column with same superscript letters are not significantly different at $p \leq 0.05$.

FC- Full cream milk yoghurt; Sk – Skim milk yoghurt; FL – Fat-filled milk yoghurt; TN – Tiger nut milk Yoghurt; CN – Commercial Yoghurt (Control), LAB – Lactic Acid Bacteria.

The microbial attributes evaluated are Total Plate Count (TPC) and Lactic Acid Bacterial (LAB) count of the yoghurt samples. This indicates the general active culture and expected probiotic loads in each sample. pH indicates the extent of acidity or alkalinity of a medium. The pH values differed significantly ($P \leq 0.05$). Tiger nut milk yoghurt exhibited the significantly lowest acidic value of 3.97, while skim milk yoghurt had the highest of 4.65.

Titrateable acidity (TA) measures acid(s) quantity in a medium. The titrateable acid levels in the samples did not differ significantly, except for tiger nut milk yoghurt, which had the highest significant value of 0.95 %.

Microbiological loads, as shown in Table 1, reveal a total microbial count ranging from 1.36×10^5 to 2.13×10^5 CFU/ml, while lactic acid bacteria (LAB) ranged from 1.063×10^5 to 1.85×10^5 CFU/ml.

Table 2: Morphological and biochemical features of yoghurt samples made from tiger nut milk and different forms of milk.

Biochemical	LAB 1	LAB 2	LAB 3
Feature			
Shape	Coci	Rod	Rod
Cell Setting	Single/ short	Circular	Paired/ long


Texture	Dry	Moist	Wet
Colour	Creamy	Off-white	Whitish
Elevation	Flat	Irregular	Raised
Appearance	Opaque	Opaque	Shinny
Catalase	-	-	-
Oxidase	-	-	-
Coagulase	+	+	+
Gram staining	+	+	+
Probable Organism	<i>Lactobacillus acidophilus</i>	<i>Lactobacillus bulgaricus</i>	<i>Streptococcus thermophilus</i>

After determining the LAB count in the samples, it became necessary to identify these bacteria further to indicate the probable organisms present in the samples. Microbial occurrences are indigenous to fermented foods, with Lactic Acids Bacteria being more dominant in yoghurts. *Lactobacillus acidophilus*, *Lactobacillus bulgaricus* and *Streptococcus thermophilus*, were identified as the probable organisms after morphological and biochemical tests were conducted in all the yoghurt samples.

Sensory evaluation

The findings of sensory attributes including colour, aroma, taste, consistency and overall acceptability of the yoghurt samples are shown in Figure 1. The result exhibits that the sensory values of all measured parameters ranged from 6.07-8.22, 6.23-7.72, 5.72-6.50, 6.91-7.74 and 6.88-7.63 for colour, aroma, taste, consistency and overall acceptability, respectively. The

colour values for tiger nut milk and skim milk ranged from 6.07 (lowest) to 8.22 (highest). The highest average aroma for the panelists was scored for full cream milk (7.72), followed by tiger nut milk (7.54) yoghurt. While the least score was observed for skim milk (6.23) yoghurt. For this study, it was shown that the least taste scores of 5.72 of fat-filled milk yoghurt was observed as compared to full-cream milk being the highest (6.50), followed by control (6.36), skim milk (6.31) and tiger nut milk (6.21) yoghurt. The control sample with 7.74 value showed the highest consistency. Next was the full cream sample of 7.4, then tiger nut of 7.12, while skim milk was the least (6.47). The observation indicates that the panelist preferred the full cream milk yoghurt (7.63) when compared to others, followed by the control sample (7.54), while skimmed milk (6.88) yoghurt was the least accepted. The familiarity of panelists with full cream milk used as a standard for yoghurt production may account for the observed preference. However, tiger nut milk yoghurt showed a high score of 7.23 and was the third most accepted yoghurt sample.



FC- Full cream milk yoghurt; Sk – Skim milk yoghurt; FL – Fat-Filled milk yoghurt; TN – Tiger nut Yoghurt;

CN – Commercial Yoghurt (Control)

Figure 1: Sensory evaluation of yoghurt samples made from tiger nut milk and forms of cow milk.

Discussion

The yoghurt samples examined in this study are within the recommended pH range of 4-4.50. However, it is essential to note that the pH of the skim milk sample differs slightly from this range, measuring at 4.65. The pH values of this study were found to be comparable to previous report [21]. This similarity is desirable because it indicates that the pH level below 3.5 may be sufficient to effectively prevent the proliferation of undesirable microorganisms, thereby extending its shelf life [11].

A progressive rise in titratable acid levels and a corresponding decrease in pH values during the fermentation process is expected [21]. It is recommended that TA levels fall into 0.85 - 0.95 % range, which agrees with the values obtained in this study (0.88 – 0.95 %). Similar results were obtained in studies by [22,11].

Quantification of LAB is critical in the yoghurt manufacturing process, necessitating the evaluation of LAB as a critical factor in determining the overall microbial composition of yoghurt products. The full cream milk yoghurt had the highest significant count in both cases, while the control sample showed the lowest. The control sample may contain chemical preservatives to regulate the microbial load and extend the shelf life of the yoghurt [21]. The

observed microbial counts in this study indicate a significant level of microbial load in the fermented yoghurts, which is thought to be a positive characteristic of the presence of active probiotic organisms [2]. The values found in this study are slightly lower than a previous study on yoghurt derived from public school sources [22]. This could have resulted from the operating procedures since the yoghurts in this study were produced in a laboratory adhering to strict hygiene control measures.

Microbial occurrences are indigenous to fermented foods, with Lactic Acids Bacteria being more dominant in yoghurts. This is because as yoghurt fermentation progresses pH decreases and inhibits the growth of many microorganisms, while LAB dominates (1). This finding is similar to that in the production of yoghurt, tiger nut milk can serve as good substitute for cow milk, if probiotic composition is the main interest [23].

The sensory values show to what extent the yoghurt samples are accepted by consumers and the possible sensory features that call for improvement [1].

Colour is a required attribute for measuring dairy and non-dairy products of milk because white or creamy colour are universally seen as milk colour. Consequently, this unconsciously affects to what extent milk or its related products are accepted. Similar colour values were obtained from tiger nut and coconut milk, ranged from 6.2-6.8 [24]. The lower tiger nut milk score from this study may be due to the brownish pigments in whole tiger nuts, which were leached during milk extraction.

The aroma finding supports previous study for comparing tiger nut milk yoghurt with fresh cow milk and other milk forms. The observation could be attributed to the diverse array of volatile aromatic chemicals and acetaldehyde produced by microbes present in tiger nut yoghurt during

the breakdown of carbohydrates enhancing its aroma [21]. While the absence of fat in the skim milk could have been a limiting factor to its aroma values.

The taste of a product determines to what extent a product appeals to the sweetness or sourness of the yoghurt. Lower taste scores (4.90-5.95) were observed [11]. This result could be so because, animal protein sources have improved sensory perception of mouthfeel when consuming products like Yoghurt. Of which fat-filled and tiger nut milk yoghurt consist of plant-based fat.

Consistency displays the smoothness, thickness as well as mouthfeel of yoghurt samples. This statistic that the fat content in milk products enhances its mouthfeel and emulsifying features [11]. Therefore, skimmed milk with no fat content exhibited the least consistency by panelist scores, although it is perceived to be healthier, many consumers show preference for sensory over nutritional attributes.

Overall acceptability reveals to what extent consumers like or dislike a product on a general note putting all tested parameters into consideration [21]. Therefore, tiger nut stands a chance of being a major source for yoghurt on the shelves also taking into account its health benefits amongst other factors such as affordability and availability [2].

Conclusion

Yoghurt made from tiger nut milk and different forms of cow milk exhibited comparable microbial and sensory attributes. Yoghurt made from full-cream milk showed the highest microbial count and sensory attributes. Moreover, tiger nut milk yoghurt showed both high sensory and acceptable microbial values. Therefore, findings from this study indicate tiger nut milk as a suitable plant-based alternative to cow milk for the production of yoghurt. It is

recommended that other plant based milk be looked into as alternatives to making cow milk yoghurt. This will result to sustainable varieties options.

References

- [1] Dhakal D, Younas T, Bhusal RP, Devkota L, Henry CJ, Dhital S. Design rules of plant-based yoghurt-mimic: Formulation, functionality, sensory profile and nutritional value. *Food Hydrocoll* [Internet]. 2023;142:108786. Available from: <https://www.sciencedirect.com/science/article/pii/S0268005X23003326>
- [2] Satir G. The effects of fermentation with water kefir grains on two varieties of tigernut (*Cyperus esculentus* L.) milk. *LWT* [Internet]. 2022;171:114164. Available from: <https://www.sciencedirect.com/science/article/pii/S0023643822010994>
- [3] Joel N, James S, Offia-Olua B. Development and comparative evaluation of storage changes in probiotic soy-yoghurt.
- [4] Ojuko OO, Tadese AA, Nwabu OA. Physico-chemical, sensory attributes and microbial analysis of Yoghurt made from Powder Milk and Tiger-nut milk blend. *World Sci News*. 2018;(108):111–22.
- [5] Meybodi NM, Mortazavian AM, Arab M, Nematollahi A. Probiotic viability in yoghurt: A review of influential factors. *Int Dairy J*. 2020;109:104793.
- [6] Nielsen SD, Jakobsen LMA, Geiker NRW, Bertram HC. Chemically acidified, live and heat-inactivated fermented dairy yoghurt show distinct bioactive peptides, free amino acids and small compounds profiles. *Food Chem*. 2022;376:131919.
- [7] Kielbasa A, Monedeiro F, Bernatowicz-Łojko U, Sinkiewicz-Darol E, Buszewski B, Gadzała-Kopciuch R. The content of selected omega-3 free fatty acids in breast milk samples and their interactions with macronutrients as well as personal characteristics

- of mothers. *Int Dairy J* [Internet]. 2024;148:105803. Available from: <https://www.sciencedirect.com/science/article/pii/S0958694623002224>
- [8] Hafiizha A, Kayaputri IL, Tensiska T, Amalia NR. The effect of skim milk concentration on sensory quality and PH of probiotic yoghurt added with red dragon fruit (*Hylocereus polyrhizus*). *Jurnal Ilmu dan Teknologi Hasil Ternak (JITEK)*. 2020;15(1):52–60.
- [9] Schmidmeier C, O’Gorman C, Drapala KP, Waldron DS, O’Mahony JA. Elucidation of factors responsible for formation of white flecks in reconstituted fat filled milk powders. *Colloids Surf A Physicochem Eng Asp* [Internet]. 2019;575:245–55. Available from: <https://www.sciencedirect.com/science/article/pii/S0927775719302298>
- [10] Assocle MS, Okidi L, Ongeng D. The effect of rosemary, ginger, or garlic on microbial shelf life and sensory acceptability of nutritionally enriched cassava-based pancake (Kabalagala). *Applied Food Research* [Internet]. 2024;4(1):100384. Available from: <https://www.sciencedirect.com/science/article/pii/S277250222300121X>
- [11] Odejobi OJ, Olawoye B, Ogundipe OR. Modelling and optimisation of yoghurt production from tigernut (*Cyperus esculentus* L.) Using Response Surface Methodology (RSM). *Asian Food Science Journal*. 2018;4(3):1–12.
- [12] Oladipo IC, Atolagbe OO, Adetiba TM. Nutritional evaluation and microbiological analysis of yoghurt produced from full cream milk, tiger-nut milk, skimmed milk and fresh cow milk. *Pensee*. 2014;76(4):30–8.
- [13] Ezeonu CS, Tatah VS, Nwokwu CD, Jackson SM. Quantification of physicochemical components in yoghurts from coconut, tiger nut and fresh cow milk. *Advances in Biotechnology and Microbiology*. 2016;1(5):555573.
- [14] Taiwo OS, Afolabi RO, Oranusi SU, Owolabi JB, Oloyede AR, Isibor PO, et al. Microbiological Assessment of Commercial Yogurt Sold in Ota Metropolis, Ogun State, Nigeria. In: *IOP Conference Series: Earth and Environmental Science*. IOP Publishing; 2018. p. 012019.

- [15] De S, Pramanik A, Das AKR, Paul S, Jana S, Pramanik P. Isolation and characterization of *Lactobacillus* spp. from curd and its pharmacological application in probiotic chocolate. *The Journal of Phytopharmacology*. 2017;6(6):335–9.
- [16] Amanullah SM, Kabir MA, Rahman MM, Halder P, Hossain SMJ, Samad MA. Isolation, identification and biochemical characterization of lactic acid bacteria from selected yogurt samples. *Bangladesh Journal of Livestock Research*. 2020;64–72.
- [17] Ngene AC, Onwuakor CE, Aguiyi JC, Ifeanyi VO, Ohaegbu CG, Okwuchukwu CP, et al. Screening of some lactic acid bacteria isolated from selected Nigerian fermented foods for vitamin production. *Adv Microbiol*. 2019;9(11):943–55.
- [18] Chinma CE, Adedeji OE, Etim II, Aniaka GI, Mathew EO, Ekeh UB, et al. Physicochemical, nutritional, and sensory properties of chips produced from germinated African yam bean (*Sphenostylisstenocarpa*). *LWT*. 2021;136:110330.
- [19] Ojukwu M, Ofoedu C, Seow EK, Easa AM. Optimization of soy protein isolate, microbial transglutaminase and glucono- δ -lactone in gluten-free rice noodles. *J Sci Food Agric*. 2021;101(9):3732–41.
- [20] Makut MD, Olokun AL, Olokun RM. Production of yoghurt from milk extract of tigernut (*Cyperus esculentus*) using lactic acid bacteria isolated from locally fermented milk (nono). *Asian Food Science Journal*. 2018;4(1):1–8.
- [21] Maduka N, Ire FS. Tigernut plant and useful application of tigernut tubers (*Cyperus esculentus*)—A review. *Curr J Appl Sci Technol*. 2018;29(3):1–23.
- [22] Bristone C, Badau MH, Igwebuike JU, Igwegbe AO. Production and evaluation of yoghurt from mixtures of cow milk, milk extract from soybean and tiger nut. *World Journal of Dairy & Food Sciences*. 2015;10(2):159–69.
- [23] Chimezie GD, Gloria L, Ebere I. Microbiological load of yoghurt sold in Omoku schools, Rivers State, Nigeria. *Afr J Microbiol Res*. 2015;9(34):1960–3.

[24]Anumba N, Joy E-E, Ishiwu C, Wabali V. Physicochemical Properties and Sensory Evaluation of a Plant-Based Yoghurt Produced from Blends of Tiger Nut, Coconut and Dates. World Journal of Food Science and Technology [Internet]. 2023. Available from: <https://doi.org/10.11648/j.wjfst.20230703.12>

UNDER PEER REVIEW