

Evaluating Species Diversity and Ecological Health of NnamdiAzikiwe University Forest, Awka

ABSTRACT

The increasing threats to forest habitats due to natural and anthropogenic factors necessitate a detailed ecological assessment. This study aims to understand the ecological dynamics and biodiversity within the forest ecosystems of NnamdiAzikiwe University, Awka, in the context of climate change and habitat destruction. Employing a stratified random sampling method and the Point Center Quarter Method, we collected data from 24 designated sample points across a 5000 m² plot. Measurements included species distance from sampling points and diameter at breast height (DBH) of trees. We computed phytosociological parameters such as density, basal area, dominance, cover, and frequency, along with relative values to determine species importance and diversity indices. A total of 96 trees comprising 71 species from 33 families were identified. Notably, Malvaceae and Fabaceae were the most represented families. Ficusbenjamina had the highest total basal area (113.89 m²/ha) and important value index (25.98), indicating its ecological significance. The Shannon-Wiener index value of 4.13 reflects robust biodiversity, highlighting the ecological richness of the forest. These findings are crucial for informing conservation efforts and sustainable management of forest resources.

Keywords: Ecological assessment, Species diversity, Forest habitat, Point Center Quarter Method (PCQM), Biodiversity

1.0 INTRODUCTION

Throughout human history, trees have been revered for their numerous contributions, ranging from serving as a means of sustenance in the form of fruits and nuts to providing shelter and raw materials for tools and construction [1]. The ancient civilizations of Nigeria recognized the role of trees in maintaining life and culture, a practice that continues to resonate today. Trees were not only sources of sustenance for the ancient civilizations but also symbols of spiritual significance, embodying beauty and life [2]. Forests are among the most important ecosystems on our planet, playing an essential role in preserving global ecological balance. They provide habitats for numerous species, regulate climate, purify air and water, and provide many resources that support both natural and human societies. A forest is a vibrant and intricate ecosystem, an exquisite web of life in which trees and their associated plants and animals form an interconnected community. Within this ecosystem, a remarkable life cycle unfolds as plants and animals interact and eventually experience the complete range of existence, including aging and death [1], [3].

Forests can be found in all places capable of supporting tree growth, at altitudes up to the tree line, except when natural fire frequency or other disturbances are too great, or where the environment has been altered by human activity [4]. Nigeria possesses land spanning 92.4 million hectares, with approximately 9.7 million hectares, constituting around 10% of the nation, designated as forest reserves [5]. The tropical rainforest, a significant component of Nigeria's forested areas, has been recognized as the most biologically diverse terrestrial ecosystem globally [6], [1], [7], [8], [9].

In forest ecology, species diversity is an important metric. Forests may have a large number of tree species in a small area (as in tropical rain and temperate deciduous forests) or a small number of species in a large area (as in taiga and arid montane coniferous forests). Tree diversity is critical to tropical forest biodiversity because trees provide homes and resources for a diverse range of plant and animal species. They contribute significantly to forest ecosystem stability, stress resistance, ecological processes (pollination, reproduction and renewal, competition and dependence, growth, and death), and ecosystem services (primary productivity, decomposition, nutrition, energy, and culture) [10], [11], [12]. As a result, they influence the design and composition of forest communities [13]. This study was carried out to determine the floristic composition and diversity of the study area and to understand how this assessment can lead to biodiversity conservation and sustainable forest management.

2.0 MATERIALS AND METHODS

2.1 Description of the study area

The study site is located at NnamdiAzikiwe University Awka forest, in Anambra State, Nigeria established in the Southeastern zone in 1991 with a mean elevation of 136 meters above sea level. The University lies within the geographical location: 6.245° and 6.283° N, 7.115° and 7.1219° E. Anambra State, with a total land area of 4,416 sq. km is situated on a generally low elevation on the eastern side of the River Niger, and shares boundaries with Kogi, Enugu, Imo, Abia, Rivers, Delta and Edo states. It lies within the following geographical locations: 5° 45' N to 6° 45' N and 6° 36' E to 7° 08' E [14]. It is bordered in the West by Delta State, on the North by Kogi State, on the East by Enugu State, and the South by Imo State. Anambra State has a high potential for agricultural development, because of stretches of fertile land on the plains in Ogbaru, Ayamelum, Oyi, Awka, and Orumba Local Government Areas. Anambra State

experiences an equatorial tropical rainforest climate, marked by two primary seasons namely: the rainy (wet) season and the dry season. The rainy season, featuring intense thunderstorms, spans from April to October, while the dry season covers the period from November to March each year [15]. The rainfall is typically intense throughout the rainy season, except for a notable decrease in August, known as the August break. This contributes to the distinctive double maxima of rainfall in this pattern. The rainy season is marked by elevated temperatures ranging from 25°C to 33°C and high relative humidity at 85% [15].

2.2 Data Collection

The initial step in conducting this research involved conducting a preliminary survey of the forest intended for sampling. A plot size of 5000 m² (50 m × 100 m), representing the minimal area, was employed. Once the plots were delineated, pegs were placed at each end to ensure proper demarcation. The forest's species composition was evaluated through floristic assessment, and this was complemented by determining the abundance of each species present at the site. However, the forest area was delineated and divided into random strata. Measurements of tree girth were then taken for trees taller than one meter. Species identification was conducted using Flora of West Tropical Africa [16] and Nigerian Trees [17].

2.2.1 Stratified Sampling

This process involves dividing the study area into relatively uniform sections and then sampling each subsection based on its area or other relevant parameter. The plotless method was utilized to estimate the species' density. This approach is also applicable for gathering data on species composition (inventory), growth, and environmental factors. The specific plotless method employed in this study is the Point Center Quarter Method. In the Point Center Quarter Method, four distances instead of one were measured at every sampling point. To establish four quarters at the sampling point, a cross was formed by two lines, one following the compass direction and the other running perpendicular to the compass direction through the sampling point. Alternatively, the cross could be randomly determined by spinning it over each sampling point. The distance to the midpoint of the nearest tree from the sampling point was measured in each quarter. The four distances between several sampling points were averaged and squared to find the mean area occupied by each tree. Cottam and Curtis [18] validated the reliability of this method across various random populations by cross-verifying the results with the plot method. The calculations for the accurate mean area per tree (MA) were observed to be applicable across various sets of mean distances. Hence, there is no necessity for a correction factor when averaging the distances from the four quarters; $MA = D^2$, where D represents the mean distance from four points to the nearest tree, measured in each of the four quarters. The mathematical validation of the effectiveness of this method was provided by Morisita (1954). According to Cottam and Curtis [18], accuracy improves with an increasing number of sampling points, and a minimum of 20 points was recommended. Newsome and Dix [19] pointed out that a limitation of this method for field application is that an individual must be located within each quarter, and the same individual must not be measured twice. Therefore, stands with individuals widely spaced pose a challenge in applying this method.

Following the sampling process, the species diversity was calculated using the data obtained from the forest sampling. The Shannon-Wiener Index of Diversity was employed to analyze and ascertain the species diversity of the sampled site utilizing the formula:

$$H' = - \sum_{i=1}^s (P_i) \times (\ln P_i)$$

$$H_{Max} = \ln s$$

$$E, \text{Equitability} = H' / H_{Max}$$

Where; Σ = summation

S = number of tree species

i-I = individual species to one

P_i = proportion of individual species

Ln P_i = natural log of the proportion of the individual species

2.3 Data Analysis

The density, frequency, dominance, and important value index (IVI) and their relative values were calculated for each tree species using the following formula:

Density = Total number of individuals of a species / Total area sampled

Relative density = (Density of a species / Sum of densities of all species) x 100

Relative Frequency = (Frequency of a species / Sum of frequencies of all species) x 100

Dominance = Total basal area of a species / Total area sampled

Relative Dominance = (Dominance of a species / Sum of dominance of all species) x 100

Important value index (IVI) = Relative frequency + Relative density + Relative dominance

3.0 RESULTS

3.1 Ecological Survey of plants in the study area

3.1.1 Forest tree species, composition, diversity, and distribution

A total of 24 sample points numbered A-X were made available using PCQM to identify trees in the study area. The distance of the species closest to the center point per quarter was estimated and the diameter at breast Height (DBH) of the tress was measured.

Table 1: Species Encountered in the sampled Points

Sample Area	Quarter	Species	DBA (cm)	Distance (m)
A	1 st	<i>Combretumillarii</i> Engl.	4.00	2.50
	2 nd	<i>Magnolia grandiflora</i> L.	3.00	1.00
	3 rd	<i>Antiaristoxicaria</i> Lesch	2.50	3.00
	4 th	<i>Ficustrichopoda</i> Baker	3.50	2.00
B	1 st	<i>Elaeiguineensis</i> Jacq	2.50	1.70
	2 nd	<i>Flacourtiarukam</i> Zoll. &Moritzi	3.00	2.90
	3 rd	<i>Ficusbenjamina</i> L.	10.20	2.50
	4 th	<i>Cecropiaobtusa</i> Trecul	5.00	2.50
C	1 st	<i>Trichiliadregeana</i> Sond.	1.10	3.22
	2 nd	<i>Theobroma cacao</i> L.	2.90	2.80
	3 rd	<i>Schizolobiumparahyba</i> (Vell.) S. F. Blake	0.30	3.48
	4 th	<i>Dubosciamacrocarya</i> Bocq.	2.00	3.40
D	1 st	<i>Garcinia gardneriana</i> (Planch. &Triana) Zapp	2.30	2.15
	2 nd	<i>Hymeneacourbarill</i> L.	0.35	4.50
	3 rd	<i>Cola cordifolia</i> (Cav.) R. Br.	0.30	2.30
	4 th	<i>Trichiliadregeana</i> Sond.	2.00	3.40
E	1 st	<i>Nuxiafloribunda</i> Benth.	0.45	2.60
	2 nd	<i>Alchorneaglandulosa</i> Poepp.	0.50	2.40
	3 rd	<i>Petiveraalliacea</i> L.	0.30	2.28
	4 th	<i>Alchorneacordifolia</i> Müll.Arg.	0.30	1.70
F	1 st	<i>Pterocarpusofficinalis</i> Jacq.	1.20	3.28
	2 nd	<i>Bixa Orellana</i> L.	1.50	2.58
	3 rd	<i>Heveabrasiliensis</i> (Willd.ex A. Juss.) Müll.Arg.	0.70	4.00
	4 th	<i>Firmianasimplex</i> (L.) W.Wight	1.00	2.10
G	1 st	<i>Populus alba</i> L.	2.00	5.00
	2 nd	<i>Coffealiberica</i> Hiern	1.39	3.69
	3 rd	<i>Heveabrasiliensis</i> (Willd. ex A. Juss.) Müll.Arg.	0.95	2.38
	4 th	<i>Derris elliptica</i> (Wall.) Benth.	2.23	1.08
H	1 st	<i>Ficusinsipida</i> Willd.	2.47	6.05
	2 nd	<i>Antiaristoxicaria</i> Lesch.	1.95	3.40
	3 rd	<i>Couropitaguianensis</i> Aubl.	0.60	2.14
	4 th	<i>Pterocarpusofficinalis</i> Jacq.	1.56	1.50
I	1 st	<i>Sterculiamacrophylla</i> Vent.	0.80	0.19
	2 nd	<i>Sterculiatragacantha</i> Lindl.	1.04	2.50
	3 rd	<i>Pterocarpusrohrii</i> Vahl	2.15	3.30
	4 th	<i>Toxicodendron succedaneum</i> (L.) Kuntze	2.20	2.10
J	1 st	<i>Mustangacecropioides</i> R. Br. ex Tedlie	3.05	6.50
	2 nd	<i>Annona purpurea</i> Moc. &Sessé ex Dunal	2.30	3.20
	3 rd	<i>Archidendronjiringa</i> (Jack) I. C. Nielsen	1.07	4.00
	4 th	<i>Pterocarpusofficinalis</i> Jacq.	1.25	2.21
K	1 st	<i>Grewiatrihocarpa</i> Hochst. ex A. Rich.	2.34	4.08
	2 nd	<i>Zanthoxylumrhoifolium</i> Lam.	2.17	1.49
	3 rd	<i>Guareaguidonia</i> (L.) Sleumer	1.96	2.72
	4 th	<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight &Arn.	3.31	2.50
L	1 st	<i>Dimocarpuslongan</i> Lour.	1.26	5.02
	2 nd	<i>Cecropiaobtusifolia</i> Bertol.	2.01	4.20
	3 rd	<i>Aphanamixispolystachya</i> (Wall.) R. Parker	1.45	0.93

M	4 th	<i>Elaeisqueensis</i> Jacq.	0.81	2.70
	1 st	<i>Viburnumtinus</i> L.	2.61	2.07
	2 nd	<i>Tetramelesnudiflora</i> R.Br.	1.90	1.25
	3 rd	<i>Huracrepitans</i> L.	2.15	0.79
N	4 th	<i>Horsfieldia kingie</i> (Hook.f.) Warb.	1.52	2.42
	1 st	<i>Stixissuaveolens</i> (Roxb.) Pierre	1.70	4.51
	2 nd	<i>Antiaristoxicaria</i> Lesch.	2.05	3.02
	3 rd	<i>Camellia sinensis</i> (L.) Kuntze	1.89	3.15
O	4 th	<i>Persea Americana</i> Mill.	2.23	2.90
	1 st	<i>Jasminummultiflorum</i> (Burm.f.) Andrews	0.81	2.37
	2 nd	<i>Annona squamosa</i> L.	0.69	4.10
	3 rd	<i>Combretumerythrophyllum</i> Sond.	2.30	2.25
P	4 th	<i>Psidiumcattleianum</i> Sabine	1.59	1.50
	1 st	<i>Tabernaemontanapachysiphon</i> Stapf	2.12	0.62
	2 nd	<i>Vepristrichocarpa</i> (Engl.) Letouzey	0.96	2.12
	3 rd	<i>Elaeisqueensis</i> Jacq.	0.82	3.32
Q	4 th	<i>Ficussycomorus</i> L.	2.33	1.82
	1 st	<i>Millettiapachycarpa</i> Benth.	2.02	2.48
	2 nd	<i>Barringtonia racemose</i> (L.) Spreng.	2.10	2.32
	3 rd	<i>Antiaristoxicaria</i> Lesch.	2.50	4.10
R	4 th	<i>Lanneacoromandelic</i> (Houtt.) Merr.	1.35	0.82
	1 st	<i>Aquilariasinensis</i> (Lour.) Spreng.	2.45	1.49
	2 nd	<i>Dracontomelonduperreanum</i> Pierre	1.83	3.26
	3 rd	<i>Antiaristoxicaria</i> Lesch.	2.40	2.52
S	4 th	<i>Firmianasimplex</i> (L.) W.Wight	0.60	4.80
	1 st	<i>Dipterocarpusturbinatus</i> C.F.Gaertn.	0.62	3.20
	2 nd	<i>Zanthoxylumzanthoxyloides</i> (Lam.) Zepern. & Timler	1.89	2.90
	3 rd	<i>Harpullia pendula</i> Planch. Ex F. Muell.	2.72	4.30
T	4 th	<i>Pterocarpus officinalis</i> Jacq.	0.32	2.45
	1 st	<i>Theobroma cacao</i> L.	0.71	3.63
	2 nd	<i>Cecropiapeltata</i> L.	1.25	2.10
	3 rd	<i>Zanthoxylumzanthoxyloides</i> (Lam.) Zepern. & Timler	1.41	1.96
U	4 th	<i>Saracaindica</i> L.	1.52	2.85
	1 st	<i>Persea americana</i> Mill.	1.32	1.76
	2 nd	<i>Ceibapentadra</i> (L.) Gaertn.	0.92	2.45
	3 rd	<i>Pisonia aculeate</i> L.	1.52	2.70
V	4 th	<i>Sterculiatragacantha</i> Lindl.	2.00	3.10
	1 st	<i>Combretumillarii</i> Browicz & Zemanek	1.27	3.50
	2 nd	<i>Cola millenii</i> K. Schum.	0.42	4.01
	3 rd	<i>Oroxylum indicum</i> (L.) Kurz	0.56	3.60
W	4 th	<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn.	2.52	3.70
	1 st	<i>Ficus insipida</i> Willd.	0.92	2.10
	2 nd	<i>Schizolobium parahyba</i> (Vell.) S. F. Blake	0.67	1.97
	3 rd	<i>Hevea brasiliensis</i> (Willd. ex A. Juss.) Müll. Arg.	1.52	3.26
X	4 th	<i>Zanthoxylum rhoifolium</i> Lam	1.28	3.00
	1 st	<i>Antiaristoxicaria</i> Lesch.	0.86	1.57
	2 nd	<i>Hymenaeacourbaril</i> L.	1.65	2.80
	3 rd	<i>Tabernaemontanadonnell - smithii</i> Rose ex J. D. Sm.	2.24	1.65
	4 th	<i>Elaeisqueensis</i> Jacq.	1.25	2.40
TOTAL				

The species mostly encountered belonged to the Malvaceae family (9 species), followed by the Fabaceae family (8 species), Moraceae family (5 species), Euphorbiaceae family (4 species), Urticaceae family (4 species), Combretaceae, Meliaceae, Rutaceae, and Anarcadiaceae (3 species each), Annonaceae, Lecythidaceae, Sapindaceae, Stilbaceae, Apocynaceae (2 species each), and Arecaceae, Theaceae, Thymelaceae, Bixaceae, Lauraceae, Rubiaceae,

Dipterocarpaceae, Salicaceae, Clusiaceae, Myristicaceae, Oleaceae, Magnoliaceae, Bignoniaceae, Phytoloccaceae, Nyctaginaceae, Myrtaceae, Resedaceae, Tetramelaceae, Adoxaceae (each having one species).
When the number of trees belonging to each family was estimated, Fabaceae was the dominant species in the forest with 13 trees, closely followed by Malvaceae having 12 trees, and Moraceae with 11 trees (Table 2).

Table 2: Family list of the forest, their number of species, and number of trees

Family Name	Scientific Names	No. of Species	Number of Trees
Moraceae	<i>Antiaristoxicaria</i>	5	11
	<i>Ficusinsipida</i>		
	<i>Ficusbenjamina</i>		
	<i>Ficussycomorus</i>		
	<i>Ficustrichopoda</i>		
Arecaceae	<i>Elaeiguineensis</i>	1	4
Fabaceae	<i>Pterocarpusofficinalis</i>	8	13
	<i>Hymenaeacourbaril</i>		
	<i>Schizolobiumparahyba</i>		
	<i>Archidendronjiringa</i>		
	<i>Millettiapachycarpa</i>		
	<i>Pterocarpusrohrrii</i>		
	<i>Saracaindica</i>		
	<i>Derris elliptica</i>		
	<i>Heveabrasiliensis</i>		
Euphorbiaceae	<i>Alchorneacordifolia</i>	4	6
	<i>Alchorneaglandulosa</i>		
	<i>Huracrepitans</i>		
Combretaceae	<i>Combretumillarii</i>	3	5
	<i>Terminalia arjuna</i>		
	<i>Combretumerythrophyllum</i>		
Malvaceae	<i>Sterculiatragacantha</i>	9	12
	<i>Firmiana simplex</i>		
	<i>Theobroma cacao</i>		
	<i>Ceibapentadra</i>		
	<i>Cola cordifolia</i>		
	<i>Cola millenii</i>		
	<i>Dubosciamacrocarpa</i>		
	<i>Grewiatrichocarpa</i>		
	<i>Sterculiamacrophylla</i>		
Lauraceae	<i>Persea Americana</i>	1	2
Meliaceae	<i>Trichiliadregeana</i>	3	4
	<i>Aphanamixispolystachya</i>		
	<i>GuareaGuidonia</i>		
Rutaceae	<i>Zanthoxylumrhoifolium</i>	3	5
	<i>Zanthoxylumzanthoxyloides</i>		
	<i>Vepristrichocarpa</i>		
Annonaceae	<i>Annona purpurea</i>	2	2
	<i>Annona squamosal</i>		
Thymelaeaceae	<i>Aquilariasinensis</i>	1	1
Lecythidaceae	<i>Barringtoniaracemose</i>	2	2
	<i>Couroupitaguianensis</i>		
Bixaceae	<i>Bixa Orellana</i>	1	1
Theaceae	<i>Camellia sinensis</i>	1	1
Urticaceae	<i>Cecropiaobtuse</i>	4	4
	<i>Cecropiaobtusifolia</i>		
	<i>Cecropiapeltata</i>		
	<i>Mustangacecropioides</i>		
Rubiaceae	<i>Coffealiberica</i>	1	1
Sapindaceae	<i>Dimocarpuslongan</i>	2	2
	<i>Harpullia pendula</i>		

Dipterocarpaceae	<i>Dipterocarpustrabinatus</i>	1	1
Anacardiaceae	<i>Dracontomelonduperreanum</i>	3	3
	<i>Lanneacoromandelica</i>		
	<i>Toxicodendron succedaneum</i>		
Salicaceae	<i>Flacourtiarukam</i>	1	1
Clusiaceae	<i>Garcinia gardneriana</i>	1	1
Myristicaceae	<i>Horsifieldia kingie</i>	1	1
Oleaceae	<i>Jasminummultiflorum</i>	1	1
Magnoliaceae	<i>Magnolia grandiflora</i>	1	1
Stilbaceae	<i>Nuxia floribunda</i>	2	2
Bignoniaceae	<i>Oroxylumindicum</i>	1	1
Phytolaccaceae	<i>Petiveraalliacea</i>	1	1
Nyctaginaceae	<i>Pisonia aculeate</i>	1	1
Myrtaceae	<i>Psidiumcattleianum</i>	1	1
Resedaceae	<i>Stixissuaveolens</i>		
Apocynaceae	<i>Tabernaemontanadonnell – smithii</i>	2	2
	<i>Tabernaemontanapachysiphon</i>		
Tetramelaceae	<i>Tetramelesnudiflora</i>	1	1
Adoxaceae	<i>Vibrurnumtinus</i>	1	1

3.2 Floristic, Species Important Value Indices, and Structural Characteristics of the Forest

The result in Table 3 shows that *Antiaristoxicaria* has the highest frequency (6) succeeded by *Elaeisqueensis*, and *Pterocarpus officinalis* (4 each). In terms of dominance, *Ficusbenjamina* had the highest value (113.89 m²/ha), followed by *Cecropiaobtusa* (27.37), and *Antiaristoxicaria* (23.94). *Ficusbenjamina* also recorded the highest important value index (25.98), followed by *Antiaristoxicaria* (17.53), *Elaeisqueensis* (10.45), and *Pterocarpus officinalis* (9.69).

Table 3: Species abundance of the forest

SPECIES	FREQ	R. F (%)	Density	R.Density (%)	Dominance	R. Dom (%)	IVI
<i>Antiaristoxicaria</i>	6	6.25	0.063	6.25	23.94	5.03	17.53
<i>Elaeisqueensis</i>	4	4.17	0.042	4.20	9.91	2.08	10.45
<i>Pterocarpusofficinalis</i>	4	4.17	0.042	4.20	6.31	1.32	9.69
<i>Heveabrasiliensis</i>	3	3.13	0.031	3.10	4.13	0.87	7.10
<i>Combretumillarii</i>	2	2.08	0.021	2.10	19.26	4.04	8.22
<i>Ficusinsipida</i>	2	2.08	0.021	2.10	7.71	1.62	5.80
<i>Firmiana simplex</i>	2	2.08	0.021	2.10	1.65	0.35	4.53
<i>Hymenaeacourbaril</i>	2	2.08	0.021	2.10	3.90	0.69	4.87
<i>Perseaamericana</i>	2	2.08	0.021	2.10	7.43	1.56	5.74
<i>Schizolobiumparahyba</i>	2	2.08	0.021	2.10	0.83	0.17	4.35
<i>Sterculiatragacantha</i>	2	2.08	0.021	2.10	5.78	1.21	5.39
<i>Terminalia arjuna</i>	2	2.08	0.021	2.10	3.16	3.99	8.17
<i>Theobroma cacao</i>	2	2.08	0.021	2.10	9.91	2.08	6.26
<i>Trichiliadregeana</i>	2	2.08	0.021	2.10	5.78	1.21	5.39
<i>Zanthoxylumrhoifolium</i>	2	2.08	0.021	2.10	6.88	1.44	5.62
<i>Zanthoxylumzanthoxyloides</i>	2	2.08	0.021	2.10	6.05	1.27	5.45
<i>Alchorneacordifolia</i>	1	1.04	0.010	1.04	0.10	0.02	2.10
<i>Alchorneaglandulosa</i>	1	1.04	0.010	1.04	0.27	0.06	2.14
<i>Annona purpurea</i>	1	1.04	0.010	1.04	5.79	1.22	3.30
<i>Annona squamosal</i>	1	1.04	0.010	1.04	0.52	0.11	2.19
<i>Aphanamixispolystachya</i>	1	1.04	0.010	1.04	2.30	0.48	2.56
<i>Aquilariasinensis</i>	1	1.04	0.010	1.04	6.57	1.38	3.46
<i>Archidendronjiringa</i>	1	1.04	0.010	1.04	1.25	0.26	2.34
<i>Barringtonia racemose</i>	1	1.04	0.010	1.04	4.83	1.01	3.09
<i>Bixa Orellana</i>	1	1.04	0.010	1.04	2.46	0.52	2.60
<i>Camellia sinensis</i>	1	1.04	0.010	1.04	3.91	0.82	2.90
<i>Cecropiaobtusa</i>	1	1.04	0.010	1.04	27.37	5.74	7.82
<i>Cecropiaobtusifolia</i>	1	1.04	0.010	1.04	4.24	0.93	3.01
<i>Cecropiapeltata</i>	1	1.04	0.010	1.04	1.71	0.36	2.44
<i>Ceibapentadra</i>	1	1.04	0.010	1.04	0.93	0.19	2.27

<i>Coffealiberica</i>	1	1.04	0.010	1.04	2.12	0.44	2.52
<i>Cola cordifolia</i>	1	1.04	0.010	1.04	0.10	0.02	2.10
<i>Cola millenii</i>	1	1.04	0.010	1.04	0.19	0.04	2.12
<i>Combretumerythrophyllum</i>	1	1.04	0.010	1.04	5.79	1.22	3.30
<i>Couroupitaguianensis</i>	1	1.04	0.010	1.04	0.39	0.08	2.16
<i>Derris elliptica</i>	1	1.04	0.010	1.04	5.44	1.14	3.22
<i>Dimocarpuslongan</i>	1	1.04	0.010	1.04	1.74	0.36	2.44
<i>Dipterocarpustrabinatus</i>	1	1.04	0.010	1.04	0.42	0.09	2.17
<i>Dracontomelonduperreanum</i>	1	1.04	0.010	1.04	3.67	0.77	2.85
<i>Dubosciamacrocarpa</i>	1	1.04	0.010	1.04	4.38	0.92	3.00
<i>Ficusbenjamina</i>	1	1.04	0.010	1.04	113.89	23.90	25.98
<i>Ficussycomorus</i>	1	1.04	0.010	1.04	5.94	1.25	3.33
<i>Ficustrichopoda</i>	1	1.04	0.010	1.04	13.41	2.81	4.89
<i>Flacourtiarukam</i>	1	1.04	0.010	1.04	9.85	2.07	4.15
<i>Garcinia gardneriana</i>	1	1.04	0.010	1.04	5.79	1.22	3.30
<i>Grewiatrihocarpa</i>	1	1.04	0.010	1.04	5.99	1.26	3.34
<i>GuareaGuidonia</i>	1	1.04	0.010	1.04	4.21	0.88	2.96
<i>Harpullia pendula</i>	1	1.04	0.010	1.04	8.10	1.70	3.78
<i>Horsifieldiakingii</i>	1	1.04	0.010	1.04	2.53	0.53	2.61
<i>Huracrepitans</i>	1	1.04	0.010	1.04	5.06	1.06	3.14
<i>Jasminummultiflorum</i>	1	1.04	0.010	1.04	0.72	0.15	2.23
<i>Lanneacoromandelica</i>	1	1.04	0.010	1.04	2.00	0.42	2.50
<i>Magnolia grandiflora</i>	1	1.04	0.010	1.04	9.85	2.07	4.15
<i>Millettiapachycarpa</i>	1	1.04	0.010	1.04	4.47	0.94	3.02
<i>Mustangacecropioides</i>	1	1.04	0.010	1.04	10.18	2.14	4.22
<i>Nuxia floribunda</i>	1	1.04	0.010	1.04	0.22	0.05	2.13
<i>Oroxylumindicum</i>	1	1.04	0.010	1.04	0.34	0.07	2.15
<i>Petiveraalliacea</i>	1	1.04	0.010	1.04	0.10	1.56	3.64
<i>Pisonia aculeate</i>	1	1.04	0.010	1.04	2.53	0.53	2.61
<i>Populus alba</i>	1	1.04	0.010	1.04	4.38	0.92	3.00
<i>Psidiumcattleianum</i>	1	1.04	0.010	1.04	2.77	0.58	2.66
<i>Pterocarpusrohrii</i>	1	1.04	0.010	1.04	5.06	1.32	3.40
<i>Saracaindica</i>	1	1.04	0.010	1.04	2.53	0.53	2.61
<i>Sterculiamacrophylla</i>	1	1.04	0.010	1.04	0.83	0.15	2.23
<i>Stixissuaveolens</i>	1	1.04	0.010	1.04	3.16	0.66	2.74
<i>Tabernaemontanadonnell-smithii</i>	1	1.04	0.010	1.04	5.49	1.15	3.23
<i>Tabernaemontanapachysiphon</i>	1	1.04	0.010	1.04	4.92	1.03	3.11
<i>Tetramelesnudiflora</i>	1	1.04	0.010	1.04	3.95	0.83	2.91
<i>Toxicodendron succedaneum</i>	1	1.04	0.010	1.04	5.30	1.11	3.19
<i>Vepristrichocarpa</i>	1	1.04	0.010	1.04	1.01	0.21	2.29
<i>Vibrurnumtinus</i>	1	1.04	0.010	1.04	7.46	1.57	3.65
TOTAL					476.45		309.1

3.3 Species Diversity Index (Shannon Weiner Index)

The analysis of the species diversity using the Shannon Weiner index of species diversity proved that the forest has a high diversity of 4.12922 approximately 4.13, and an evenness of 0.97. This showed that the forest has a large number of species that are relatively evenly distributed and almost equally abundant (Table 4).

Table 4: Shannon Weiner Index of Species Diversity

S/N	Species	N	N	Pi	ln(Pi)	pi*ln(pi)	-Σ(pi)*ln(pi)
1	<i>Antiaristoxicaria</i>	6	96	0.06250	-2.77259	-0.17329	H ¹ = 4.12922
2	<i>Elaeisguineensis</i>	4	96	0.04167	-3.17805	-0.13242	
3	<i>Pterocarpusofficinalis</i>	4	96	0.04167	-3.17805	-0.13242	
4	<i>Heveabrasiliensis</i>	3	96	0.03125	-3.46574	-0.10830	
5	<i>Combretumillarii</i>	2	96	0.02083	-3.87120	-0.08065	
6	<i>Ficusinsipida</i>	2	96	0.02083	-3.87120	-0.08065	

7	<i>Firmiana simplex</i>	2	96	0.02083	-3.87120	-0.08065
8	<i>Hymenaeacourbaril</i>	2	96	0.02083	-3.87120	-0.08065
9	<i>Persea Americana</i>	2	96	0.02083	-3.87120	-0.08065
10	<i>Schizolobiumparahyba</i>	2	96	0.02083	-3.87120	-0.08065
11	<i>Sterculiatragacantha</i>	2	96	0.02083	-3.87120	-0.08065
12	<i>Terminalia arjuna</i>	2	96	0.02083	-3.87120	-0.08065
13	<i>Theobroma cacao</i>	2	96	0.02083	-3.87120	-0.08065
14	<i>Trichiliadregeana</i>	2	96	0.02083	-3.87120	-0.08065
15	<i>Zanthoxylumrhoifolium</i>	2	96	0.02083	-3.87120	-0.08065
16	<i>Zanthoxylumzanthoxyloides</i>	2	96	0.02083	-3.87120	-0.08065
17	<i>Alchorneacordifolia</i>	1	96	0.01042	-4.56435	-0.04755
18	<i>Alchorneaglandulosa</i>	1	96	0.01042	-4.56435	-0.04755
19	<i>Annona purpurea</i>	1	96	0.01042	-4.56435	-0.04755
20	<i>Annona squamosal</i>	1	96	0.01042	-4.56435	-0.04755
21	<i>Aphanamixispolystachya</i>	1	96	0.01042	-4.56435	-0.04755
22	<i>Aquilariasinensis</i>	1	96	0.01042	-4.56435	-0.04755
23	<i>Archidendronjiringa</i>	1	96	0.01042	-4.56435	-0.04755
24	<i>Barringtoniaracemosa</i>	1	96	0.01042	-4.56435	-0.04755
25	<i>Bixa Orellana</i>	1	96	0.01042	-4.56435	-0.04755
26	<i>Camellia sinensis</i>	1	96	0.01042	-4.56435	-0.04755
27	<i>Cecropiaobtusa</i>	1	96	0.01042	-4.56435	-0.04755
28	<i>Cecropiaobtusifolia</i>	1	96	0.01042	-4.56435	-0.04755
29	<i>Cecropiapeltata</i>	1	96	0.01042	-4.56435	-0.04755
30	<i>Ceibapentadra</i>	1	96	0.01042	-4.56435	-0.04755
31	<i>Coffealiberica</i>	1	96	0.01042	-4.56435	-0.04755
32	<i>Cola cordifolia</i>	1	96	0.01042	-4.56435	-0.04755
33	<i>Cola millenii</i>	1	96	0.01042	-4.56435	-0.04755
34	<i>Combretumerythrophyllum</i>	1	96	0.01042	-4.56435	-0.04755
35	<i>Couroupitaguianensis</i>	1	96	0.01042	-4.56435	-0.04755
36	<i>Derris elliptica</i>	1	96	0.01042	-4.56435	-0.04755
37	<i>Dimocarpuslongan</i>	1	96	0.01042	-4.56435	-0.04755
38	<i>Dipterocarpesturbinatus</i>	1	96	0.01042	-4.56435	-0.04755
39	<i>Dracontomelonduperreanum</i>	1	96	0.01042	-4.56435	-0.04755
40	<i>Dubosciamacrocarpa</i>	1	96	0.01042	-4.56435	-0.04755
41	<i>Ficusbenjamina</i>	1	96	0.01042	-4.56435	-0.04755
42	<i>Ficussycomorus</i>	1	96	0.01042	-4.56435	-0.04755
43	<i>Ficustrichopoda</i>	1	96	0.01042	-4.56435	-0.04755
44	<i>Flacourtiarukam</i>	1	96	0.01042	-4.56435	-0.04755
45	<i>Garcinia gardneriana</i>	1	96	0.01042	-4.56435	-0.04755
46	<i>Grewiatrichocarpa</i>	1	96	0.01042	-4.56435	-0.04755
47	<i>Guarea Guidonia</i>	1	96	0.01042	-4.56435	-0.04755
48	<i>Harpullia pendula</i>	1	96	0.01042	-4.56435	-0.04755
49	<i>Horsifieldia kingie</i>	1	96	0.01042	-4.56435	-0.04755
50	<i>Huracrepitans</i>	1	96	0.01042	-4.56435	-0.04755
51	<i>Jasminummultiflorum</i>	1	96	0.01042	-4.56435	-0.04755
52	<i>Lanneacoromandelica</i>	1	96	0.01042	-4.56435	-0.04755
53	<i>Magnolia grandiflora</i>	1	96	0.01042	-4.56435	-0.04755
54	<i>Millettiapachycarpa</i>	1	96	0.01042	-4.56435	-0.04755
55	<i>Mustangacecropioides</i>	1	96	0.01042	-4.56435	-0.04755
56	<i>Nuxia floribunda</i>	1	96	0.01042	-4.56435	-0.04755
57	<i>Oroxylumindicum</i>	1	96	0.01042	-4.56435	-0.04755
58	<i>Petiveraallicea</i>	1	96	0.01042	-4.56435	-0.04755
59	<i>Pisonia aculeate</i>	1	96	0.01042	-4.56435	-0.04755
60	<i>Populus alba</i>	1	96	0.01042	-4.56435	-0.04755
61	<i>Psidiumcattleianum</i>	1	96	0.01042	-4.56435	-0.04755
62	<i>Pterocarpusrohrii</i>	1	96	0.01042	-4.56435	-0.04755
63	<i>Saracaindica</i>	1	96	0.01042	-4.56435	-0.04755
64	<i>Sterculiamacrophylla</i>	1	96	0.01042	-4.56435	-0.04755
65	<i>Stixissuaveolens</i>	1	96	0.01042	-4.56435	-0.04755

			96			
66	<i>Tabernaemontanadonnell-smithii</i>	1		0.01042	-4.56435	-0.04755
67	<i>Tabernaemontanapachysiphon</i>	1	96	0.01042	-4.56435	-0.04755
68	<i>Tetramelesnudiflora</i>	1	96	0.01042	-4.56435	-0.04755
69	<i>Toxicodendron succedaneum</i>	1	96	0.01042	-4.56435	-0.04755
70	<i>Vepristrichocarpa</i>	1	96	0.01042	-4.56435	-0.04755
71	<i>Vibrurnumtinus</i>	1	96	0.01042	-4.56435	-0.04755
	Total					-4.12922

$$\text{Equitability} = (H^1 / H_{\text{max}})$$

Where H^1 = Shannon Weiner Diversity Index = $-\sum(\pi_i) \cdot \ln(\pi_i) = -(-4.12922) = 4.12922$

$H_{\text{max}} = \ln(71) = 4.2627$

$H^1 / H_{\text{max}} = 4.12922 / 4.2627$

= 0.9686

4.0 DISCUSSION

This study aimed to conduct a comprehensive ecological assessment of the species diversity and evaluate the physicochemical parameters of the soil within the forest of NnamdiAzikiwe University, Awka. A total of 96 trees comprising 71 species and 33 families were documented during the study. The high number of species and families recorded in the study area highlights the rich biodiversity of the studied forest which is consistent with research findings that show that tropical rainforests are the most biologically diverse terrestrial ecosystem in the world [1], [8].

Among the families observed, Malvaceae and Fabaceae were the most dominant species, having 9 and 8 species respectively, closely followed by Moraceae with 5 species, Euphorbiaceae and Urticaceae with 4 species each, and Combretaceae, Meliaceae, and Rutaceae having 3 species each. This observation suggests that these families play a vital role in shaping the forest ecosystem. The presence of many species recorded in these families demonstrates their ecological significance in the forest. The species richness of the Malvaceae family observed in this study agrees with the work of Olaoti-Laaroet *al* [20] in their study of tree species diversity and distribution in the natural forest of Onigambari Forest Reserve, Oyo State, Nigeria. They observed that Malvaceae was the most abundant among other families. However, regarding the number of individual trees within a family, Fabaceae was more dominant with 13 trees followed by Malvaceae which had 12 trees, Moraceae with 11 trees, Euphorbiaceae with 6 trees, and Combretaceae and Rutaceae having 5 trees each. This may be due to their ability to germinate quickly along with the symbiotic properties that have allowed them to establish in habitats rapidly. This observation was consistent with the work of Dekaet *al* [21] on the vegetative assessment of tree species in Takamanda forest, in Cameroon. Patrick *et al* [22] in their work on diversity, distribution, and conservation status of forest tree species in cross river state, Nigeria also reported Fabaceae to be the dominant species, but in addition, Malvaceae, Moraceae, Euphorbiaceae, were also among the families observed to be dominant, which is in agreement with the findings of this work. These families may be dominant in the study area because of their ability to adapt to their habitat and suitable environmental conditions that favour their root penetration and nutrient absorption from the subsurface. This is confirmed by Austin *et al* [23] who discovered that an ecosystem's species richness and establishment can be significantly influenced by edaphic factors such as soil nutrients.

When analyzing the species found in the study area, *Antiaristoxicaria* had the highest frequency of 6, this suggests that *A. toxicaria* is a frequently encountered plant in the study area. This may be due to their ability to grow in diverse habitats, thrive in tropical and subtropical regions across Africa, Asia, and the Pacific Islands, and tolerate a range of soil types and climatic conditions [24]. The large number of seeds produced by *Antiaristoxicaria*, and its rapid growth rate enabling them to reach maturity quickly as reported by Orwaet *al* [25] may be another reason why it was frequently encountered in this study area. Aside from *A. toxicaria*, other species such as *Elaeisqueensis*, and *Pterocarpusofficinalis* with four occurrences each were also recorded. In terms of dominance, *Ficusbenjamina* had the highest value (113.89 m²/ha), followed by *Cecropiaobtusa* (27.37 m²/ha), and *Antiaristoxicaria* (23.94 m²/ha). Other species like *Combretumillarii*, *Elaeisqueensis*, *Theobroma cacao*, *Ficustrichopoda*, and *Mustangacecropioides* had 19.26 m²/ha, 9.91 m²/ha, 9.91 m²/ha, 13.41 m²/ha, and 10.18 m²/ha respectively. These values offer insights into the important ecological functioning and structure of the ecosystem and how these species can play vital roles in preserving ecological balance by providing habitats and foods for the various faunas inhabiting the forest. According to Finegan [26], the structural framework of a forest is frequently formed by dominant tree species, which also create a variety of microhabitats on the forest floor, understory, and canopy. Supporting numerous plant and animal species, including birds, animals, insects, and epiphytes. The dominance pattern observed in this study aligns with previous research works on tropical forests. Condit *et al* [27] in Panama's Barro Colorado Island, demonstrated that a few species exerted more influence on the overall shape and function of the ecosystem. These species with distinct demographic characteristics played a significant role in shaping the

demographic space analyzed, indicating that certain species may have a disproportionate impact on the dynamics of the ecosystem compared to others which is in line with the findings of this research. However, the findings of Adekunle *et al* [28] are in contrast to the observation recorded in this study area. In their work on Forest reserves in Southwestern Nigeria, they observed that different environmental conditions and management regimes have a significant influence on plant species composition and richness, which in turn affects forest growth and yield.

In terms of important value index, *Ficusbenjamina* also recorded the highest value of 25.98, followed by *Antiaristoxicaria*(17.53), *Elaeisqueensis* (10.45), and *Pterocarpusofficinalis* (9.69). The high IVI value of *Ficusbenjamina* indicates that it is a species exerting an influence on the structure of the forest availability of habitats and distribution of resources. Its prominent presence signifies its importance in maintaining balance and fostering biodiversity [28].

The Importance Value Index (IVI) is vital in ecological research as it offers valuable insights into a species' overall significance within a specific ecosystem. IVI is calculated by summing the values of relative frequency, relative density, and relative dominance of species making it an indispensable tool for understanding the ecological roles of species and their impact on forest structure and dynamics.

The species diversity index employed in this research is the Shannon-Weiner Diversity index (H'). Since the Shannon-Weiner diversity index considers both species richness and evenness in a community, it has been widely used by researchers to study ecosystem diversity [28], [29], [30], [31], [32]. Species diversity is typically one of the most significant indices used to assess an ecosystem. An ecosystem with a low value (H') will have low species diversity, whereas an ecosystem with great species diversity and richness has a significant value (H') [33]. This present study has a high species diversity of 4.13, this may be due to the limited or low exploitation of the species in the forest habitat and the species' ability to withstand unfavorable environmental variables that are common in the forest. Patrick *et al*[22] recorded a much higher species diversity in the Okwangwo Division of Cross River National Park, and in the Oban Division of Cross River National Park (4.75 and 4.68 respectively). The diversity indices of this study were higher than those reported for protected rainforests in Nigeria [1], [34], [35], [36].

Bush *et al* [37] and Richlefs and Schluter [38] found that environmental complexity or heterogeneity leads to increased species diversity. Their study found that environmental factors that benefit a specific group of organisms may not benefit another. Therefore, to predict the impact of environmental structure on biodiversity, it is important to understand the ecological requirements of species.

5.0 CONCLUSION

The results from this study reveal that the forest of NnamdiAzikiwe University, Awka has high species diversity. Some families of trees were more dominant than others, and these families include Malvaceae, Fabaceae, and Moraceae. The abundance of these families is probably a result of their ability to cope with prevalent environmental conditions in the study site. The abundance of trees in the study site demonstrated that the vegetation is relatively natural playing an important role in shaping the forest ecosystem. Some of the dominant species found in the study site include: *Antiaristoxicaria*, *Elaeisqueensis*, *Pterocarpusofficinalis*, *Heveabrasiliensis*, *Combretumillarrii*, *Ficusinsipida*, *Firmiana simplex*, *Hymenaeacourbaril*. *Ficusinsipida*, *Firmiana simplex*, while some of the least encountered plant species during the study include: *Alchorneacordifolia*, *Alchorneaglandulosa*, *Annona purpurea*, *Annona squamosal*, *Aphanamixispolystachya*, *Aquilariasinensis*, *Archidendronjiringa*, *Ficusbenjamina*.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that AI technologies such as Natural Language Processors, etc was used to proofread the manuscripts for grammatical correctness.

Details of the AI usage are given below:

- 1.Quillbot grammar checker
- 2.Ref n write
3. Grammarly.com

REFERENCES

1. Onyekwelu J C, Mosandl R &Stimm B. Tree Species Diversity and Soil Status of Primary and Degraded Tropical Rainforest Ecosystems in South-Western Nigeria. *Journal of Tropical Forest Science*. 2008; **20**(3): 193-204.
2. Okereke C N, Mbaekwe E I, Nnabude P C, Ekwealor K U, Nwonumara G N, Iroka C F, &Ukpaka C G. Comparative Evaluation of the Species Richness and Diversity of Three Parallel Forest Ecosystems in South-Eastern Nigeria. *Journal of Agriculture and Ecology Research International*. 2016;**8**(4): 1-12.
3. Ekwealor K U, Iroka C F, Ukpaka G C, Okafor P N, Okeke P N, Okereke K E. An Investigation on the Forest and Savannah Vegetation Types in NnamdiAzikiwe University, Awka Campus in Anambra State of Nigeria. *Natural resources and conservation*. 2020; **8**(1)7-18.
4. Okereke C N, Nnabude P C, Mbaekwe E I, Ekwealor K U, Uhuo C A and Nwanchor K C. The Use of Ecological Methods in Vegetative Studies of Plant Species and Abundance in South-eastern Nigeria. *African Journal of Plant Sciences*. 2014; **8**(9): 441-449.
5. ITTO. Status of Tropical Forest Management. Technical series.2011; **38**: 420.
6. Turner M G. Disturbance and Landscape Dynamics in a Changing World. *Ecology*. 2010; **91**(10): 2833–2849.
7. Schmitt C B, Burgess N D, Coad L, Belokurov A, Besançon C, Boisrobert L, Campbell A, Fish L, Gliddon D, Humphries K, Kapos V, Loucks C, Lysenko I, Miles L, Mills C, Minnemeyer S, Pistorius T, Ravilious C, Steininger M, &Winkel G. Global Analysis of the Protection Status of the World's Forests. *Biological Conservation*, 2009; **142**(10): 2122-2130.
8. FAO. Global Forest Resources Assessment 2010. Rome: FAO. Available online at: <http://www.fao.org/docrep/013/i1757e/i1757e.pdf> (Accessed 10/11/2015). 2010.
9. IUCN. Plants under Pressure – a Global Assessment. The First Report of the IUCN Sampled Red List Index for Plants. Royal Botanic Gardens, Kew, UK, Natural History Museum, London, and International Union for Conservation of Nature. 2010.
10. Wang H X, Hu YB, & Zhao Z H. A simple Method for the Estimation of Tree Species Spatial Diversity Index (TSS). *Journal of Northwest Forestry University*. 2013; **28**(4): 184–187.
11. Brockerhoff E G, Barbaro L, Castagneyrol B, Forrester D I, Gardiner B, González-Olabarria J R, Lyver P O B, Meurisse N, Oxbrough A, Taki H, Thompson I D, Van Der Plas F, Jactel H. Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodiversity Conservation*. 2017;**26**(13):3005–3035.
12. Moradi A, Afsharzadeh S, Hamzehee B, Mozaffarian V. Study of Plant Diversity and Floristics in the Westernmost Hyrcanian Forests. *Journal of Forestry Research*. 2020; **31**(5):1589–1598.
13. Akwaji P I & Edu E A. Population Frequency, Density, Abundance and Diversity of Tree Species in Ten Communal Forests of Northern Cross River State, Nigeria. *International Journal of Current Research*, 2017; **9**(10), 59581- 59596.
14. Egboka B C E. Erosion, Gullies, and Landslide Ravage in Awka, Anambra State. ASECA. 215 pp. 1993.
15. Nwosu M C. Studies on the Physico-Chemical Parameters and Plankton Abundance in Relation to Fish Production of Some Lakes in Anambra State, Nigeria. Ph D. Thesis. 203 pp. 2003.
16. Hutchinson J &Dalzel J M. *Flora of West Tropical Africa. Vol 2. 2nd Edition. Crown Agent Publishers. Milbank, London. 544 pp. 1963.*
17. Keay RWJ, Onochie CFA &Standfield DP. *Nigerian Trees Vol 2. Department of Forestry Research Ibadan. Nigeria. 512 pp. 1964.*
18. Cottam G & Curtis JT. The Use of Distance Measures in Phytosociological Sampling. *Ecology*. 1956; **37**:451-460.
19. Newsome RD & Dix RI. The forest of Cypress Hills. Alberta and Saskatchewan. Canada. *Midland Naturalist*. 1968; **80**: 118-185.
20. Olaoti-laaro S O, Taiwo D M, Adeniji K A, Odewo SA, Animashaun ZT, Woghiren A I, Adedoyin E D. Tree Species Diversity and Distribution in the Natural Forest of Onigambari Forest Reserve, Oyo State, Nigeria. *Journal of Applied Sciences and Environmental Management*. 2021; **25**(5): 829 - 834.
21. Deka J, Tripathi PO, Khan L M. High Dominance of ShorearobustaGaertn in Alluvial Plain Kamrup Sal Forest of Assam, N. E. India. *International Journal of Ecosystem*.2012; **2**(4): 67-73.
22. Patrick AI, Oden G N, Onah D O, Okon E I, Ajikah L B &Akomaye F A. Diversity, distribution, and conservation status of forest tree species in Cross River State, Nigeria. *Sustainability and Biodiversity Conservation*. 2022; **1**(1):42-83.

23. Austin M P, Pausas J G & Nicholls A O. Patterns of tree species richness in relation to environment in South Eastern New South Wales. *Australian Journal of Ecology*. 1996; **21**:154-164.
24. Kokwaro J O. *Medicinal Plants of East Africa*. 2nd Ed. Kenya Literature Bureau, Nairobi. 1993.
25. Orwa C, Mutua A, Kindt R, Jamnadass R & Anthony S. *Agroforestry Database: a tree reference and selection guide version 4.0*. World Agroforestry Centre, Kenya. 2009.
26. Finegan B. Pattern and Process in Neotropical Secondary Rainforests: The first 100 years of succession. *Trends in Ecology & Evolution*. 1996; **11**(3), 119-124.
27. Condit R, Hubbell S P & Foster R B. Assessing the response of plant functional types to climatic change in tropical forests. *Journal of Vegetation Science*. 1996; **7**(3):405-416.
28. Adekunle V A J, Olagoke AO & Akindele SO. Tree species diversity and structure of a Nigerian Strict Nature Reserve. *Tropical Ecology*. 2013; **54**(3): 275-289.
29. Onyekwelu J. Site index curves for site quality assessment of *Naucleadiderrichii* monoculture plantations in Omo Forest Reserve, Nigeria. 17. 2005.
30. Adeduntan S A. and Olusola J A. Variation in the abundance of trees and insect species in selected forest reserves in Ondo state. *Journal of Sustainable Technology*. 2015; **6**:63-75.
31. Oke OS, Akindele SO & Onyekwelu JC. Tree species richness, diversity and structure of a strict conservation natural tropical rainforest ecosystem in Nigeria. *Forests and Forest Products Journal*. 2017; **10**:39-51.
32. Amonum JI, Jonathan BA, Japheth HD. Structure and Diversity of Tree Species at the College of Forestry and Fisheries, University of Agriculture Makurdi, Benue State, Nigeria. *International Journal of Forestry and Horticulture (IJFH)*. 2019; **5**:20-27.
33. Sobuj NA & Rahman M. Assessment of plant diversity in Khadimnagar National Park of Bangladesh. *International Journal of Environmental Science*. 2011; **2**(1): 1-13.
34. Bello AG, Isah AD and Ahmad B. Tree species diversity analysis of Kogo Forest Reserve in North Western Nigeria. *International Journal of Plant, Animal and Environmental Sciences*. 2013; **3**(3): 189-196.
35. Saka MG, Jatau DF & Olaniyi WA. Status of indigenous tree species in Girei forest reserve of Adamawa state. *Journal of Research in Forestry, Wildlife and Environment*. 2013; **5**(1): 28-40.
36. Ikyagba TE, Tee TN, Dagba BI, Ancha UP, Ngibo KD & Tume C. Activities on vegetation of Olokemeji Forest Reserve. *Global Nest: The International Journal*. 2015; **6**(2): 130-139.
37. Bush MD, Pipemo D R & Colivanx PA. A 6000 year history of Amazonian Maize cultivation. *Journal of plant science*. 1989; **23**(6): 123-148.
38. Richlefs R.E & Schluter D. *Species diversity in ecological community of Chicago*. The University of Chicago Press. USA. 75 pp. 1993.